

Changes in Mental Status in Patient with Parkinson's Disease With or Without Psychosis

Mellisa Calt, MD, Liz Joseph, MSIII, Tiffany Battagliola, MSIII, Alexandra De Pastene, MSIII, Matthew O'Reilly, MSIII, Ryan Babayev, MS III, and Adriana Fitzsimmons, MD
Department of Psychiatry, Jersey Shore University Medical Center, Neptune NJ



Hackensack
Meridian Health

LEARNING OBJECTIVES

1. Identify diagnostic conditions in patients with Parkinson's Disease who are admitted with a change in mental status.
2. Discuss treatment approaches



CASE PRESENTATION

- Male patient in his 70s, with 5 year history of Parkinson's Disease treated with levodopa-carbidopa, was admitted medically after falling and injuring his hip and arm.
- Patient has a past psychiatric history of OCD, depression, social anxiety and PTSD, treated with clomipramine, quetiapine, and fluoxetine.
- Patient has a past medical history of lymphoma, HTN, DM, and HLD.
- Prior to admission, his outpatient psychiatrist was focused on managing the patient's OCD and anger issues which had progressed recently.
- At his first outpatient visit, his lamotrigine was increased from 50mg to 100mg daily, which he stopped taking due to a reported adverse effect of hypnagogic hallucinations.
- CT head on arrival was negative for any acute intracranial events.
- It was unclear if the patient's fluoxetine was effective and was subsequently started on clomipramine 75mg PO daily for his OCD symptoms, with a plan to taper off fluoxetine.
- The patient was continued on quetiapine at 150mg PO daily for psychosis.
- While hospitalized, the patient continued his home dose of psychiatric medications, was optimized medically, and his mental status and QTc were monitored.
- Psychiatry was consulted for acute onset of hallucinations and delusions after accidentally being restarted on lamotrigine, an old medication.

PSYCHIATRY CONSULT

- He was AAOX2, calm and cooperative, no hallucinations or delusions of assessment, diminished recent memory, and poor attention.
- He has a history of delirium with hallucinations attributed to either his levodopa-carbidopa or his fluctuating glucose levels.

TRIGGERS AND TREATMENT OPTIONS



Lenka, Abhishek, et al. "Approach to the Management of Psychosis in Parkinson's Disease" *Annals of Movement Disorders*, Annals of Movement Disorders, 4 Dec. 2019, www.aomd.in/article.asp?issn=2590-3446;year=2019;volume=2;issue=3;page=83;epage=90;au=Lenka.

CONCLUSIONS

- Approximately sixty percent of patients with Parkinson's Disease develop psychosis.
- Potential causes of psychosis in this population include infection, dehydration, metabolic abnormalities, and medications. Other triggers include changes in sleep, environment, or nutrition.
- Treatment approaches include general measures such as re-establishment of circadian rhythms and environment, addressing coexisting medical conditions, and reducing the use of anticholinergic, anticholinergic, and sedating drugs.
- Tapering anti-parkinson medications may be considered.
- Clozapine and quetiapine have low D2-receptor affinity and have shown to improve psychotic symptoms without worsening movement symptoms.
- Typical antipsychotics should be avoided as they may worsen motor symptoms.
- In 2006, AAN guidelines identified the need for a new antipsychotic without dopamine antagonist effects. Pimavanserin, an atypical antipsychotic, was developed and approved for the treatment of hallucinations and delusions in PD-associated patients.

SUMMARY

- Patients with Parkinson's Disease can develop psychosis with various risk factors.
- It is important to adequately treat Parkinson's disease psychosis to prevent the worsening of movement symptoms.
- While low D2-receptor affinity drugs have helped to improve conditions without worsening movement symptoms, new medications which target different receptors, Pimavanserin, have also shown to improve psychosis without worsening movement symptoms.

References:

1. Sawada H, Oeda T, Yamamoto K, Umemura A, Tomita S, Hyashi R, Kohsaka M, Kawamura T(2013) Trigger medications and patient-related risk factors for Parkinson disease psychosis requiring anti-psychotic drugs: a retrospective cohort study. *BMC Neurol*13:145. doi:10.1186/1471-2377-13-145
2. Schleisman, Andrew. "Treatment of Parkinson's Disease Psychosis." *U.S. Pharmacist - The Leading Journal in Pharmacy*, 17 Nov. 2016, www.uspharmacist.com/article/treatment-of-parkinsons-disease-psychosis
3. Seppi K, Weintraub D, Coelho M, Perez-Lloret S, Fox SH, Katzenschlager R, Hametner EM, Poewe W, Rascol O, Goetz CG, Sampaio C (2011) The Movement Disorder Society Evidence-Based Medicine Review Update: treatments for the non-motor symptoms of Parkinson's disease. *Mov Disord* 26(Suppl 3):S42-S80. doi:10.1002/mds.23884