

Vector Algebra

EE24Btech11022 - Eshan sharma

I. MCQs WITH ONE CORRECT ANSWER

- 1) The unit vector which is orthogonal to the vector $3\hat{i} + 2\hat{j} + 6\hat{k}$ and is coplanar with vectors $2\hat{i} + \hat{j} + \hat{k}$ and $\hat{i} - \hat{j} + \hat{k}$ is (2004S)
 - (a) $\frac{2\hat{i}-6\hat{j}+\hat{k}}{\sqrt{41}}$
 - (b) $\frac{2\hat{i}-3\hat{j}}{\sqrt{13}}$
 - (c) $\frac{3\hat{i}-\hat{k}}{\sqrt{10}}$
 - (d) $\frac{4\hat{i}+3\hat{j}-3\hat{k}}{\sqrt{34}}$
- 2) A variable plane at a distance of the one unit from the origin cuts the coordinate axes at A, B and C. If the centroid D (x, y, z) of triangle ABC satisfies the relation $\frac{1}{x^2} + \frac{1}{y^2} + \frac{1}{z^2} = k$, then the value k is (2005S)
 - (a) 3
 - (b) 1
 - (c) $\frac{1}{3}$
 - (d) 9
- 3) If $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are three non-zero, non-coplanar vectors and $\mathbf{b}_1 = \mathbf{b} - \frac{\mathbf{b} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a}$, $\mathbf{b}_2 = \mathbf{b} + \frac{\mathbf{b} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a}$, $\mathbf{c}_1 = \mathbf{c} - \frac{\mathbf{c} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a} + \frac{\mathbf{b} \cdot \mathbf{c}}{|\mathbf{c}|^2} \mathbf{b}_1$, $\mathbf{c}_2 = \mathbf{c} - \frac{\mathbf{c} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a} + \frac{\mathbf{b}_1 \cdot \mathbf{c}}{|\mathbf{b}_1|^2} \mathbf{b}_1$, $\mathbf{c}_3 = \mathbf{c} - \frac{\mathbf{c} \cdot \mathbf{a}}{|\mathbf{c}|^2} \mathbf{a} + \frac{\mathbf{b} \cdot \mathbf{c}}{|\mathbf{c}|^2} \mathbf{b}_1$, $\mathbf{c}_4 = \mathbf{c} - \frac{\mathbf{c} \cdot \mathbf{a}}{|\mathbf{c}|^2} \mathbf{a} = \frac{\mathbf{b} \cdot \mathbf{c}}{|\mathbf{b}|^2} \mathbf{b}_1$, then the set of orthogonal vectors is (2005S)
 - (a) $(\mathbf{a}, \mathbf{b}_1, \mathbf{c}_3)$
 - (b) $(\mathbf{a}, \mathbf{b}_1, \mathbf{c}_2)$
 - (c) $(\mathbf{a}, \mathbf{b}_1, \mathbf{c}_1)$
 - (d) $(\mathbf{a}, \mathbf{b}_2, \mathbf{c}_2)$
- 4) A plane which is perpendicular to two planes $2x - 2y + z = 0$ and $x - y + 2z = 4$ passes through $(1, -2, 1)$. The distance of the plane from the point $(1, 2, 2)$ is (2006 -3M,-1)
 - (a) 0
 - (b) 1
 - (c) $\sqrt{2}$
 - (d) $2\sqrt{2}$
- 5) Let $\mathbf{a} = \hat{i} + 2\hat{j} + \hat{k}$, $\mathbf{b} = \hat{i} - \hat{j} + \hat{k}$ and $\mathbf{c} = \hat{i} + \hat{j} - \hat{k}$. A vector in the plane of \mathbf{a} and \mathbf{b} whose projection on \mathbf{c} is $\frac{1}{\sqrt{3}}$, is (2006-3M,-1)
 - (a) $4\hat{i} - \hat{j} + 4\hat{k}$
 - (b) $3\hat{i} + \hat{j} - 3\hat{k}$
 - (c) $2\hat{i} + \hat{j} - 2\hat{k}$
 - (d) $4\hat{i} + \hat{j} - 4\hat{k}$
- 6) The number of distinct real values of λ , for which the vectors $-\lambda^2\hat{i} + \hat{j} + \hat{k}$, $\hat{i} - \lambda^2\hat{j} + \hat{k}$ and $\hat{i} + \hat{j} - \lambda^2\hat{k}$ are coplanar, is (2007 - 3marks)
 - (a) zero
 - (b) one
 - (c) two
 - (d) three
- 7) let $\mathbf{a}, \mathbf{b}, \mathbf{c}$ be unit vectors such that $\mathbf{a} + \mathbf{b} + \mathbf{c} = \mathbf{0}$. Which of the following are correct? (2007-3marks)
 - (a) $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} = \mathbf{c} \times \mathbf{a} = \mathbf{0}$
 - (b) $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} = \mathbf{c} \times \mathbf{a} \neq \mathbf{0}$
 - (c) $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} = \mathbf{a} \times \mathbf{c} \neq \mathbf{0}$
 - (d) $\mathbf{a} \times \mathbf{b}, \mathbf{b} \times \mathbf{c}, \mathbf{c} \times \mathbf{a}$ are mutually perpendicular
- 8) The edges of a parallelopiped are of unit length and are parallel to non-coplanar unit vectors $\hat{a}, \hat{b}, \hat{c}$ such that $\hat{a} \cdot \hat{b} = \hat{b} \cdot \hat{c} = \hat{c} \cdot \hat{a} = \frac{1}{2}$. Then, the volume of the parallelopiped is (2008)
 - (a) $\frac{1}{\sqrt{2}}$
 - (b) $\frac{1}{2\sqrt{2}}$
 - (c) $\frac{\sqrt{3}}{2}$
 - (d) $\frac{1}{\sqrt{3}}$
- 9) Let two non-collinear unit vectors \hat{a} and \hat{b} form an acute angle. A point P moves so that at any time t the position vector \overrightarrow{OP} (where O is the origin) is given by $\hat{a} \cos t + \hat{b} \sin t$. When P is farthest from origin O, let M be the length of \overrightarrow{OP} and \hat{u} be the unit vector along \overrightarrow{OP} . Then, (2008)
 - (a) $\hat{u} = \frac{\hat{a} + \hat{b}}{|\hat{a} + \hat{b}|}$ and $M = (1 + \hat{a} \cdot \hat{b})^{1/2}$
 - (b) $\hat{u} = \frac{\hat{a} - \hat{b}}{|\hat{a} - \hat{b}|}$ and $M = (1 + \hat{a} \cdot \hat{b})^{1/2}$
 - (c) $\hat{u} = \frac{\hat{a} + \hat{b}}{|\hat{a} + \hat{b}|}$ and $M = (1 + 2\hat{a} \cdot \hat{b})^{1/2}$
 - (d) $\hat{u} = \frac{\hat{a} - \hat{b}}{|\hat{a} - \hat{b}|}$ and $M = (1 + 2\hat{a} \cdot \hat{b})^{1/2}$
- 10) Let P(3,2,6) be a point in space and Q be a point on the line $\mathbf{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu(-3\hat{i} + \hat{j} + 5\hat{k})$. Then the value of μ for which the vector \overrightarrow{PQ}

is parallel to the plane $x - 4y + 3z = 1$ is (2009)

- (a) $\frac{1}{4}$
- (b) $-\frac{1}{4}$
- (c) $\frac{1}{8}$
- (d) $-\frac{1}{8}$

11) If $\mathbf{a}, \mathbf{b}, \mathbf{c}$, and \mathbf{d} are unit vectors such that $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = 1$ and $\mathbf{a} \cdot \mathbf{c} = \frac{1}{2}$, then (2009)

- (a) $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are non-coplanar
- (b) $\mathbf{b}, \mathbf{c}, \mathbf{d}$ are non-coplanar
- (c) \mathbf{b}, \mathbf{d} are non-parallel
- (d) \mathbf{a}, \mathbf{d} are parallel and \mathbf{b}, \mathbf{c} are parallel

12) A line with positive direction cosines passes through the point $P(2, -1, 2)$ and makes equal angles with the coordinate axes. The line meets the plane $2x + y + z = 9$ at point Q . The length of the line segment PQ equals (2009)

- (a) 1
- (b) $\sqrt{2}$
- (c) $\sqrt{3}$
- (d) 2

13) Let P, Q, R and S be the points on the plane with position vectors $-2\hat{i} - \hat{j}, 4\hat{i}, 3\hat{i} + 3\hat{j}$ and $-3\hat{i} + 2\hat{j}$ respectively. The quadrilateral $PQRS$ must be a (2010)

- (a) parallelogram, which is neither a rhombus nor a rectangle
- (b) square
- (c) rectangle, but not a square
- (d) rhombus, but not a square

14) Equation of the plane containing the straight line $\frac{x}{2} = \frac{y}{3} = \frac{z}{4}$ and perpendicular to the plane containing the straight lines $\frac{x}{3} = \frac{y}{4} = \frac{z}{2}$ and $\frac{x}{4} = \frac{y}{2} = \frac{z}{3}$ is (2010)

- (a) $x + 2y - 2z = 0$
- (b) $3x + 2y - 2z = 0$
- (c) $x - 2y + z = 0$
- (d) $5x + 2y - 4z = 0$

15) If the distance of the point $P(1, -2, 1)$ from the plane $x + 2y - 2z = \alpha$, where $\alpha > 0$, is 5, then the foot of the perpendicular from P to the plane is (2010)

- (a) $\left(\frac{8}{3}, \frac{4}{3}, -\frac{7}{3}\right)$
- (b) $\left(\frac{4}{3}, -\frac{4}{3}, \frac{1}{3}\right)$
- (c) $\left(\frac{1}{3}, \frac{2}{3}, \frac{10}{3}\right)$
- (d) $\left(\frac{2}{3}, -\frac{1}{3}, \frac{5}{2}\right)$