

Parameter provenance and conditional justification of the reconstructed AE4 model

Technical report for the AE4 project

15 September 2026

Abstract

This report inventories the actual Task 40 production objects, rather than treating dataclass defaults as the model parameter set. Each numerical value, protocol setting and inactive legacy field is classified by its role and provenance. The inventory contains 137 records, of which 109 are marked active; these are not independently fitted physiological parameters. Published kinetic coefficients, transferred model settings, quantities derived from WT constraints, effective assumptions, numerical controls and the separate Task 41 inverse parameter are distinguished. No parameter is refitted, no experimental uncertainty interval is invented, and no historical model or result is changed. Several quantities are mathematically derived but remain conditional on unmeasured assumptions, most notably the NBC capacity constructed from a 70/30 chloride loading partition. The accompanying full system analysis provides local sensitivity evidence, not a substitute for experimental parameter identification.

1 Source of the audit

The audit uses repository <https://github.com/esig626/ae4-salivary-transport-control> at main revision `c4d4f207f702d8eee09ab94d2d90ae90552dd641`. It loads the actual nested Task 40 model through its historical validation helper and checks the complete active parameter hashes against `results/40_ae4_equal_cation_routing/frozen_inputs.json`. It does not overwrite the helper or weaken its historical numerical checks. The canonical initial condition is the separately frozen WT resting state, not the constructor default tuple.

The model lineage includes the original AE4 reconstruction [Vera-Sigüenza et al., 2018], the Palk and Benjamin NKCC1 law [Palk et al., 2010, Benjamin and Johnson, 1997], and the Cha NHE kinetic model [Cha et al., 2009]. The experimental evidence for AE4 transport and regulation constrains mechanism and direction, but does not automatically identify every abundance, gain or timescale in this implementation [Peña Münzenmayer et al., 2015, 2016, 2021, Catalán et al., 2025]. This report preserves that distinction.

2 What counts as a justification

A published numerical coefficient has a source, but transfer from another tissue or model still needs an applicability argument. A value determined by a balance equation is justified conditional on the concentrations, fluxes and structural assumptions used in that balance. A value selected to keep a reference trajectory within physiological limits is an effective modelling choice; acceptance of that trajectory does not make the value measured, unique, or known within an uncertainty interval. A numerical inversion bracket is not a physiological observation. A parameter selected using the knockout phenotype must remain labelled as an inverse design choice.

The inventory records value, units, original metadata, active role, classification, exact code locations, the equation role and source, mathematical justification, default value and whether the active value differs from that default. Separate fields state what constrains each quantity, what does not justify it, its logical domain, its uncertainty status, its relevance to manuscript conclusions, and an experiment that could constrain it. Classification is conservative: inherited metadata identifies a lineage, not a newly verified measurement of salivary transporter abundance. Each physiological uncertainty interval remains explicitly null in the machine readable inventory. Logical positivity or fractional bounds are not substituted for experimental uncertainty.

The current categories include physical constants; published kinetic coefficients; inherited model conversions or settings; reference and WT constraint derived values; WT architecture derived values; effective assumptions; numerical brackets and regularisation; protocol and law selections; inactive legacy values; constructor seeds; and the target selected Task 41 parameter. These categories should not be collapsed into a single “from the literature” column.

3 Parameters that can be explained mathematically

3.1 NKCC1 scale: derived from a modelled resting flux

The production law has the form

$$J_N = \alpha_N m_N(t) \psi([Na]_i, [K]_i, [Cl]_i), \quad \psi = \frac{A_1 - A_2[Na]_i[K]_i[Cl]_i^2}{A_3 + A_4[Na]_i[K]_i[Cl]_i^2}.$$

At the accepted resting state, the scale is determined by

$$\alpha_N = \frac{J_{N,rest}}{\psi(c_{rest})} = 0.017334746894096052 \text{ fmol/s.} \quad (1)$$

Equation (1) makes the choice reproducible and avoids an independent extra fit. However, $J_{N,rest}$ is itself an inherited model flux. The scale is not a directly measured salivary carrier amount. The much larger legacy field `homeostasis.nkcc1_capacity_fmol_s` is inactive for this selected law and must not appear in a publication table as its active capacity.

The four Palk coefficients contain fourth order concentration factors. Their conversion to mM is essential. They are effective coefficients for the adopted fixed bath, so an uncertainty study that changes bath composition cannot automatically reuse them without rechecking the underlying reduction.

3.2 NHE1 carrier amount: a WT resting pH calibration

The active NHE1 law is the eight state Cha reduction with its Mod2 term. Printed rate and binding constants are retained, including the documented rounding discrepancy in the fourth reverse rate. The salivary carrier amount is

$$N_H = 2.339370005697548 \times 10^{-5} \text{ fmol.}$$

This was determined against a resting pH target of 6.91 on the selected R09 background. The derivation fixes one scale conditional on that background and target. It does not identify a cardiac carrier count as a salivary measurement, nor does it prove unique joint identification of the carrier amount, buffer pool and other pH pathways.

The completion audit also records the fixed Mod2 exponents $n_H = 3$ and $m_H = 1$, which were absent from the earlier inventory because they are module constants rather than dataclass fields. Both are part of the selected kinetic law. Their inclusion corrects an inventory omission without changing the model.

Legacy tanh NHE capacity and older NHE kinetic fields remain in the dataclass for other models, but are inactive in production. The active stimulus multiplier is 2.3. Calling it fixed before the test documents absence of retuning; it does not establish that its transfer to every tissue and stimulus is exact. The mathematical companion report shows explicitly how the alkalinity bound depends on this multiplier and N_H .

3.3 NBC capacity: derived conditional on an assumed partition

The NBC source imports one sodium and two bicarbonate equivalents per inward cycle. Its capacity is

$$G_B = 0.11570913197464398 \text{ fmol/s.}$$

The construction begins from the assumed fraction $s = 0.70$ of positive WT chloride loading assigned to NKCC1 at the Task 31 reference scale. If that NKCC1 chloride flux is L_N , the corresponding AE4 demand is

$$J_{4,req} = \frac{1-s}{s} L_N. \quad (2)$$

Using $L_N = 0.2562444047732706 \text{ fmol/s}$ and the selected stimulated NHE flux, the approximate reference requirement is $B_{req} = J_{4,req} - H_{stim}/2$, before evaluating the full coupled electrical closure. The capacity is then chosen so the NBC law supplies that cycle rate at the reference chemical state with its self consistent voltage. The derivation is recorded in the unchanged Task 36 design and NBC module.

This is a conditional structural construction, not a knockout fit and not a measured transporter density. In particular, equation (2) does not turn the 70/30 partition into experimental evidence. The subsequently evolved trajectory need not retain that partition exactly. The manuscript should state the assumed partition, how it sets the capacity, and what changes when that assumption changes. A broad uncertainty interval for s has not been established by this audit.

The NBC increment is zero at the declared resting calcium and fully recruited at $0.25 \mu\text{M}$. Its width parameter is effective. Its molecular identity is deliberately not assigned to a particular salivary SLC4 isoform.

3.4 Fixed charge and osmotic pools

Reference electroneutrality determines the fixed intracellular anion equivalent amount from

$$X = n_{Na,i} + n_{K,i} - n_{Cl,i} - n_{A,i}.$$

The active value is $78.06176220594485 \text{ fmol equivalents}$. Its status is reference constraint derived, conditional on the reference tuple and carbonate buffer model. It is not a directly counted molecular pool.

The other impermeant osmotic pool is $124.69936937124379 \text{ fmol}$ on the selected background, rather than the smaller dataclass default. It is an inherited effective osmotic setting. The separate finite intracellular buffer amount is 30 fmol and is counted exactly once in the osmolarity. Effective osmotic closure and charge closure are different roles and should not be conflated.

4 Parameters that remain assumptions or transferred settings

The AE4 carrier amount and transition attempt rates are inherited effective model settings. Their thermodynamic construction constrains rate ratios and reversal properties of the parent cycle; it does not supply independent measurements of the absolute carrier number or all attempt rates. Moreover, fixed source substitution in the later mechanism panel does not itself establish thermodynamic consistency of the inherited scalar kinetics with each substituted cycle.

Channel conductances, pump capacity and apical fractions determine current closure and the distribution of secretion. Their active values are explicit in the inventory, but physiological feasibility at one reference point is not evidence that each value is unique. The same applies to the effective outlet rate and dead volume. The code itself describes the outlet law as a transparent compliance closure rather than a calibrated secretion law.

The water coefficients are inherited model conversions with stated units. They should be described as transferred model parameters, not newly measured hydraulic permeabilities of the present preparation. The effective beta activation time of 30 s and AE4 gain of 0.25 likewise need to remain explicit assumptions. Experimental activation of a pathway does not identify all intermediate time constants [Peña Münzenmayer et al., 2021].

The NKCC1 algebraic recruitment reaches its endpoint at 0.10 μM calcium, while NBC reaches its endpoint at 0.25 μM . The production input fully recruits both. This difference is preserved and recorded, not silently reconciled. Any analysis of submaximal calcium would need to account for it.

The numerical pH bracket from 3 to 11 only enables inversion; the actual inherited physiological criterion is pH 6.6 to 7.3. The tiny positive luminal buffer amount 10^{-9} fmol is numerical regularisation approximating no fixed luminal buffer, not an abundance measurement. Constructor concentrations and volumes do not replace the frozen solved initial state used by the production experiments.

Finally, the Task 41 residual CaCC recruitment coefficient $b = 0.10511872843288446$ is target selected. It is not an active parameter of the parent Task 40 model and is not reclassified as literature supported merely because the resulting trajectory has an appropriate secretion magnitude.

The CCh and IPR dose fields, 0.3 and 5 μM respectively, are experiment labels in this implementation. The selected input callable returns prescribed calcium and beta activation from the input arm and time window; it does not map those dose values through a dose response law. They are therefore retained as protocol metadata but marked inactive for the selected right hand side. The previous inventory incorrectly marked both active. Changing the nominal dose labels alone cannot change the model trajectory.

The frozen initial state is now recorded separately, with all twelve conserved amount and volume coordinates and the AE4 regulatory fraction. The twelve constructor reference fields remain distinct. The upstream NBC design quantities, selected transport laws, R1 regulatory family and WT construct are also explicit. These additions explain the larger record count; they add no fitted degrees of freedom.

The externally passed WT genotype object and the three prescribed AE4 expression values, 1, 0.05 and 0, are also recorded. The other transporter expression multipliers remain one in those comparisons. These values define interventions, not an uncertainty interval for protein abundance. The prescribed Task 41 family requires $0 < b \leq 1$, equivalently $0 \leq \rho < 1$; the endpoint $b = 0$ is outside its implemented domain.

The additional file `output/numerical_controls.json` records the production integration specification, conservation tolerances, pH inversion tolerances and voltage bracket. In particular, $[-0.5, 0.5]$ V is the implemented voltage search bracket and not a measured physiological voltage interval. Integration begins at 10^{-6} s with the shared frozen state; the distinct resting observable at exactly $t = 0$ is preserved. Exact unit conversions are stated separately from physiological settings.

5 What the mathematical analysis adds to justification

The separate Task 44 report uses the unchanged reference parameter set and retains carbon, alkalinity, volume, lumen and electrical feedback. It supplies exact structural identities and a conditional NHE supply bound, then evaluates local stationary sensitivities. These calculations

help identify which assumptions affect which conclusions. They do not estimate experimental uncertainty intervals.

An important distinction is between statements independent of capacities, such as the cancellation of the explicit equal routing AE4 term in the chloride balance, and numerical conclusions that depend on the adopted kinetics and scales, such as large NKCC1 replacement. Even an exact balance identity does not determine the magnitude of the knockout flow deficit. The dependence of pump and NHE fluxes on state remains part of the mechanism.

For later manuscript development, every nonliterature value should therefore be accompanied by either its conditional derivation or a plainly stated effective role. Where several values were selected together on the same WT background, their correlation should be retained in future uncertainty work. Independently perturbing every value as though each had its own measured error bar would create an unsupported uncertainty model.

The remaining parameter work is substantive rather than typographical: independently justify or measure ranges for the effective settings; define a WT feasible set using evidence not reused from the knockout comparison; and assess which conclusions survive across that set. A local derivative or a collection of physiologically passing simulations does not prove robustness over all plausible parameterisations. This audit has performed no new fitting and established no calibrated uncertainty intervals.

5.1 Completed sensitivity crosswalk

The completed companion calculation examines sixteen uncertain or effective settings. All 128 nearby WT and AE4 null equilibrium solves pass the inherited physiological gates. Parameter differences of 10^{-3} and 5×10^{-4} in log coordinates are numerical verification steps, not uncertainty ranges. The machine readable crosswalk reads the verified results directly from immutable Task 44 numerical source commit `f783785a863440df459e9c5530beba4c7eb59110` and records SHA256 hashes of all seven imported source files. Later report-only commits need not have that same head SHA. It maps every companion parameter name back to the actual inventory value and classification.

At the adopted constant full stimulus equilibrium, WT secretion is 0.0015840671 pL/s and intracellular pH is 7.0503863. The AE4 null stationary secretion deficit is 0.94617%. This stationary deficit is distinct from the frozen Task 40 cumulative deficit of 3.8575% at 600 s. Intracellular storage, changing volumes, finite lumen composition and regulation remain active in the companion analysis.

In the table, $E_Q = d \log Q_{WT} / d \log p$, $E_D = d \log D / d \log p$ for the positive stationary null deficit D , and $E_\tau = d \log \tau_{slow, WT} / d \log p$. The pH column uses $dpH / d \log p$ because pH is already logarithmic. A large elasticity of a small deficit is not a large absolute percentage point change.

The largest local sensitivities of the stationary null deficit are to pump capacity ($E_D = 4.634$), NBC capacity (2.085), NBC saturation width (-1.502), and NKCC effective capacity (-1.494). These values identify priorities for experimental constraint. They do not establish a finite parameter change that recovers the experimental phenotype. The common stationary effects of NHE carrier amount and its stimulated multiplier reflect their product under full stimulation; separate resting and stimulated measurements are needed to distinguish them.

At the WT equilibrium, NBC supplies 95.48% of the AE4 alkalinity demand. Under the stated bath composition and inherited intracellular pH lower gate of 6.6, the conditional NHE/AE2 capacity calculation covers at most 57.96% of that current demand without NBC. This comparison holds the required AE4 demand at the analysed level; it does not rule out a lower supported flux, another admissible state or an additional pathway. The capacity ratio has local elasticity 0.931 to NHE carrier amount and to the stimulated NHE multiplier. Direct measurements of maximal acid extrusion are therefore essential before treating the conditional

Setting	E_Q	$dpH/d \log p$	E_D	E_τ
NBC capacity	0.019923	0.064832	2.0848	0.10701
AE4 carrier	0.00081137	-0.072461	0.084942	-0.063968
NKCC scale	0.055361	-0.001558	-1.4937	0.19015
NHE carrier	-0.0032611	0.070838	-0.22928	-0.014728
pump capacity	0.52593	0.12061	4.6337	-1.5222
CaCC conductance	0.03247	-0.0055239	-0.22116	0.0082736
buffer pool	-0.0015749	0.0013957	-0.20631	0.30856
water apical	0.0021295	-8.3167e-05	-0.003558	-3.7398e-05
AE4 tau	0	0	0	0
AE4 gain	0.00016227	-0.014492	0.016988	-0.012794
apical pump fraction	0.049564	-0.0035713	-0.059826	0.045169
NHE stimulated multiplier	-0.0032611	0.070838	-0.22928	-0.014728
CO2 basolateral permeability	-0.004448	-0.083186	-0.43378	-0.068179
AE2 capacity	-7.3346e-05	0.00036297	-0.030565	7.5082e-05
NBC log width	-0.014352	-0.046703	-1.5016	-0.072634
NKCC stimulated multiplier	0.055361	-0.001558	-1.4937	0.19015

Table 1: Local stationary sensitivities imported from Task 44. No measured uncertainty intervals are implied.

obstruction as a biological conclusion.

The companion continuation also finds a physiological equilibrium with zero NBC capacity: pH 6.96833 and secretion 0.00153598 pL/s, only 3.035% below the full NBC equilibrium. Its AE4 cycle flux falls from 0.0750724 to 0.00632651 fmol/s. Consequently, any earlier broad claim that NBC is structurally necessary for secretion must be qualified: the proved obstruction concerns sustaining the specified AE4 loading level. It does not imply that all physiological secretion fails without NBC.

The WT and null slowest local decay times are approximately 346 and 460 s. The WT decay time is particularly sensitive to pump capacity ($E_\tau = -1.522$) and the finite intracellular buffer pool (0.309). The model therefore already contains slow modes. The effective AE4 regulatory time has zero local stationary influence in the selected R1 family while the slower chemical mode remains dominant; this is not evidence that time dependent regulation is irrelevant. Acute pH, TIC and volume measurements alongside secretion would constrain those chemical modes and their observable effects.

The separate inverse calculation stays inside the one prescribed Task 41 family. Its first numerical 600 s crossing requires an AE4 dependent stimulated CaCC fraction $\rho \simeq 0.89507$, approximately 89.51%. The original selected value $\rho = 1 - b = 0.89488127$ gives the frozen 30.2612% deficit, slightly below the exact 30.3% comparison threshold. The threshold is 35% minus the reported standard error of 4.7 percentage points in the 2015 comparison; it is not a confidence bound. Adaptive integration and the original sampled trapezoidal observable are retained separately in the companion files. This is the first crossing found in the ordered numerical scan. Monotonicity between samples and exclusion of earlier unsampled crossings are unproved; it is neither a global lower bound nor a universal recruitment requirement.

Inverse sensitivities hold the inherited conserved WT initial state fixed, perturb the WT denominator and null system together, and perform no new resting solve. A buffer perturbation therefore changes the initial algebraic pH. Within that family, the adaptive crossing has local log elasticity 0.09833 to CaCC conductance and -0.03558 to pump capacity, larger in magnitude than its elasticities to the other six tested settings. The relative sensitivity of the residual independent recruitment $1 - \rho$ is also recorded, because a small change in ρ can be a larger relative change in the remaining approximately 10.49%. Under continued full stimulation, both the selected and crossing null cases eventually violate the inherited upper pH gate at

about 1485 s. This diagnostic does not change their declared 600 s results, but prevents an unsupported long duration interpretation.

The most urgent experiments consequently differ by claim. Functional pump turnover and membrane distribution, NBC capacity and saturation shape, and ion conditioned NKCC uptake constrain secretion and compensation. Maximal NHE acid extrusion constrains the conditional alkalinity obstruction. Buffer titration with TIC and volume time series constrains slow relaxation. Matched CaCC current measurements during acute AE4 perturbation would directly test the very large recruitment dependence required by the inverse family. Strong modelled NKCC compensation remains in tension with the isolated 2015 NKCC assay; changing the assay context in the model does not erase that discrepancy.

6 Reproducibility and use in the manuscript

Run `python analysis/43_parameter_provenance/audit.py` from the repository root. It verifies the production parameter hashes and writes both CSV and JSON inventories under this task’s `output` directory. Each record contains source paths and justification notes beyond the abbreviated table below. `prepare_report.py` typesets the table from those records. The report and the full inventory belong on the Task 43 branch, separate from the mathematical analysis and from the main manuscript.

The inventory has 137 records and 109 active flags. This is an inventory of configuration and roles, not a count of independent degrees of freedom: it includes physical constants, protocol settings, law selections, derived values and other nonfit quantities. Five legacy fields and twelve constructor seed fields are explicitly separated. The Task 41 inverse coefficient is also outside the parent model.

Run `analysis/43_parameter_provenance/verify.py` to compare the saved CSV/JSON products with a fresh reconstruction, check every nested production field, verify source hashes, confirm the fixed Cha exponents and demonstrate that changing dose labels leaves the selected input unchanged. The audit performs no fresh sensitivity simulations. With the companion commit available in the fetched Git history, `import_crosswalk.py--check` independently reconstructs the saved crosswalk from that exact commit; no adjacent Task 44 working directory is required. The report tables are also checked against their machine readable sources. The source manifest contains SHA256 values for the production modules and frozen inputs; verification also checks that protected paths are unchanged from recovered Task 43 checkpoint 693f44c0b05e92faec737aa7793c3a99ec76445c.

No model source, historical analysis, frozen trajectory or main manuscript is modified by this audit. The exact original metadata is retained in the machine readable inventory, while the new classification states what that metadata does and does not justify.

A Complete active object and legacy inventory

The “Used” column denotes use by the selected production configuration as classified by the audit. “No” for constructor seeds means they are not the production initial state; the constructor still uses that tuple for reference consistency checks. Full units, source locations, defaults and detailed justifications are in `parameter_inventory.csv` and `parameter_inventory.json`.

Setting	Value	Used	Classification
<code>constants.temperature_K</code>	310.15	Yes	effective assumption
<code>constants.gas_constant_J_mol_K</code>	8.31446262	Yes	physical constant
<code>constants.faraday_C_mol</code>	96485.3321	Yes	physical constant
<code>acid_base.carbon_pka1</code>	6.1	Yes	inherited model setting
<code>acid_base.carbon_pka2</code>	10.3	Yes	inherited model setting

Setting	Value	Used	Classification
acid_base.water_pkw	14	Yes	inherited model setting
acid_base.cell_buffer_pka	7	Yes	effective assumption
acid_base.lumen_buffer_pka	7	Yes	effective assumption
acid_base.ph_lower	3	Yes	numerical bracket
acid_base.ph_upper	11	Yes	numerical bracket
bath.na_mM	145	Yes	effective assumption
bath.k_mM	5	Yes	effective assumption
bath.cl_mM	126.161593	Yes	reference constraint derived
bath.tic_mM	25	Yes	effective assumption
bath.ph	7.4	Yes	effective assumption
bath.buffer_total_mM	0	Yes	effective assumption
bath.untracked_osmolyte_mM	0	Yes	effective assumption
geometry.cell_buffer_total_fmole	30	Yes	effective assumption
geometry.lumen_buffer_total_fmole	1e-09	Yes	numerical regularisation
geometry.fixed_cell_anion_equivalents_fmole	78.0617622	Yes	reference constraint derived
geometry.cell_impermeant_osmoles_fmole	124.699369	Yes	inherited effective setting
homeostasis.nkcc1_capacity_fmole_s	0.32	No	inactive legacy
homeostasis.nhe1_capacity_fmole_s	0.00744098059	No	inactive legacy
homeostasis.nhe1_model	cha_2009_eight_state_mod2	Yes	law selection
homeostasis.nhe1_published_g_fmole_s	0.0305	No	inactive legacy
homeostasis.nhe1_published_k_h_mM	0.00045	No	inactive legacy
homeostasis.nhe1_published_k_na_mM	15	No	inactive legacy
homeostasis.nhe1_cha_carrier_amount_fmole	2.33937001e-05	Yes	WT constraint derived
homeostasis.ae2_capacity_fmole_s	0.005	Yes	effective assumption
homeostasis.thermodynamic_saturation_log_width	2	Yes	effective assumption
homeostasis.co2_basolateral_permeability_fmole_s_mM	0.02	Yes	effective assumption
homeostasis.co2_apical_permeability_fmole_s_mM	0.01	Yes	effective assumption
membranes.nak_capacity_fmole_s	0.08	Yes	effective assumption
membranes.nak_na_half_mM	10	Yes	inherited model setting
membranes.nak_k_half_mM	1.5	Yes	inherited model setting
membranes.apical_pump_fraction	0.075075	Yes	effective assumption
membranes.apical_k_fraction	0.3	Yes	effective assumption
membranes.g_k_total_S	1.4e-08	Yes	effective assumption
membranes.g_cl_apical_S	3.14e-08	Yes	effective assumption
membranes.calcium_half_uM	0.26	Yes	effective assumption
membranes.calcium_hill	1.46	Yes	effective assumption
membranes.g_para_na_S	3e-10	Yes	inherited model setting
membranes.g_para_k_S	5e-11	Yes	inherited model setting
membranes.g_para_cl_S	2.5e-10	Yes	effective assumption
membranes.g_para_hco3_S	5e-11	Yes	effective assumption
membranes.g_apical_background_S	0	Yes	effective assumption
membranes.g_basolateral_background_S	0	Yes	effective assumption

Setting	Value	Used	Classification
water.apical_hydraulic_pL_s_mOsm	0.00432	Yes	published model conversion
water.basolateral_hydraulic_pL_s_mOsm	0.0515	Yes	published model conversion
water.paracellular_hydraulic_pL_s_mOsm	0.00026	Yes	published model conversion
water.lumen_dead_volume_pL	0.1	Yes	effective assumption
water.outflow_rate_s	0.2	Yes	effective assumption
initial.cell_na_mM	15	No	constructor seed
initial.cell_k_mM	140	No	constructor seed
initial.cell_cl_mM	40	No	constructor seed
initial.cell_tic_mM	20	No	constructor seed
initial.cell_ph	7.2	No	constructor seed
initial.cell_volume_pL	1	No	constructor seed
initial.lumen_na_mM	145	No	constructor seed
initial.lumen_k_mM	5	No	constructor seed
initial.lumen_cl_mM	126.161593	No	constructor seed
initial.lumen_tic_mM	25	No	constructor seed
initial.lumen_ph	7.4	No	constructor seed
initial.lumen_volume_pL	0.1	No	constructor seed
ae4.carrier_amount_fmole	0.150126427	Yes	inherited effective setting
ae4.common_cl_attempt_rate_s	1	Yes	inherited effective setting
ae4.na_loaded_attempt_rate_s	0.1	Yes	inherited effective setting
ae4.k_loaded_attempt_rate_s	1.9	Yes	inherited effective setting
ae4.cooperative_gate	None	No	inherited effective setting
nbc.capacity_fmole_s	0.115709132	Yes	WT architecture derived
nbc.log_width	2	Yes	effective assumption
nbc.resting_calcium_uM	0.058	Yes	effective assumption
nbc.fully_recruited_calcium_uM	0.25	Yes	effective assumption
nbc.nhe1_stimulated_multiplier	2.3	Yes	effective assumption
nkcc.alpha_eff_fmole_s	0.0173347469	Yes	WT constraint derived
nkcc.A1	157.5	Yes	published kinetic coefficient
nkcc.A2	2.0096e-05	Yes	published kinetic coefficient
nkcc.A3	1.0306	Yes	published kinetic coefficient
nkcc.A4	1.3852e-06	Yes	published kinetic coefficient
cha.k1_plus_per_ms	10.5	Yes	published kinetic coefficient
cha.k1_minus_per_ms	0.201	Yes	published kinetic coefficient
cha.k2_plus_per_ms	15.8	Yes	published kinetic coefficient
cha.k_na_o_mM	195	Yes	published kinetic coefficient
cha.k_na_i_mM	16.2	Yes	published kinetic coefficient
cha.k_h_o_mM	0.00162	Yes	published kinetic coefficient
cha.k_h_i_mM	0.000605	Yes	published kinetic coefficient
cha.modifier_k_i_mM	3.07e-05	Yes	published kinetic coefficient

Setting	Value	Used	Classification
cha.modifier_k_o_mM	4.8e-07	Yes	published kinetic coefficient
cha.k2_minus_table_per_ms	183	Yes	published kinetic coefficient
regulation.tau_ae4_s	30	Yes	effective assumption
regulation.basal_capacity_multiplier	1	Yes	effective assumption
regulation.fully_activated_increment	0.25	Yes	effective assumption
regulation.coupling_scale	1	Yes	effective assumption
nkcc_activation.fully_activated_multiplier	1.75	Yes	inherited effective setting
nkcc_activation.resting_calcium_uM	0.058	Yes	inherited effective setting
nkcc_activation.stimulated_calcium_uM	0.1	Yes	inherited effective setting
protocol.arm	CCH_IPR	Yes	protocol setting
protocol.onset_s	0	Yes	protocol setting
protocol.duration_s	600	Yes	protocol setting
protocol.cch_dose_uM	0.3	No	protocol setting
protocol.ipr_dose_uM	5	No	protocol setting
protocol.resting_calcium_uM	0.058	Yes	protocol setting
protocol.stimulated_calcium_uM	0.25	Yes	protocol setting
protocol.beta_occupancy_on	1	Yes	protocol setting
Task41.b	0.105118728	No	target selected
cha.intracellular_modifier_hill	3	Yes	published kinetic coefficient
cha.extracellular_modifier_hill	1	Yes	published kinetic coefficient
nkcc.law	palk_benjamin_2010_eq17	Yes	law selection
ae4.law	equal_routing_adapter	Yes	law selection
regulation.family	R1	Yes	law selection
regulation.construct	WT	Yes	protocol setting
nkcc_activation.family	N1_CCH_ALGEBRAIC	Yes	law selection
nbc_design.TARGET_NKCC1_POSITIVE_CL_SHARE	0.7	No	effective assumption
nbc_design.TASK31_R09_NKCC1_CL_FMOL_S	0.256244405	No	WT architecture derived
nbc_design.TASK31_R09_NHE1_FMOL_S	0.00789058049	No	WT architecture derived
nbc_design.DERIVED_AE4_CL_FMOL_S	0.109819031	No	WT architecture derived
nbc_design.DERIVED_STIMULATED_NHE1_FMOL_S	0.0181483351	No	WT architecture derived
nbc_design.DERIVED_REQUIRED_NBC_CYCLE_FMOL_S	0.100744863	No	WT architecture derived
frozen_initial.na_i_fmol	16.7203587	Yes	WT constraint derived
frozen_initial.k_i_fmol	170.002009	Yes	WT constraint derived
frozen_initial.cl_i_fmol	88.031802	Yes	WT constraint derived
frozen_initial.tic_i_fmol	8.26351181	Yes	WT constraint derived
frozen_initial.alk_i_fmol	20.6288031	Yes	WT constraint derived
frozen_initial.volume_i_pL	1.45333439	Yes	WT constraint derived
frozen_initial.na_l_fmol	13.1277687	Yes	WT constraint derived
frozen_initial.k_l_fmol	2.71897261	Yes	WT constraint derived

Setting	Value	Used	Classification
frozen_initial.cl_l_fmol	15.3786879	Yes	WT constraint derived
frozen_initial.tic_l_fmol	0.540262316	Yes	WT constraint derived
frozen_initial.alk_l_fmol	0.468053423	Yes	WT constraint derived
frozen_initial.volume_l_pL	0.105388407	Yes	WT constraint derived
frozen_initial.ae4_effective_activation_fraction	0	Yes	WT constraint derived
genotype.name	WT	No	protocol setting
genotype.ae4_expression	1	Yes	protocol setting
genotype.ae2_expression	1	Yes	protocol setting
genotype.nkcc1_expression	1	Yes	protocol setting
genotype.nhe1_expression	1	Yes	protocol setting
protocol.ae4_expression_cases	[1.0, 0.05, 0.0]	Yes	protocol setting

References

- B. A. Benjamin and E. A. Johnson. A quantitative description of the Na–K–2Cl cotransporter and its conformity to experimental data. *American Journal of Physiology-Renal Physiology*, 273(3):F473–F482, 1997. doi: 10.1152/ajprenal.1997.273.3.F473.
- Marcelo A. Catalán, Lisandra Flores-Aldama, Fernanda Fernández, Daniel Bustos, Natalia Apablaza, Ailen Hidalgo, Yuliet Mazola, Ella Matamala, Li Yo Kao, Ira Kurtz, Carlos Spichiger, José M. Sarmiento, Sebastián Brauchi, Wendy González, Leandro Zúñiga, and Gaspar Peña Münzenmayer. Molecular determinants of bicarbonate and cation transport in the human cation-dependent chloride–bicarbonate exchanger AE4. *American Journal of Physiology-Cell Physiology*, 328(6):C2070–C2084, 2025. doi: 10.1152/ajpcell.00346.2024.
- Chae Young Cha, Chiaki Oka, Yung E. Earm, Shigeo Wakabayashi, and Akinori Noma. A model of sodium–proton exchanger and its central role in regulation of pH and sodium in cardiac myocytes. *Biophysical Journal*, 97(10):2674–2683, 2009. doi: 10.1016/j.bpj.2009.08.053.
- Laurence Palk, James Sneyd, Trevor J. Shuttleworth, David I. Yule, and Edmund J. Crampin. A dynamic model of saliva secretion. *Journal of Theoretical Biology*, 266(4):625–640, 2010. doi: 10.1016/j.jtbi.2010.06.027.
- Gaspar Peña Münzenmayer, Marcelo A. Catalán, Yusuke Kondo, Yasna Jaramillo, Frances Liu, Gary E. Shull, and James E. Melvin. Ae4 (Slc4a9) anion exchanger drives chloride-uptake-dependent fluid secretion by mouse submandibular gland acinar cells. *Journal of Biological Chemistry*, 290(17):10677–10688, 2015. doi: 10.1074/jbc.M114.612895.
- Gaspar Peña Münzenmayer, A. T. George, Gary E. Shull, James E. Melvin, and Marcelo A. Catalán. Ae4 (Slc4a9) is an electroneutral monovalent cation-dependent chloride–bicarbonate exchanger. *Journal of General Physiology*, 147(5):423–436, 2016. doi: 10.1085/jgp.201611571.
- Gaspar Peña Münzenmayer, Yusuke Kondo, Constanza Salinas, José Sarmiento, Sebastián Brauchi, and Marcelo A. Catalán. Activation of the Ae4 (Slc4a9) cation-driven chloride–bicarbonate exchanger by the camp-dependent protein kinase in salivary gland acinar cells. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, 321(6):G628–G638, 2021. doi: 10.1152/ajpgi.00145.2021.
- Elías Vera-Sigüenza, Marcelo A. Catalán, Gaspar Peña Münzenmayer, James E. Melvin, and James Sneyd. A mathematical model supports a key role for Ae4 (Slc4a9) in salivary gland secretion. *Bulletin of Mathematical Biology*, 80:255–282, 2018. doi: 10.1007/s11538-017-0370-6.