


Attractors and contact differences in a four-dimensional atlas

Direct/direct similarities · Equal contractions $w = 1/2$ · Six linked parameters


A $\rho \approx 1.467780$

$$\alpha/\pi = \frac{107}{150} \quad \gamma/\pi = \frac{62}{75}$$




C $\rho \approx 1.633886$

$$\alpha/\pi = \frac{17}{50} \quad \gamma/\pi = \frac{44}{75}$$




E $\rho \approx 1.474340$

$$\alpha/\pi = \frac{1}{50} \quad \gamma/\pi = \frac{34}{75}$$




B $\rho \approx 1.554649$

$$\alpha/\pi = \frac{109}{150} \quad \gamma/\pi = \frac{83}{75}$$





D $\rho \approx 1.532699$

$$\alpha/\pi = \frac{67}{150} \quad \gamma/\pi = \frac{39}{25}$$



F $\rho \approx 1.518192$

$$\alpha/\pi = \frac{1}{50} \quad \gamma/\pi = \frac{16}{75}$$



$$f_{-,p}(z) = -1 + \lambda_- z, \quad f_{+,p}(z) = 1 + \lambda_+ z$$

$$\lambda_- = \rho^{-1} e^{i(\alpha-\gamma)}, \quad \lambda_+ = \rho^{-1} e^{-i(\alpha+\gamma)}$$

$$D_p = \lambda_- K_p - \lambda_+ K_p$$

$$K_p \text{ connected} \iff 2 \in D_p$$

$$f_{-,p}(K_p)$$

$$f_{+,p}(K_p)$$

D_p : shared finite probability-density scale

0 0.005 0.02 0.1 ≥ 0.475

Gray: finite convex hulls. Rings: branch fixed points. Scale bars: one complex-plane unit.

Finite numerical atlas: hue records a selected pair-address code. Each attractor uses all length-22 words.

The torus is an angular display; its central hole is imposed by the projection. Exact contact is not certified by these samples.