

Proof Theory of Modal Logic

Lecture 3 (and a half): Hypersequent calculi (part I)

Tiziano Dalmonte & Marianna Girlando

Free University of Bozen-Bolzano
ILLC, University of Amsterdam

ESSLLI 2024, Leuven,
5-9 August 2024

- ▶ Lecture 1: Sequent calculi
- ▶ Lecture 2: Labelled sequent calculi
- ▶ In this lecture we start looking at **structured calculi**, that extend sequent calculi with **additional structural connectives**

In particular, we now look at **hypersequent calculi**

- ▶ Simple generalisation of sequent calculi
- ▶ Introduced by [Mints, 1968] [Pottinger, 1983], [Avron, 1987] to provide cut-free calculi for modal and relevant logics

 In this lecture we mostly focus on modal logic S5

Axiomatisation of S5

$$\begin{array}{lcl} K + & t \Box A \rightarrow A & \\ & 4 \Box A \rightarrow \Box \Box A & \text{or} \\ & b A \vee \Box \neg \Box A & \end{array} \quad \begin{array}{lcl} K + & t \Box A \rightarrow A & \\ & 5 \Box A \vee \Box \neg \Box A & \end{array}$$

Semantics of S5

Kripke models with equivalence relation

Complexity of S5

The validity/derivability problem for S5 is coNP-complete

Recap

- ▶ No cut-free, Gentzen-style **sequent calculus** for S5 (Lecture 1)
- ▶ **Cut-free labelled calculus** for S5 (Lecture 2)
- ▶ What about an internal, structured calculus for S5?

A hypersequent calculus for S5

Main reference for this calculus

- ▶ *A cut-free simple sequent calculus for modal logic S5*
[Poggiolesi, 2008]: Definition of the calculus and structural analysis

Further references

- ▶ [Lellmann, 2016]: Optimal proof-search procedure in the calculus
- ▶ [Restall, 2007]: A version of the calculus with explicit structural rules


Hypersequent Finite **multiset of sequents**, written

$$\Gamma_1 \Rightarrow \Delta_1 \mid \dots \mid \Gamma_n \Rightarrow \Delta_n$$

where $\Gamma_1 \Rightarrow \Delta_1, \dots, \Gamma_n \Rightarrow \Delta_n$ are the **components** of the hypersequent

Formula interpretation

$$\begin{aligned} i(\Gamma_1 \Rightarrow \Delta_1 \mid \dots \mid \Gamma_n \Rightarrow \Delta_n) \\ = \\ \Box(\bigwedge \Gamma_1 \rightarrow \bigvee \Delta_1) \vee \dots \vee \Box(\bigwedge \Gamma_n \rightarrow \bigvee \Delta_n) \end{aligned}$$

 Differently from labelled sequents, hypersequents can be interpreted as formulas

Initial hypersequents and propositional hypersequent rules

$$\text{init } p, \Gamma \Rightarrow \Delta, p \quad \leadsto \quad \text{init } \mathcal{H} \mid p, \Gamma \Rightarrow \Delta, p$$

$$\vee_R \frac{\Gamma \Rightarrow \Delta, A, B}{\Gamma \Rightarrow \Delta, A \vee B} \quad \leadsto \quad \vee_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A, B}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A \vee B}$$

Modal rules for S5

$$\Box_L \frac{\mathcal{H} \mid \Box A, \Gamma \Rightarrow \Delta \mid A, \Sigma \Rightarrow \Pi}{\mathcal{H} \mid \Box A, \Gamma \Rightarrow \Delta \mid \Sigma \Rightarrow \Pi} \quad \Box_L^t \frac{\mathcal{H} \mid A, \Box A, \Gamma \Rightarrow \Delta}{\mathcal{H} \mid \Box A, \Gamma \Rightarrow \Delta}$$

$$\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A}$$

Example. Derivation of axiom B

$$\frac{\frac{\frac{A \Rightarrow A \mid \Box A \Rightarrow}{\Rightarrow A \mid \Box A \Rightarrow} \Box_L}{\Rightarrow A \mid \Rightarrow \neg \Box A} \neg_R}{\Rightarrow A, \Box \neg \Box A} \Box_R}{\Rightarrow A \vee \Box \neg \Box A} \vee_R$$

Exercise. Derive axioms k, t, 4, 5

Soundness

Theorem. If $\vdash_{\mathbf{HS5}} \mathcal{H}$, then $\vdash_{\mathbf{S5}} i(\mathcal{H})$

Proof sketch (i). We consider simple instances of the rules

$$\Box_L \frac{\Box A \Rightarrow | A \Rightarrow B}{\Box A \Rightarrow | \Rightarrow B}$$

- i. $\vdash \Box \neg \Box A \vee \Box(A \rightarrow B)$ ($i(P)$)
- ii. $\vdash \Box \neg \Box A \vee \neg \Box \neg \Box A$ (CPL)
- iii. $\vdash \neg \Box \neg \Box A \rightarrow \Box A$ (axiom 5)
- iv. $\vdash \Box A \wedge \Box(A \rightarrow B) \rightarrow \Box B$ (axiom k)
- v. $\vdash \Box \neg \Box A \vee \Box B = i(C)$ (by classical reasoning)

Soundness

Theorem. If $\vdash_{\text{HS5}} \mathcal{H}$, then $\vdash_{\text{S5}} i(\mathcal{H})$

Proof sketch (ii). We consider simple instances of the rules

$$\Box_R \frac{B \Rightarrow C \mid \Rightarrow A}{B \Rightarrow C, \Box A}$$

- | | | |
|-------|--|--------------------------------|
| i. | $\vdash \Box(B \rightarrow C) \vee \Box A$ | $(i(P))$ |
| ii. | $\vdash (B \rightarrow C) \rightarrow (B \rightarrow C \vee \Box A)$ | (CPL) |
| iii. | $\vdash \Box(B \rightarrow C) \rightarrow \Box(B \rightarrow C \vee \Box A)$ | $(\text{ii, by K valid rule})$ |
| iv. | $\vdash \Box A \rightarrow (B \rightarrow C \vee \Box A)$ | (CPL) |
| v. | $\vdash \Box \Box A \rightarrow \Box(B \rightarrow C \vee \Box A)$ | $(\text{iv, by K valid rule})$ |
| vi. | $\vdash \Box A \rightarrow \Box \Box A$ | (axiom 4) |
| vii. | $\vdash \Box A \rightarrow \Box(B \rightarrow C \vee \Box A)$ | (from iv, vi) |
| viii. | $\vdash \Box(B \rightarrow C \vee \Box A) = i(C)$ | $(\text{from i, iii, vii})$ |

Soundness

Theorem. If $\vdash_{\mathbf{HS5}} \mathcal{H}$, then $\vdash_{\mathbf{S5}} i(\mathcal{H})$

Proof sketch (ii). We consider simple instances of the rules

$$\Box_R \frac{B \Rightarrow C \mid \Rightarrow A}{B \Rightarrow C, \Box A}$$

- | | | |
|-------|--|--------------------------------|
| i. | $\vdash \Box(B \rightarrow C) \vee \Box A$ | $(i(P))$ |
| ii. | $\vdash (B \rightarrow C) \rightarrow (B \rightarrow C \vee \Box A)$ | (CPL) |
| iii. | $\vdash \Box(B \rightarrow C) \rightarrow \Box(B \rightarrow C \vee \Box A)$ | $(\text{ii, by K valid rule})$ |
| iv. | $\vdash \Box A \rightarrow (B \rightarrow C \vee \Box A)$ | (CPL) |
| v. | $\vdash \Box \Box A \rightarrow \Box(B \rightarrow C \vee \Box A)$ | $(\text{iv, by K valid rule})$ |
| vi. | $\vdash \Box A \rightarrow \Box \Box A$ | (axiom 4) |
| vii. | $\vdash \Box A \rightarrow \Box(B \rightarrow C \vee \Box A)$ | (from iv, vi) |
| viii. | $\vdash \Box(B \rightarrow C \vee \Box A) = i(C)$ | $(\text{from i, iii, vii})$ |

Exercise. Prove soundness of all the rules of **HS5**

Theorem. All rules of **HS5** are **hp-invertible**

Sketch of proof. For each rule, the proof proceeds by induction on the height h of the derivation of the conclusion.

Theorem. All rules of **HS5** are **hp-invertible**

Sketch of proof. For each rule, the proof proceeds by induction on the height h of the derivation of the conclusion. We consider as an example the rule \Box_R

$$\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A}$$

Theorem. All rules of **HS5** are **hp-invertible**

Sketch of proof. For each rule, the proof proceeds by induction on the height h of the derivation of the conclusion. We consider as an example the rule \Box_R

$$\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A}$$

(base case) If $h = 0$, then the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$ of \Box_R is an initial hypersequent. There are three possibilities:

1. \mathcal{H} is an initial hypersequent
2. $p \in \Gamma \cap \Delta$ for some p
3. $\perp \in \Gamma \cap \Delta$

Theorem. All rules of **HS5** are **hp-invertible**

Sketch of proof. For each rule, the proof proceeds by induction on the height h of the derivation of the conclusion. We consider as an example the rule \Box_R

$$\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A}$$

(base case) If $h = 0$, then the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$ of \Box_R is an initial hypersequent. There are three possibilities:

1. \mathcal{H} is an initial hypersequent
2. $p \in \Gamma \cap \Delta$ for some p
3. $\perp \in \Gamma \cap \Delta$

In each of these cases, the premiss $\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A$ of \Box_R is an initial hypersequent, hence it is derivable with height 0.

(inductive step) If $h > 0$, we need to consider the last rule application in the derivation of the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$.

(inductive step) If $h > 0$, we need to consider the last rule application in the derivation of the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$. Since \mathcal{H} can be any hypersequent, and Γ, Δ can contain any formula, the last rule applied can be any rule, hence one needs to consider all of them ...

(inductive step) If $h > 0$, we need to consider the last rule application in the derivation of the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$. Since \mathcal{H} can be any hypersequent, and Γ, Δ can contain any formula, the last rule applied can be any rule, hence one needs to consider all of them ... Here we consider as an example the case where the last rule applied is \Box_R itself

(inductive step) If $h > 0$, we need to consider the last rule application in the derivation of the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$. Since \mathcal{H} can be any hypersequent, and Γ, Δ can contain any formula, the last rule applied can be any rule, hence one needs to consider all of them ... Here we consider as an example the case where the last rule applied is \Box_R itself

There are two possibilities

1. $\Box A$ is principal in the last rule application
2. $\Box A$ is not principal in the last rule application

(inductive step) If $h > 0$, we need to consider the last rule application in the derivation of the conclusion $\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A$. Since \mathcal{H} can be any hypersequent, and Γ, Δ can contain any formula, the last rule applied can be any rule, hence one needs to consider all of them ... Here we consider as an example the case where the last rule applied is \Box_R itself

There are two possibilities

1. $\Box A$ is principal in the last rule application
2. $\Box A$ is not principal in the last rule application

(case 1.) If $\Box A$ is principal in the last rule application, then the last rule application is precisely

$$\frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R$$

which means that the premiss $\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A$ has a derivation of height $h - 1$.

(case 2.) If $\Box A$ is not principal in the last rule application, then the last rule application has the form

$$\frac{\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B}{\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R$$

where $\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B = \mathcal{H}$

(case 2.) If $\Box A$ is not principal in the last rule application, then the last rule application has the form

$$\frac{\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B}{\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R$$

where $\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B = \mathcal{H}$ and the premiss $\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B$ has a derivation of height $h - 1$.

(case 2.) If $\Box A$ is not principal in the last rule application, then the last rule application has the form

$$\frac{\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B}{\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R$$

where $\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B = \mathcal{H}$ and the premiss $\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B$ has a derivation of height $h - 1$.

(Alternatively, we may also have $\Box B \in \Delta$, the proof is analogous in this case.)

(case 2.) If $\Box A$ is not principal in the last rule application, then the last rule application has the form

$$\frac{\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B}{\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R$$

where $\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B = \mathcal{H}$ and the premiss $\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B$ has a derivation of height $h - 1$.

(Alternatively, we may also have $\Box B \in \Delta$, the proof is analogous in this case.)

Then, by inductive hypothesis, $\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta \mid \Rightarrow B \mid \Rightarrow A$ has a derivation of height $\leq h - 1$

(case 2.) If $\Box A$ is not principal in the last rule application, then the last rule application has the form

$$\frac{\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B}{\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R$$

where $\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B = \mathcal{H}$ and the premiss $\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow B$ has a derivation of height $h - 1$.

(Alternatively, we may also have $\Box B \in \Delta$, the proof is analogous in this case.)

Then, by inductive hypothesis, $\mathcal{H}' \mid \Sigma \Rightarrow \Pi \mid \Gamma \Rightarrow \Delta \mid \Rightarrow B \mid \Rightarrow A$ has a derivation of height $\leq h - 1$ and by applying \Box_R to this hypersequent, we obtain a derivation of height $\leq h$ of $\mathcal{H}' \mid \Sigma \Rightarrow \Pi, \Box B \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A = \mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A$.

qed

Structural rules

$$\text{wk}_L \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta}{\mathcal{H} \mid A, \Gamma \Rightarrow \Delta}$$

$$\text{wk}_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A}$$

$$\text{ctr}_L \frac{\mathcal{H} \mid A, A, \Gamma \Rightarrow \Delta}{\mathcal{H} \mid A, \Gamma \Rightarrow \Delta}$$

$$\text{ctr}_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A, A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A}$$

$$\text{wk}_{\text{ext}} \frac{\mathcal{H}}{\mathcal{H} \mid \Gamma \Rightarrow \Delta}$$

$$\text{ctr}_{\text{ext}} \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Gamma \Rightarrow \Delta}{\mathcal{H} \mid \Gamma \Rightarrow \Delta}$$

👉 Note: external forms of weakening and contraction

The cut rule

$$\text{cut} \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A \quad \mathcal{H}' \mid A, \Gamma' \Rightarrow \Delta'}{\mathcal{H} \mid \mathcal{H}' \mid \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

Theorem. Left, right and external weakening and contraction are **hp-admissible** in **HS5**

Sketch of proof. By induction on the height of the derivation of the premiss (*exercise*)

Theorem. Left, right and external weakening and contraction are **hp-admissible** in **HS5**

Sketch of proof. By induction on the height of the derivation of the premiss (*exercise*)

Hint. In order to prove the hp-admissibility of some structural rules you may need the following (nice) rule

$$\text{merge} \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Sigma \Rightarrow \Pi}{\mathcal{H} \mid \Gamma, \Sigma \Rightarrow \Delta, \Pi}$$

Theorem. The rule **merge** is hp-admissible in **HS5**

Sketch of proof. By induction on the height of the derivation of the premiss (*exercise*)

Theorem. Cut is admissible in **HS5**

Proof sketch. By induction on the complexity of the cut formula and subinduction on the cut height.

As an example, consider the following derivation, with the cut formula $\Box A$ principal in the last rule application in both premisses of cut

$$\frac{\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A} \quad \frac{\mathcal{H}' \mid A, \Box A, \Gamma' \Rightarrow \Delta'}{\mathcal{H}' \mid \Box A, \Gamma' \Rightarrow \Delta'} \Box_L^t}{\mathcal{H} \mid \mathcal{H}' \mid \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \text{cut}$$

Converted into the following, with one application of cut at a **lower height**, and one application of cut with a cut formula of **lower complexity**

$$\frac{\frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \mathcal{H} \mid \mathcal{H}' \mid \Gamma \Rightarrow \Delta \mid \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \quad \frac{\frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A \quad \mathcal{H}' \mid A, \Box A, \Gamma' \Rightarrow \Delta'}{\mathcal{H} \mid \mathcal{H}' \mid \Gamma, \Gamma', A \Rightarrow \Delta, \Delta'} \text{cut}}{\mathcal{H} \mid \mathcal{H} \mid \mathcal{H}' \mid \Gamma, \Gamma' \Rightarrow \Delta, \Delta' \mid \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \text{cut} \quad \frac{\text{wk}^*}{\text{ctr}_{\text{ext}} \times 2}$$

(where wk^* denotes multiple applications of (left and right) weakening

Completeness

Theorem. If $\vdash_{S5} A$, then $\vdash_{HS5} A$

Proof sketch.

- ▶ All axioms of S5 are derivable in **HS5** (*exercise*)
- ▶ The necessitation rule is admissible in **HS5** (*exercise*)
- ▶ Modus ponens is simulated by cut

So far, purely syntactical analysis.
What about a semantics for the calculus?

Two semantics for S5

1. Kripke models with equivalence relation, or


2. **Universal semantics** $\mathcal{M} = \langle W, v \rangle$

► No binary relation

► $\mathcal{M}, w \Vdash \Box A$ iff for all $u \in W$, $\mathcal{M}, u \Vdash A$

👉 $\Box A$ true somewhere iff A true everywhere

👉 Corresponds to choosing one cluster of a model with equivalence relation



Notation. We denote \mathcal{U} the class of all universal models

- ▶ Different **components** \leadsto different **worlds**
- ▶ For each component, formulas on the **left true**, formulas on the **right false** in the corresponding world

$$\Box_L \frac{\mathcal{H} \mid \Box A, \Gamma \Rightarrow \Delta \mid \textcolor{brown}{A}, \Sigma \Rightarrow \Pi}{\mathcal{H} \mid \Box A, \Gamma \Rightarrow \Delta \mid \Sigma \Rightarrow \Pi} \quad \uparrow$$



- ▶ Different **components** \leadsto different **worlds**
- ▶ For each component, formulas on the **left true**, formulas on the **right false** in the corresponding world

$$\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A} \uparrow$$



Valid hypersequent $\mathcal{M} \models \mathcal{H}$ iff $\exists \Gamma \Rightarrow \Delta \in \mathcal{H} : \mathcal{M} \models \Gamma \Rightarrow \Delta$

- 👉 Semantically, a hypersequent is a **disjunction of validities**
- 👉 That is, \mathcal{H} valid iff one component is valid

Soundness

Theorem. If $\vdash_{\mathbf{HS5}} \mathcal{H}$, then $\models_{\mathcal{U}} \mathcal{H}$

Proof sketch. One needs to show that the initial hypersequent are valid in \mathcal{U} (trivial) and that all rules of **HS5** preserve validity in universal models (*exercise*).

We now prove the opposite direction (completeness of **HS5**), namely that

$$\text{if } \models_{\mathcal{U}} \mathcal{H}, \text{ then } \vdash_{\mathbf{HS5}} \mathcal{H}$$

1. First, we define a terminating (optimal) **proof-search procedure** in **HS5**
2. Then, we show that every failed proof constructed according to this procedure provides a **countermodel** of the root hypersequent: that is, if $\not\vdash_{\mathbf{HS5}} \mathcal{H}$, then $\not\models_{\mathcal{U}} \mathcal{H}$

A proof-search procedure in **HS5**

Main reference [\[Lellmann, 2016\]](#)

As a first step, we consider a **cumulative formulation** of **HS5**

Cumulative formulation of a rule

The principal formula is copied to the premiss(es)

e.g.

$$\vee_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A \vee B, A, B}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, A \vee B}$$

$$\vee_L \frac{\mathcal{H} \mid A, A \vee B, \Gamma \Rightarrow \Delta \quad \mathcal{H} \mid B, A \vee B, \Gamma \Rightarrow \Delta}{\mathcal{H} \mid A \vee B, \Gamma \Rightarrow \Delta}$$

$$\Box_R \frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A}$$

 Remark. The rules \Box_L and \Box_L^t are already in cumulative form

Notation. We call **HS5_{cum}** the calculus defined by the cumulative formulation of the rules of **HS5**

Theorem (Soundness). If $\vdash_{\mathbf{HS5}_{\text{cum}}} \mathcal{H}$, then $\models_{\mathcal{U}} \mathcal{H}$

Proof. The cumulative rules are **admissible** in **HS5**.

Example: Admissibility of the cumulative version of \Box_R in **HS5**

$$\frac{\frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A \mid \Rightarrow A} \text{ by invertibility of } \Box_R}{\frac{\mathcal{H} \mid \Gamma \Rightarrow \Delta \mid \Rightarrow A}{\mathcal{H} \mid \Gamma \Rightarrow \Delta, \Box A} \Box_R} \text{ctr}_{\text{ext}}$$

Therefore: if $\vdash_{\mathbf{HS5}_{\text{cum}}} \mathcal{H}$, then $\vdash_{\mathbf{HS5}} \mathcal{H}$, hence $\models_{\mathcal{U}} \mathcal{H}$

Clearly, the complexity of hypersequents is **not reduced** by backward applications of cumulative rules

👉 In order to ensure termination of backward proof-search, one needs to **avoid redundant rule applications**

Clearly, the complexity of hypersequents is **not reduced** by backward applications of cumulative rules

👉 In order to ensure termination of backward proof-search, one needs to **avoid redundant rule applications**

Local loop-checking condition (LLCC)

An application of a hypersequent rule with premisses $\mathcal{G}_1, \dots, \mathcal{G}_n$ and conclusion \mathcal{H} satisfies the **local loop checking condition** if for each premiss \mathcal{G}_i , there exists a component $\Gamma \Rightarrow \Delta$ in \mathcal{G}_i such that for no component $\Sigma \Rightarrow \Pi$ of the conclusion \mathcal{H} we have $\text{set}(\Gamma) \subseteq \text{set}(\Sigma)$ and $\text{set}(\Delta) \subseteq \text{set}(\Pi)$

Example: the following rule applications violate the LLCC

$$\frac{\Rightarrow p \wedge q, q, p}{\Rightarrow p \wedge q, q} \wedge_R \quad \frac{p \Rightarrow q \mid r \Rightarrow \Box q \mid \Rightarrow q}{p \Rightarrow q \mid r \Rightarrow \Box q} \Box_R$$

👉 The LLCC prevents the applications of rules that do not add additional information to the hypersequents

Saturated hypersequent

A hypersequent which is not initial and such that no rule is backward applicable to it without violating the LLCC

Backward proof-search with LLCC for \mathcal{H}

The construction of a derivation tree from the root to the leaves such that the root is labelled with the hypersequent \mathcal{H} , and the branches are expanded by applying at each step a backwards applicable rule that **satisfies the LLCC**. The construction terminates when all leaves are labelled with hypersequents that are either **initial or saturated**

The LLCC restricts the backward applicability of the rules

The LLCC restricts the backward applicability of the rules

👉 Is proof-search with LLCC still complete?

The LLCC restricts the backward applicability of the rules

👉 Is proof-search with LLCC still complete?

👉 We now prove that proof-search with LLCC is complete by showing that every hypersequent \mathcal{H} on which it fails is not valid in the universal semantics

The LLCC restricts the backward applicability of the rules

👉 Is proof-search with LLCC still complete?

👉 We now prove that proof-search with LLCC is complete by showing that every hypersequent \mathcal{H} on which it fails is not valid in the universal semantics

👉 In particular, we show that from every failed proof for \mathcal{H} we can extract a countermodel of \mathcal{H}

The LLCC restricts the backward applicability of the rules


👉 Is proof-search with LLCC still complete?

👉 We now prove that proof-search with LLCC is complete by showing that every hypersequent \mathcal{H} on which it fails is not valid in the universal semantics

👉 In particular, we show that from every failed proof for \mathcal{H} we can extract a countermodel of \mathcal{H}

👉 More precisely, we show that each saturated hypersequent occurring in a failed proof of \mathcal{H} provides the information needed to build such countermodel

The procedure in a picture



Let $\mathcal{H} = \Gamma_1 \Rightarrow \Delta_1 \mid \dots \mid \Gamma_n \Rightarrow \Delta_n$ be a **saturated hypersequent** occurring in a **failed proof for \mathcal{G}**

Countermodel extracted from a saturated hypersequent

We define $\mathcal{M} = \langle W, v \rangle$ on the basis of \mathcal{H} as follows

- ▶ $W = \{k \mid \Gamma_k \Rightarrow \Delta_k \in \mathcal{H}\}$
- ▶ For all $p \in \text{Atm}$, $v(p) = \{k \in W \mid p \in \Gamma_k\}$

Countermodel lemma

For all formulas A , for all components $\Gamma_k \Rightarrow \Delta_k$,

- ▶ if $A \in \Gamma_k$, then $k \Vdash A$
- ▶ if $A \in \Delta_k$, then $k \nVdash A$

Proved by induction on the construction of A (*exercise*)

Let $\mathcal{H} = \Gamma_1 \Rightarrow \Delta_1 \mid \dots \mid \Gamma_n \Rightarrow \Delta_n$ be a **saturated hypersequent** occurring in a **failed proof for \mathcal{G}** , and \mathcal{M} be the model defined on the basis of \mathcal{H} as in the previous slide

The countermodel lemma implies that


- ▶ for all $\Gamma_k \Rightarrow \Delta_k \in \mathcal{H}$, $k \not\models \bigwedge \Gamma_k \rightarrow \bigvee \Delta_k$
- ▶ hence, $\mathcal{M} \not\models \mathcal{H}$

Moreover, since all rules are **cumulative**, we have

for all $\Sigma \Rightarrow \Pi \in \mathcal{G}$, there is $\Gamma_k \Rightarrow \Delta_k \in \mathcal{H}$ s.t. $\Sigma \subseteq \Gamma_k$ and $\Pi \subseteq \Delta_k$

Therefore

- ▶ for all $\Sigma \Rightarrow \Pi \in \mathcal{G}$, there is $k \in W$ s.t. $k \not\models \bigwedge \Sigma \rightarrow \bigvee \Pi$
- ▶ hence, $\mathcal{M} \not\models \mathcal{G}$

 \mathcal{M} is a countermodel of the root hypersequent \mathcal{G}

Example of failed proof


$$\begin{array}{c}
 \begin{array}{c} w_1 \quad w_2 \quad w_3 \\
 \hline
 p, p \vee q, \dots \Rightarrow \dots \mid q, \dots \Rightarrow p \mid p, \dots \Rightarrow q \quad \mathcal{X} \\
 \hline
 p \vee q, \dots \Rightarrow \dots \mid q, p \vee q \Rightarrow p \mid p, p \vee q \Rightarrow q \quad \mathcal{X} \\
 \hline
 p \vee q, \Box(p \vee q) \Rightarrow \dots \mid q, p \vee q \Rightarrow p \mid p \vee q \Rightarrow q \\
 \hline
 p \vee q, \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \mid p \vee q \Rightarrow p \mid p \vee q \Rightarrow q \\
 \hline
 p \vee q, \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \mid p \vee q \Rightarrow p \mid p \vee q \Rightarrow q \\
 \hline
 p \vee q, \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \mid p \vee q \Rightarrow p \mid \Rightarrow q \\
 \hline
 p \vee q, \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \mid \Rightarrow p \mid \Rightarrow q \\
 \hline
 \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \mid \Rightarrow p \mid \Rightarrow q \\
 \hline
 \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \mid \Rightarrow p \\
 \hline
 \Box(p \vee q) \Rightarrow \Box p \vee \Box q, \Box p, \Box q \quad \mathcal{X} \\
 \hline
 \Box(p \vee q) \Rightarrow \Box p \vee \Box q \quad \mathcal{X}
 \end{array}
 \end{array}$$

w_1 $p \quad \neg q$ $p \vee q$	w_2 $q \quad \neg p$ $p \vee q$	w_3 $p \quad \neg q$ $p \vee q$
---	---	---

$$\begin{array}{l}
 w_1, w_2, w_3 \models \Box(p \vee q) \\
 w_1, w_2, w_3 \not\models \Box p \vee \Box q
 \end{array}$$

Theorem. Backward proof-search with LLCC in **HS5** provides a NP decision procedure for non derivability in S5

- At each step, non deterministically chose an applicable rule satisfying the LLCC and a correct premiss



This result relies on two key remarks:

1. The length of branches in a proof built by backward proof-search with LLCC is polynomially bounded by the length of the root hypersequent (see next slide)
2. Verifying the LLCC takes polynomial time

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof.

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch.

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula.

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} .

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R .

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R . Moreover, the creation of new identical components is prevented by the LLCC.

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R . Moreover, the creation of new identical components is prevented by the LLCC. Hence, the number of components is bounded by the number of \Box -subformulas of \mathcal{H} .

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R . Moreover, the creation of new identical components is prevented by the LLCC. Hence, the number of components is bounded by the number of \Box -subformulas of \mathcal{H} .

Length of each component: Because of analyticity, each component can only contain subformulas of \mathcal{H} .

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R . Moreover, the creation of new identical components is prevented by the LLCC. Hence, the number of components is bounded by the number of \Box -subformulas of \mathcal{H} .

Length of each component: Because of analyticity, each component can only contain subformulas of \mathcal{H} . Moreover, repetitions of same formulas in the same component is prevented by the LLCC.

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R . Moreover, the creation of new identical components is prevented by the LLCC. Hence, the number of components is bounded by the number of \Box -subformulas of \mathcal{H} .

Length of each component: Because of analyticity, each component can only contain subformulas of \mathcal{H} . Moreover, repetitions of same formulas in the same component is prevented by the LLCC. Hence, the length of each component is bounded by the number of subformulas of \mathcal{H} .

Lemma. The length of branches in a proof for a hypersequent \mathcal{H} built by backward proof-search with LLCC is polynomially bounded by the length n of \mathcal{H}

Sketch of proof. The length of a branch depends on the number of rule applications in that branch. Each backward rule application of a cumulative rule introduces a new formula. Hence, the number of possible rule applications is bounded by the maximal length of a hypersequent that can occur in a proof for \mathcal{H} . In turn, the length of a hypersequent depends on the number of components and the length of each component.

Number of components: New components are introduced only by the rule \Box_R . Moreover, the creation of new identical components is prevented by the LLCC. Hence, the number of components is bounded by the number of \Box -subformulas of \mathcal{H} .

Length of each component: Because of analyticity, each component can only contain subformulas of \mathcal{H} . Moreover, repetitions of same formulas in the same component is prevented by the LLCC. Hence, the length of each component is bounded by the number of subformulas of \mathcal{H} .

Therefore, the length of hypersequents in the proof, hence the length of branches, is in $O(n^2)$

Mono- vs. Multi-modal logics

How many modalities can sequent calculi support?


Sequent and labelled sequent calculi can be **extended to multimodal logics** without essential modifications.

Example. Let K_n be the logic with n K-modalities \Box_1, \dots, \Box_n . The calculus **G3K_n** can be defined considering, for each $i \leq n$, the rule

$$k_i \frac{\Gamma \Rightarrow A}{\Gamma', \Box_i \Gamma \Rightarrow \Box_i A, \Delta}$$

Similarly, a labelled calculus for K_n can be defined considering relational symbols R_1, \dots, R_n and, for each $i \leq n$, the rules

$$\Box_L \frac{xR_i y, x : \Box_i A, y : A, \Gamma \Rightarrow \Delta}{xR_i y, x : \Box_i A, \Gamma \Rightarrow \Delta} \quad \Box_R \frac{xR_i y, \Gamma \Rightarrow \Delta, y : A}{\Gamma \Rightarrow \Delta, x : \Box_i A} (y!)$$

 The properties of the sequent and the labelled calculus for K hold also for the sequent and the labelled calculus for K_n

The same is **not possible** in **HS5**

- ▶ The hypersequent construct | can represent **only one** S5 modality
- ▶ After all, a model can have only one universal modality

However, the universal modality can be **combined with other kinds of modalities**

Example.

Let $K_{\mathcal{U}}$ be the logic with a K modality \Box and a universal modality \blacksquare

Semantics $\mathcal{M} = \langle W, R, v \rangle$, with

- ▶ $\mathcal{M}, w \Vdash \Box A$ iff for all u s.t. wRu , $\mathcal{M}, u \Vdash A$
- ▶ $\mathcal{M}, w \Vdash \blacksquare A$ iff for all u , $\mathcal{M}, u \Vdash A$

(Redundant but complete) **axiomatisation** (cf. [Goranko, Passy, 1992])

- ▶ K axiomatisation for \Box
- ▶ S5 axiomatisation for \blacksquare
- ▶ $\blacksquare A \rightarrow \Box A$

Hypersequent calculus S5 hypersequent calculus for \blacksquare ,
extended with the hypersequent formulation of the rule k for \Box :

$$k \frac{\mathcal{H} \mid \Sigma \Rightarrow A}{\mathcal{H} \mid \Gamma, \Box \Sigma \Rightarrow \Box A, \Delta}$$

Exercise. Derive the axiom $\blacksquare A \rightarrow \Box A$

As we have seen, a sequent $\Gamma \Rightarrow \Delta$ represents a **consequence relation** between the antecedent Γ (the assumptions) and the consequent Δ

But... **which kind of assumptions?**

Global vs. local modal consequence relation

Syntactically $\Gamma \vdash A$ (Hilbert systems)

- ▶ **Global** Both propositional and modal rules (necessitation) can be applied to the assumptions
- ▶ **Local** Only propositional rules be applied to the assumptions

Semantically $\Gamma \models A$

- ▶ **Global** For all \mathcal{M} , $\mathcal{M} \models \bigwedge \Gamma$ implies $\mathcal{M} \models A$
- ▶ **Local** For all \mathcal{M} , for all w , $\mathcal{M}, w \models \bigwedge \Gamma$ implies $\mathcal{M}, w \models A$

Remark.

- The sequent rule


$$\rightarrow_R \frac{A, \Gamma \Rightarrow \Delta, B}{\Gamma \Rightarrow \Delta, A \rightarrow B}$$

expresses the **deduction theorem**, that holds (in this form) for local consequence only

$$\begin{aligned} A \models_{local} B &\leadsto \models_{local} A \rightarrow B \\ A \models_{global} B &\not\leadsto \models_{global} A \rightarrow B \end{aligned}$$

e.g. $A \models_{global} \Box A$ but $\not\models_{global} A \rightarrow \Box A$

- Indeed, validity of modal sequents is defined exactly as the local consequence

 Modal sequents represent **local consequence** relations

The **hypersequent calculus** can be used to **reasoning under global assumptions**

Indeed, reasoning under global assumptions in K:


$$B_1, \dots, B_n \vdash_{global} A$$

can be reduced to

$$\vdash_{K_U} \blacksquare B_1 \wedge \dots \wedge \blacksquare B_n \rightarrow A$$

which is expressed in **HK_U** with the sequent

$$\blacksquare B_1, \dots, \blacksquare B_n \Rightarrow A$$

 We now show that **HK_U** provides a decision procedure for reasoning under global assumptions in K

Main references

- ▶ Avron, [A constructive analysis of RM](#), Journal of Symbolic Logic, 52(4), 1987. 939–951.
- ▶ Avron, [The method of hypersequents in the proof theory of propositional non-classical logics](#), in Logic: From Foundations to Applications, Oxford University Press, 1996
- ▶ Goranko, Passy, [Using the universal modality: gains and questions](#), Journal of Logic and Computation, 2(1), 1992. 5–30.
- ▶ Lellmann, [Hypersequent rules with restricted contexts for propositional modal logics](#), Theoretical Computer Science, 656, 2016. 76–105.
- ▶ Mints, [On some calculi of modal logic](#), Proceedings of the Steklov Institute of Mathematics, 98, 1968. 97–124.
- ▶ Poggiolesi, [A cut-free simple sequent calculus for modal logic S5](#), Review of Symbolic Logic, 2008.
- ▶ Pottinger, [Uniform, cut-free formulations of T, S4 and S5 \(abstract\)](#), Journal of Symbolic Logic, 48(3), 1983. 900.
- ▶ Restall, [Proofnets for S5: sequents and circuits for modal logic](#), in Logic Colloquium 2005, Cambridge University Press, 2007.