

# Proof Theory of Modal Logic

## Lecture 2: Labelled Proof Systems

Tiziano Dalmonte, Marianna Girlando

Free University of Bozen-Bolzano, University of Amsterdam

ESSLLI 2024

Leuven, 5-9 August 2024

## Partial references:


- ▶ [Kanger, 1957] Spotted formulas for S5
- ▶ [Fitting, 1983], [Goré 1998] Tableaux + labels
- ▶ [Simpson, 1994], [Viganò, 1998] Natural deduction + labels
- ▶ [Mints, 1997], [Viganò, 2000], [Negri,2005] Sequent calculus + labels

## We follow the approach of Negri:

- ▶ *Proof analysis in modal logics* [Negri, 2005]
- ▶ *Contraction-free sequent calculi for geometric theories with an application to Barr's theorem* [Negri, 2003]

- ▶ Labelled calculi for K
- ▶ Frame conditions: a general recipe
- ▶ Semantic completeness

## Labelled sequent calculus for K



$$A, B ::= p \mid \perp \mid A \wedge B \mid A \vee B \mid A \rightarrow B \mid \Box A \mid \Diamond A$$

$$A, B ::= p \mid \perp \mid A \wedge B \mid A \vee B \mid A \rightarrow B \mid \Box A \mid \Diamond A$$

Take countably many variables  $x, y, z, \dots$  (the **lables**)

$$A, B ::= p \mid \perp \mid A \wedge B \mid A \vee B \mid A \rightarrow B \mid \Box A \mid \Diamond A$$

Take countably many variables  $x, y, z, \dots$  (the **lables**)

### Labelled formulas

- ▶  $xRy$  meaning ' $x$  has access to  $y$ ' (relational atoms)
- ▶  $x:A$  meaning ' $x$  satisfies  $A$ '

$$A, B ::= p \mid \perp \mid A \wedge B \mid A \vee B \mid A \rightarrow B \mid \Box A \mid \Diamond A$$

Take countably many variables  $x, y, z, \dots$  (the **lables**)

## Labelled formulas

- ▶  $xRy$  meaning ‘ $x$  has access to  $y$ ’ (relational atoms)
- ▶  $x:A$  meaning ‘ $x$  satisfies  $A$ ’

## Labelled sequent

$$\mathcal{R}, \Gamma \Rightarrow \Delta$$

where

- ▶  $\mathcal{R}$  is a multiset of relational atoms;
- ▶  $\Gamma, \Delta$  are multisets of labelled formulas *without* relational atoms.



$$A, B ::= p \mid \perp \mid A \wedge B \mid A \vee B \mid A \rightarrow B \mid \Box A \mid \Diamond A$$

Take countably many variables  $x, y, z, \dots$  (the **lables**)

### Labelled formulas

- ▶  $xRy$  meaning ' $x$  has access to  $y$ ' (relational atoms)
- ▶  $x:A$  meaning ' $x$  satisfies  $A$ '

### Labelled sequent

$$\mathcal{R}, \Gamma \Rightarrow \Delta$$

where

- ▶  $\mathcal{R}$  is a multiset of relational atoms;
- ▶  $\Gamma, \Delta$  are multisets of labelled formulas *without* relational atoms.

Labelled sequents lack a formula interpretation

$$\text{init} \frac{}{\mathcal{R}, x:p, \Gamma \Rightarrow \Delta, x:p}$$

$$\perp_L \frac{}{\mathcal{R}, x:\perp, \Gamma \Rightarrow \Delta}$$

$$\text{init} \frac{}{\mathcal{R}, x:p, \Gamma \Rightarrow \Delta, x:p}$$

$$\wedge_R \frac{\mathcal{R}, x:A, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \wedge B, \Gamma \Rightarrow \Delta}$$

$$\vee_L \frac{\mathcal{R}, x:A, \Gamma \Rightarrow \Delta \quad \mathcal{R}, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \vee B, \Gamma \Rightarrow \Delta}$$

$$\rightarrow_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad \mathcal{R}, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \rightarrow B, \Gamma \Rightarrow \Delta}$$

$$\perp_L \frac{}{\mathcal{R}, x:\perp, \Gamma \Rightarrow \Delta}$$

$$\wedge_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad \mathcal{R}, \Gamma \Rightarrow \Delta, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \wedge B}$$

$$\vee_R \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \vee B}$$

$$\rightarrow_R \frac{x:A, \mathcal{R}, \Gamma \Rightarrow \Delta, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \rightarrow B}$$

$$\text{init} \frac{}{\mathcal{R}, x:p, \Gamma \Rightarrow \Delta, x:p}$$

$$\wedge_R \frac{\mathcal{R}, x:A, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \wedge B, \Gamma \Rightarrow \Delta}$$

$$\vee_L \frac{\mathcal{R}, x:A, \Gamma \Rightarrow \Delta \quad \mathcal{R}, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \vee B, \Gamma \Rightarrow \Delta}$$

$$\rightarrow_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad \mathcal{R}, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \rightarrow B, \Gamma \Rightarrow \Delta}$$

$$\Box_L \frac{xRy, \mathcal{R}, y:A, x:\Box A, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, x:\Box A, \Gamma \Rightarrow \Delta}$$

$$\perp_L \frac{}{\mathcal{R}, x:\perp, \Gamma \Rightarrow \Delta}$$

$$\wedge_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad \mathcal{R}, \Gamma \Rightarrow \Delta, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \wedge B}$$

$$\vee_R \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \vee B}$$

$$\rightarrow_R \frac{x:A, \mathcal{R}, \Gamma \Rightarrow \Delta, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \rightarrow B}$$

$$\Box_R \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, y:A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A} \text{ } y \text{ fresh}$$

$y$  fresh means  $y$  does not occur in  $\mathcal{R} \cup \Gamma \cup \Delta$

$$\begin{array}{c}
 \text{init} \frac{}{\mathcal{R}, x:p, \Gamma \Rightarrow \Delta, x:p} \\
 \wedge_R \frac{\mathcal{R}, x:A, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \wedge B, \Gamma \Rightarrow \Delta} \\
 \vee_L \frac{\mathcal{R}, x:A, \Gamma \Rightarrow \Delta \quad \mathcal{R}, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \vee B, \Gamma \Rightarrow \Delta} \\
 \rightarrow_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad \mathcal{R}, x:B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:A \rightarrow B, \Gamma \Rightarrow \Delta} \\
 \Box_L \frac{xRy, \mathcal{R}, y:A, x:\Box A, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, x:\Box A, \Gamma \Rightarrow \Delta} \\
 \Diamond_L \frac{xRy, \mathcal{R}, y:A, \Gamma \Rightarrow \Delta}{\mathcal{R}, x:\Diamond A, \Gamma \Rightarrow \Delta} y \text{ fresh}
 \end{array}
 \qquad
 \begin{array}{c}
 \perp_L \frac{}{\mathcal{R}, x:\perp, \Gamma \Rightarrow \Delta} \\
 \wedge_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad \mathcal{R}, \Gamma \Rightarrow \Delta, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \wedge B} \\
 \vee_R \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \vee B} \\
 \rightarrow_R \frac{x:A, \mathcal{R}, \Gamma \Rightarrow \Delta, x:B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \rightarrow B} \\
 \Box_R \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, y:A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A} y \text{ fresh} \\
 \Diamond_R \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, x:\Diamond A, y:A}{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, x:\Diamond A}
 \end{array}$$

$y$  fresh means  $y$  does not occur in  $\mathcal{R} \cup \Gamma \cup \Delta$

We write  $\vdash_{\text{labK}} \mathcal{R}, \Gamma \Rightarrow \Delta$  if there is a derivation of  $\mathcal{R}, \Gamma \Rightarrow \Delta$  in labK.

Example:  $\vdash_{\text{labK}} \Rightarrow x:(\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)$

$$\begin{array}{c}
 \begin{array}{c}
 \text{init} \frac{}{xRy, y:p \Rightarrow y:q, x:\Diamond p, y:p} \\
 \Diamond_R \frac{}{xRy, y:A \Rightarrow y:q, x:\Diamond p}
 \end{array}
 \quad
 \begin{array}{c}
 \text{init} \frac{}{xRy, x:\Box q, y:q, y:p \Rightarrow y:q} \\
 \Box_L \frac{}{xRy, x:\Box q, y:p \Rightarrow y:q}
 \end{array} \\
 \hline
 \begin{array}{c}
 \rightarrow_L \frac{}{xRy, x:\Diamond p \rightarrow \Box q, y:p \Rightarrow y:q} \\
 \rightarrow_R \frac{}{xRy, x:\Diamond p \rightarrow \Box q \Rightarrow y:p \rightarrow q} \\
 \Box_R \frac{}{x:\Diamond p \rightarrow \Box q \Rightarrow x:\Box(p \rightarrow q)} \\
 \rightarrow_R \frac{}{\Rightarrow x:(\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)}
 \end{array}
 \end{array}$$



Given a sequent  $\mathcal{S} = \mathcal{R}, \Gamma \Rightarrow \Delta$ , and a model  $\mathcal{M} = \langle W, R, v \rangle$ , let  $\text{Lb}(\mathcal{S}) = \{x \mid x \in \mathcal{R} \cup \Gamma \cup \Delta\}$ , and  $\rho : \text{Lb}(\mathcal{S}) \rightarrow W$  (interpretation).



Given a sequent  $\mathcal{S} = \mathcal{R}, \Gamma \Rightarrow \Delta$ , and a model  $\mathcal{M} = \langle W, R, v \rangle$ , let  $\text{Lb}(\mathcal{S}) = \{x \mid x \in \mathcal{R} \cup \Gamma \cup \Delta\}$ , and  $\rho : \text{Lb}(\mathcal{S}) \rightarrow W$  (interpretation).

Satisfiability of labelled formulas at  $\mathcal{M}$  under  $\rho$  :

$$\mathcal{M}, \rho \Vdash xRy \quad \text{iff} \quad \mathcal{M} \Vdash \rho(x)R\rho(y)$$

$$\mathcal{M}, \rho \Vdash x:A \quad \text{iff} \quad \mathcal{M}, \rho(x) \Vdash A$$

Given a sequent  $\mathcal{S} = \mathcal{R}, \Gamma \Rightarrow \Delta$ , and a model  $\mathcal{M} = \langle W, R, v \rangle$ , let  $\text{Lb}(\mathcal{S}) = \{x \mid x \in \mathcal{R} \cup \Gamma \cup \Delta\}$ , and  $\rho : \text{Lb}(\mathcal{S}) \rightarrow W$  (interpretation).

Satisfiability of labelled formulas at  $\mathcal{M}$  under  $\rho$  :

$$\mathcal{M}, \rho \Vdash xRy \quad \text{iff} \quad \mathcal{M} \Vdash \rho(x)R\rho(y)$$

$$\mathcal{M}, \rho \Vdash x:A \quad \text{iff} \quad \mathcal{M}, \rho(x) \Vdash A$$

Satisfiability of sequents at  $\mathcal{M}$  under  $\rho$  ( $\varphi$  is  $xRy$  or  $x:A$ ):

$$\mathcal{M}, \rho \Vdash \mathcal{R}, \Gamma \Rightarrow \Delta \quad \text{iff}$$

*if* for all  $\varphi \in \mathcal{R} \cup \Gamma$  it holds that  $\mathcal{M}, \rho \Vdash \varphi$ ,

*then* for some  $x:D \in \Delta$  it holds that  $\mathcal{M}, \rho \Vdash x:D$ .

Given a sequent  $\mathcal{S} = \mathcal{R}, \Gamma \Rightarrow \Delta$ , and a model  $\mathcal{M} = \langle W, R, v \rangle$ , let  $\text{Lb}(\mathcal{S}) = \{x \mid x \in \mathcal{R} \cup \Gamma \cup \Delta\}$ , and  $\rho : \text{Lb}(\mathcal{S}) \rightarrow W$  (interpretation).

Satisfiability of labelled formulas at  $\mathcal{M}$  under  $\rho$  :

$$\mathcal{M}, \rho \Vdash xRy \quad \text{iff} \quad \mathcal{M} \Vdash \rho(x)R\rho(y)$$

$$\mathcal{M}, \rho \Vdash x:A \quad \text{iff} \quad \mathcal{M}, \rho(x) \Vdash A$$

Satisfiability of sequents at  $\mathcal{M}$  under  $\rho$  ( $\varphi$  is  $xRy$  or  $x:A$ ):

$$\mathcal{M}, \rho \Vdash \mathcal{R}, \Gamma \Rightarrow \Delta \quad \text{iff}$$

*if* for all  $\varphi \in \mathcal{R} \cup \Gamma$  it holds that  $\mathcal{M}, \rho \Vdash \varphi$ ,

*then* for some  $x:D \in \Delta$  it holds that  $\mathcal{M}, \rho \Vdash x:D$ .

A sequent  $\mathcal{R}, \Gamma \Rightarrow \Delta$  has a countermodel iff there are  $\mathcal{M}, \rho$  such that:

- ▷  $\mathcal{M}, \rho \Vdash \varphi$ , for all  $\varphi \in \mathcal{R} \cup \Gamma$ , and
- ▷  $\mathcal{M}, \rho \not\Vdash x:D$ , for all  $x:D \in \Delta$ .

Given a sequent  $\mathcal{S} = \mathcal{R}, \Gamma \Rightarrow \Delta$ , and a model  $\mathcal{M} = \langle W, R, v \rangle$ , let  $\text{Lb}(\mathcal{S}) = \{x \mid x \in \mathcal{R} \cup \Gamma \cup \Delta\}$ , and  $\rho : \text{Lb}(\mathcal{S}) \rightarrow W$  (interpretation).

Satisfiability of labelled formulas at  $\mathcal{M}$  under  $\rho$  :

$$\mathcal{M}, \rho \Vdash xRy \quad \text{iff} \quad \mathcal{M} \Vdash \rho(x)R\rho(y)$$

$$\mathcal{M}, \rho \Vdash x:A \quad \text{iff} \quad \mathcal{M}, \rho(x) \Vdash A$$

Satisfiability of sequents at  $\mathcal{M}$  under  $\rho$  ( $\varphi$  is  $xRy$  or  $x:A$ ):

$$\mathcal{M}, \rho \Vdash \mathcal{R}, \Gamma \Rightarrow \Delta \quad \text{iff}$$

*if* for all  $\varphi \in \mathcal{R} \cup \Gamma$  it holds that  $\mathcal{M}, \rho \Vdash \varphi$ ,

*then* for some  $x:D \in \Delta$  it holds that  $\mathcal{M}, \rho \Vdash x:D$ .

A sequent  $\mathcal{R}, \Gamma \Rightarrow \Delta$  has a countermodel iff there are  $\mathcal{M}, \rho$  such that:

- ▷  $\mathcal{M}, \rho \Vdash \varphi$ , for all  $\varphi \in \mathcal{R} \cup \Gamma$ , and
- ▷  $\mathcal{M}, \rho \not\Vdash x:D$ , for all  $x:D \in \Delta$ .

Validity of sequents in a class of frames  $\mathcal{X}$  :

$$\models_{\mathcal{X}} \mathcal{R}, \Gamma \Rightarrow \Delta \quad \text{iff} \quad \text{for any } \rho \text{ and any } \mathcal{M} \in \mathcal{X}, \mathcal{M}, \rho \Vdash \mathcal{R}, \Gamma \Rightarrow \Delta$$

**Theorem (Soundness).** If  $\vdash_{\text{labK}} \mathcal{R}, \Gamma \Rightarrow \Delta$  then  $\models \mathcal{R}, \Gamma \Rightarrow \Delta$





Substitution on labelled formulas:

$$xRy[z/y] \quad := \quad xRz$$

$$y:A[z/y] \quad := \quad z:A$$

Substitution on multisets of labelled formulas  $\Gamma[z/y]$



Substitution on labelled formulas:

$$xRy[z/y] \quad := \quad xRz$$

$$y:A[z/y] \quad := \quad z:A$$

Substitution on multisets of labelled formulas  $\Gamma[z/y]$

**Lemma (Substitution).** Rule subst is hp-admissible.

$$\text{subst} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}[y/x], \Gamma[y/x] \Rightarrow \Delta[y/x]}$$

Substitution on labelled formulas:

$$xRy[z/y] \quad := \quad xRz$$

$$y:A[z/y] \quad := \quad z:A$$

Substitution on multisets of labelled formulas  $\Gamma[z/y]$

**Lemma (Substitution).** Rule subst is hp-admissible.

$$\text{subst} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}[y/x], \Gamma[y/x] \Rightarrow \Delta[y/x]}$$

**Lemma (Weakening).** Rules  $\text{wk}_L$ ,  $\text{wk}_R$  are hp-admissible ( $\varphi$  is  $xRy$  or  $x:A$ ).

$$\text{wk}_L \frac{\mathcal{R}, \Gamma \Rightarrow \Delta}{\varphi, \mathcal{R}, \Gamma \Rightarrow \Delta}$$

$$\text{wk}_R \frac{\mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta, \varphi}$$

**Lemma (Invertibility).**

For every rule  $r$ , if the conclusion of  $r$  is derivable with a derivation of height  $h$ , then each of its premisses is derivable, with at most the same  $h$ .

### Lemma (Invertibility).

For every rule  $r$ , if the conclusion of  $r$  is derivable with a derivation of height  $h$ , then each of its premisses is derivable, with at most the same  $h$ .

Lemma (Contraction). Rules  $\text{ctr}_L$ ,  $\text{ctr}_R$  are hp-admissible ( $\varphi$  is  $xRy$  or  $x:A$ ).

$$\text{ctr}_L \frac{\varphi, \varphi, \mathcal{R}, \Gamma \Rightarrow \Delta}{\varphi, \mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{ctr}_R \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, \varphi, \varphi}{\mathcal{R}, \Gamma \Rightarrow \Delta, \varphi}$$

**Lemma (Cut).** The cut rule is admissible.

$$\text{cut} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad x:A, \mathcal{R}', \Gamma' \Rightarrow \Delta'}{\mathcal{R}, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

**Proof.** By induction on  $(c(A), h_1 + h_2)$ .

**Lemma (Cut).** The cut rule is admissible.

$$\text{cut} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad x:A, \mathcal{R}', \Gamma' \Rightarrow \Delta'}{\mathcal{R}, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

**Proof.** By induction on  $(c(A), h_1 + h_2)$ .

$$\text{cut} \frac{\begin{array}{c} \text{R} \\ \text{R} \end{array} \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, y:A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A} \quad \begin{array}{c} \text{L} \\ \text{L} \end{array} \frac{xRz, \mathcal{R}', x:\Box A, z:A, \Gamma' \Rightarrow \Delta'}{xRz, \mathcal{R}', x:\Box A, \Gamma' \Rightarrow \Delta'}}{\mathcal{R}, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

**Lemma (Cut).** The cut rule is admissible.

$$\text{cut} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad x:A, \mathcal{R}', \Gamma' \Rightarrow \Delta'}{\mathcal{R}, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

**Proof.** By induction on  $(c(A), h_1 + h_2)$ .

$$\text{cut} \frac{\begin{array}{c} \Box_R \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, y:A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A} \\ \Box_L \frac{xRz, \mathcal{R}', x:\Box A, z:A, \Gamma' \Rightarrow \Delta'}{xRz, \mathcal{R}', x:\Box A, \Gamma' \Rightarrow \Delta'} \end{array}}{\mathcal{R}, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

$$\text{cut} \frac{\begin{array}{c} \text{cut} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A \quad xRz, \mathcal{R}', x:\Box A, z:A, \Gamma' \Rightarrow \Delta'}{xRz, \mathcal{R}, \mathcal{R}', z:A, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \\ xRz, \mathcal{R}, \Gamma \Rightarrow \Delta, z:A \end{array}}{\begin{array}{c} \text{ctr}_L, \text{ctr}_R \frac{\mathcal{R}, \mathcal{R}, xRz, xRz, \mathcal{R}', \Gamma, \Gamma, \Gamma' \Rightarrow \Delta, \Delta, \Delta'}{\mathcal{R}, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \end{array}}$$

**Lemma (Cut).** The cut rule is admissible.

$$\text{cut} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad x:A, \mathcal{R}', \Gamma' \Rightarrow \Delta'}{\mathcal{R}, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

**Proof.** By induction on  $(c(A), h_1 + h_2)$ .

$$\begin{array}{c} \frac{\frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta, y:A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A} \quad \frac{xRz, \mathcal{R}', x:\Box A, z:A, \Gamma' \Rightarrow \Delta'}{xRz, \mathcal{R}', x:\Box A, \Gamma' \Rightarrow \Delta'} \quad \text{cut}}{\mathcal{R}, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \end{array}$$

$$\begin{array}{c} \frac{\frac{xRz, \mathcal{R}, \Gamma \Rightarrow \Delta, z:A}{\mathcal{R}, \mathcal{R}, xRz, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \quad \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:\Box A \quad xRz, \mathcal{R}', x:\Box A, z:A, \Gamma' \Rightarrow \Delta'}{xRz, \mathcal{R}, \mathcal{R}', z:A, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \quad \text{cut}}{\mathcal{R}, \mathcal{R}, xRz, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \quad \text{ctr}_L, \text{ctr}_R} \\ \mathcal{R}, xRz, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta' \end{array}$$


For  $\Gamma$  set of formulas and  $x:\Gamma = \{x:G \mid \text{for each } G \in \Gamma\}$ :

**Theorem (Syntactic Completeness).** If  $\Gamma \vdash_K A$  then  $\vdash_{\text{labK}} x:\Gamma \Rightarrow x:A$ .





Frame conditions: a general recipe



What do we mean by modularity?

---

## What do we mean by modularity?

---

Let  $K = \text{CPL} \cup \{k, \text{nec}\}$ . Logic  $K$  is characterised by the class of all Kripke frames.

## What do we mean by modularity?

---

Let  $K = \text{CPL} \cup \{k, \text{nec}\}$ . Logic K is characterised by the class of all Kripke frames.

Name	Axiom	Frame condition	
d	$\Box A \rightarrow \Diamond A$	Seriality	$\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity	$\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry	$\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity	$\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness	$\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

## What do we mean by modularity?

Let  $K = \text{CPL} \cup \{k, \text{nec}\}$ . Logic  $K$  is characterised by the class of all Kripke frames.

Name	Axiom	Frame condition	
d	$\Box A \rightarrow \Diamond A$	Seriality	$\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity	$\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry	$\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity	$\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness	$\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Take  $X \subseteq \{d, t, b, 4, 5\}$ .

## What do we mean by modularity?

Let  $K = \text{CPL} \cup \{k, \text{nec}\}$ . Logic  $K$  is characterised by the class of all Kripke frames.

Name	Axiom	Frame condition
d	$\Box A \rightarrow \Diamond A$	Seriality $\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity $\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry $\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity $\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness $\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Take  $X \subseteq \{d, t, b, 4, 5\}$ .

We write  $\Gamma \vdash_{K \cup X} A$  iff  $A$  is derivable from  $\Gamma$  in the axiom system  $K \cup X$ .

## What do we mean by modularity?

Let  $K = \text{CPL} \cup \{k, \text{nec}\}$ . Logic  $K$  is characterised by the class of all Kripke frames.

Name	Axiom	Frame condition
d	$\Box A \rightarrow \Diamond A$	Seriality $\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity $\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry $\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity $\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness $\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Take  $X \subseteq \{d, t, b, 4, 5\}$ .

We write  $\Gamma \vdash_{K \cup X} A$  iff  $A$  is derivable from  $\Gamma$  in the axiom system  $K \cup X$ .

We denote by  $\mathcal{X}$  the class of frames satisfying properties in  $X$ .

We write  $\Gamma \models_{\mathcal{X}} A$  iff  $A$  is logical consequence of  $\Gamma$  in the class of frames  $\mathcal{X}$ .



## What do we mean by modularity?

Let  $K = \text{CPL} \cup \{k, \text{nec}\}$ . Logic  $K$  is characterised by the class of all Kripke frames.

Name	Axiom	Frame condition
d	$\Box A \rightarrow \Diamond A$	Seriality $\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity $\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry $\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity $\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness $\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Take  $X \subseteq \{d, t, b, 4, 5\}$ .

We write  $\Gamma \vdash_{K \cup X} A$  iff  $A$  is derivable from  $\Gamma$  in the axiom system  $K \cup X$ .

We denote by  $\mathcal{X}$  the class of frames satisfying properties in  $X$ .

We write  $\Gamma \models_{\mathcal{X}} A$  iff  $A$  is logical consequence of  $\Gamma$  in the class of frames  $\mathcal{X}$ .

**Theorem.** For  $X \subseteq \{d, t, b, 4, 5\}$ ,  $\Gamma \vdash_{K \cup X} A$  iff  $\Gamma \models_{\mathcal{X}} A$ .

Name	Axiom	Frame condition	
d	$\Box A \rightarrow \Diamond A$	Seriality	$\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity	$\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry	$\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity	$\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness	$\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Name	Axiom	Frame condition	
d	$\Box A \rightarrow \Diamond A$	Seriality	$\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity	$\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry	$\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity	$\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness	$\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Frame conditions can be characterised by first-order logic formulas, in the language consisting of a single predicate symbol,  $R(x, y)$ .

Name	Axiom	Frame condition	
d	$\Box A \rightarrow \Diamond A$	Seriality	$\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity	$\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry	$\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity	$\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness	$\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Frame conditions can be characterised by first-order logic formulas, in the language consisting of a single predicate symbol,  $R(x, y)$ .

Proof systems for geometric theories, [Negri, 2003]:

“axioms-as-rules”

How to transform axioms of geometric theories (geometric implications) into rules, preserving the structural properties of the calculus.

Name	Axiom	Frame condition	
d	$\Box A \rightarrow \Diamond A$	Seriality	$\forall x \exists y (xRy)$
t	$\Box A \rightarrow A$	Reflexivity	$\forall x (xRx)$
b	$A \rightarrow \Box \Diamond A$	Symmetry	$\forall x \forall y (xRy \rightarrow yRx)$
4	$\Box A \rightarrow \Box \Box A$	Transitivity	$\forall x \forall y \forall z ((xRy \wedge yRz) \rightarrow xRz)$
5	$\Diamond A \rightarrow \Box \Diamond A$	Euclideaness	$\forall x \forall y \forall z ((xRy \wedge xRz) \rightarrow yRz)$

Frame conditions can be characterised by first-order logic formulas, in the language consisting of a single predicate symbol,  $R(x, y)$ .

Proof systems for geometric theories, [Negri, 2003]:

“axioms-as-rules”

How to transform axioms of geometric theories (geometric implications) into rules, preserving the structural properties of the calculus.

The first-order logic formulas corresponding to the frame conditions above (and many more!) are geometric implications



A first-order signature is a tuple  $\sigma = \langle c, d, \dots, f, g, \dots p, q, \dots \rangle$

- ▶ Constant symbols  $c, d, \dots$
- ▶ Function symbols  $f, g, \dots$ , each with arity  $> 0$
- ▶ Predicate symbols  $p, q, \dots$ , each with arity  $\geq 0$

A first-order signature is a tuple  $\sigma = \langle c, d, \dots, f, g, \dots, p, q, \dots \rangle$

- ▶ Constant symbols  $c, d, \dots$
- ▶ Function symbols  $f, g, \dots$ , each with arity  $> 0$
- ▶ Predicate symbols  $p, q, \dots$ , each with arity  $\geq 0$

A first-order language over a signature  $\sigma$ , denoted  $\mathcal{L}(\sigma)$ , consists of:

- ▶ The **terms** generated from a countably many variables  $x, y, \dots$  using the constants and function symbols of  $\sigma$ ;
- ▶ The **formulas** generated from the terms of  $\mathcal{L}(\sigma)$  and predicate symbols of  $\sigma$  using the operators  $\perp, \wedge, \vee, \rightarrow, \forall, \exists$ .



A first-order signature is a tuple  $\sigma = \langle c, d, \dots, f, g, \dots, p, q, \dots \rangle$

- ▶ Constant symbols  $c, d, \dots$
- ▶ Function symbols  $f, g, \dots$ , each with arity  $> 0$
- ▶ Predicate symbols  $p, q, \dots$ , each with arity  $\geq 0$

A first-order language over a signature  $\sigma$ , denoted  $\mathcal{L}(\sigma)$ , consists of:

- ▶ The **terms** generated from a countably many variables  $x, y, \dots$  using the constants and function symbols of  $\sigma$ ;
- ▶ The **formulas** generated from the terms of  $\mathcal{L}(\sigma)$  and predicate symbols of  $\sigma$  using the operators  $\perp, \wedge, \vee, \rightarrow, \forall, \exists$ .

A first-order language with equality over a signature  $\sigma$ , denoted  $\mathcal{L}^=(\sigma)$ , additionally comprises a binary predicate for equality.

A first-order signature is a tuple  $\sigma = \langle c, d, \dots, f, g, \dots, p, q, \dots \rangle$

- ▶ Constant symbols  $c, d, \dots$
- ▶ Function symbols  $f, g, \dots$ , each with arity  $> 0$
- ▶ Predicate symbols  $p, q, \dots$ , each with arity  $\geq 0$

A first-order language over a signature  $\sigma$ , denoted  $\mathcal{L}(\sigma)$ , consists of:

- ▶ The **terms** generated from a countably many variables  $x, y, \dots$  using the constants and function symbols of  $\sigma$ ;
- ▶ The **formulas** generated from the terms of  $\mathcal{L}(\sigma)$  and predicate symbols of  $\sigma$  using the operators  $\perp, \wedge, \vee, \rightarrow, \forall, \exists$ .

A first-order language with equality over a signature  $\sigma$ , denoted  $\mathcal{L}^=(\sigma)$ , additionally comprises a binary predicate for equality.

**Example.**

$\mathcal{L}^=(0, \text{succ}^1, +^2, \times^2)$  is the language of arithmetic

$\mathcal{L}(R^2)$  is the language we use to express frame conditions

Fix a first-order language  $\mathcal{L}(\sigma)$  (with or without equality).

Fix a first-order language  $\mathcal{L}(\sigma)$  (with or without equality).

A **first-order theory** over  $\mathcal{L}(\sigma)$  is a set of closed formulas of  $\mathcal{L}(\sigma)$ .

**Example.** Peano Arithmetic and Robinson Arithmetic are first-order theories over  $\mathcal{L}^=(0, suc, +, \times)$ .

Fix a first-order language  $\mathcal{L}(\sigma)$  (with or without equality).

A **first-order theory** over  $\mathcal{L}(\sigma)$  is a set of closed formulas of  $\mathcal{L}(\sigma)$ .

**Example.** Peano Arithmetic and Robinson Arithmetic are first-order theories over  $\mathcal{L}^=(0, suc, +, \times)$ .

A **geometric formula** is a formula of  $\mathcal{L}(\sigma)$  which does not contain  $\rightarrow$  or  $\forall$ .

Fix a first-order language  $\mathcal{L}(\sigma)$  (with or without equality).

A **first-order theory** over  $\mathcal{L}(\sigma)$  is a set of closed formulas of  $\mathcal{L}(\sigma)$ .

**Example.** Peano Arithmetic and Robinson Arithmetic are first-order theories over  $\mathcal{L}^=(0, suc, +, \times)$ .

A **geometric formula** is a formula of  $\mathcal{L}(\sigma)$  which does not contain  $\rightarrow$  or  $\forall$ .

A **geometric implication** is closed formula of  $\mathcal{L}(\sigma)$  of the shape:

$$\forall \vec{x}(A \rightarrow B), \quad \text{for } A, B \text{ geometric formulas}$$

Fix a first-order language  $\mathcal{L}(\sigma)$  (with or without equality).

A **first-order theory** over  $\mathcal{L}(\sigma)$  is a set of closed formulas of  $\mathcal{L}(\sigma)$ .

**Example.** Peano Arithmetic and Robinson Arithmetic are first-order theories over  $\mathcal{L}^=(0, suc, +, \times)$ .

A **geometric formula** is a formula of  $\mathcal{L}(\sigma)$  which does not contain  $\rightarrow$  or  $\forall$ .

A **geometric implication** is closed formula of  $\mathcal{L}(\sigma)$  of the shape:

$$\forall \vec{x}(A \rightarrow B), \quad \text{for } A, B \text{ geometric formulas}$$

A **geometric theory** over  $\mathcal{L}(\sigma)$  is a first-order theory over  $\mathcal{L}(\sigma)$  whose formulas are geometric implications.

**Example.** Robinson arithmetic is a geometric theory over the language  $\mathcal{L}^=(0, suc, +, \times)$ .

Geometric implications can be expressed as conjunctions of **geometric axioms**, i.e., closed formulas of  $\mathcal{L}(\sigma)$  having the form:

$$\forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \cdots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

- ▶  $\vec{x}, \vec{y}_1, \dots, \vec{y}_m$  are (possibly empty) vectors of variables;
- ▶  $m \geq 0$ ;
- ▶  $P, Q_1, \dots, Q_m$  are (possibly empty) conjunctions of atomic formulas of  $\mathcal{L}(\sigma)$ ;
- ▶  $\vec{y}_1, \dots, \vec{y}_m$  do not occur in  $P$ .



Geometric implications can be expressed as conjunctions of **geometric axioms**, i.e., closed formulas of  $\mathcal{L}(\sigma)$  having the form:

$$\forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \dots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

- ▶  $\vec{x}, \vec{y}_1, \dots, \vec{y}_m$  are (possibly empty) vectors of variables;
- ▶  $m \geq 0$ ;
- ▶  $P, Q_1, \dots, Q_m$  are (possibly empty) conjunctions of atomic formulas of  $\mathcal{L}(\sigma)$ ;
- ▶  $\vec{y}_1, \dots, \vec{y}_m$  do not occur in  $P$ .

Geometric axioms can be turned into sequent calculus rules:

$$\text{GA} \frac{\Xi_1[\vec{z}_1/\vec{y}_1], \Pi, \Gamma \Rightarrow \Delta \quad \dots \quad \Xi_m[\vec{z}_m/\vec{y}_m], \Pi, \Gamma \Rightarrow \Delta}{\Pi, \Gamma \Rightarrow \Delta}$$

- ▶  $\Pi$  is the multiset of atomic formulas in  $P$ ;
- ▶  $\Xi_i$  is the multiset of atomic formulas in  $Q_i$ , for each  $i \leq m$ ;
- ▶  $\vec{z}_1, \dots, \vec{z}_m$  do not occur in  $\Gamma \cup \Delta$ .

Geometric implications can be expressed as conjunctions of **geometric axioms**, i.e., closed formulas of  $\mathcal{L}(\sigma)$  having the form:

$$\forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \dots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

- ▶  $\vec{x}, \vec{y}_1, \dots, \vec{y}_m$  are (possibly empty) vectors of variables;
- ▶  $m \geq 0$ ;
- ▶  $P, Q_1, \dots, Q_m$  are (possibly empty) conjunctions of atomic formulas of  $\mathcal{L}(\sigma)$ ;
- ▶  $\vec{y}_1, \dots, \vec{y}_m$  do not occur in  $P$ .

Geometric axioms can be turned into sequent calculus rules:

$$\text{GA} \frac{\Xi_1[\vec{z}_1/\vec{y}_1], \Pi, \Gamma \Rightarrow \Delta \quad \dots \quad \Xi_m[\vec{z}_m/\vec{y}_m], \Pi, \Gamma \Rightarrow \Delta}{\Pi, \Gamma \Rightarrow \Delta}$$

- ▶  $\Pi$  is the multiset of atomic formulas in  $P$ ;
- ▶  $\Xi_i$  is the multiset of atomic formulas in  $Q_i$ , for each  $i \leq m$ ;
- ▶  $\vec{z}_1, \dots, \vec{z}_m$  do not occur in  $\Gamma \cup \Delta$ .

$$\forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \cdots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

$$\text{GA} \frac{\Xi_1[\vec{z}_1/\vec{y}_1], \Pi, \mathcal{R}, \Gamma \Rightarrow \Delta \quad \cdots \quad \Xi_m[\vec{z}_m/\vec{y}_m], \Pi, \mathcal{R}, \Gamma \Rightarrow \Delta}{\Pi, \mathcal{R}, \Gamma \Rightarrow \Delta}$$

$$\begin{array}{c}
 \text{ser} \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \quad y \text{ fresh} \quad \text{ref} \frac{xRx, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{sym} \frac{yRx, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta} \\
 \\
 \text{tr} \frac{xRz, xRy, yRz, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, yRz, \mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{euc} \frac{yRz, xRy, xRz, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, xRz, \mathcal{R}, \Gamma \Rightarrow \Delta}
 \end{array}$$

$$\begin{array}{c}
 \text{ser} \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \quad y \text{ fresh} \quad \text{ref} \frac{xRx, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{sym} \frac{yRx, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta} \\
 \\
 \text{tr} \frac{xRz, xRy, yRz, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, yRz, \mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{euc} \frac{yRz, xRy, xRz, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, xRz, \mathcal{R}, \Gamma \Rightarrow \Delta}
 \end{array}$$

For  $X \subseteq \{d, t, b, 4, 5\}$ ,  $\text{labK} \cup X$  is defined by adding to  $\text{labK}$  the rules for frame conditions corresponding to elements of  $X$ , plus the rules obtained by to satisfy the **closure condition** (contracted instances of the rules):

$$\text{euc} \frac{yRy, xRy, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta} \rightsquigarrow \text{euc}' \frac{yRy, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}$$

**Example:**  $\text{labK} \cup \{5\}$  denotes the proof system  $\text{labK} \cup \{\text{euc}, \text{euc}'\}$ .

$$\begin{array}{c}
 \text{ser} \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \quad y \text{ fresh} \quad \text{ref} \frac{xRx, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{sym} \frac{yRx, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta} \\
 \\
 \text{tr} \frac{xRz, xRy, yRz, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, yRz, \mathcal{R}, \Gamma \Rightarrow \Delta} \quad \text{euc} \frac{yRz, xRy, xRz, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, xRz, \mathcal{R}, \Gamma \Rightarrow \Delta}
 \end{array}$$

For  $X \subseteq \{d, t, b, 4, 5\}$ ,  $\text{labK} \cup X$  is defined by adding to  $\text{labK}$  the rules for frame conditions corresponding to elements of  $X$ , plus the rules obtained by to satisfy the **closure condition** (contracted instances of the rules):

$$\text{euc} \frac{yRy, xRy, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta} \rightsquigarrow \text{euc}' \frac{yRy, xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}$$

**Example:**  $\text{labK} \cup \{5\}$  denotes the proof system  $\text{labK} \cup \{\text{euc}, \text{euc}'\}$ .

We denote by  $\vdash_{\text{labK} \cup X} S$  derivability of labelled sequent  $S$  in  $\text{labK} \cup X$ .

For  $X \subseteq \{d, t, b, 4, 5\}$ :

**Theorem (Soundness).** If  $\vdash_{\text{labK} \cup X} \mathcal{R}, \Gamma \Rightarrow \Delta$  then  $\models_X \mathcal{R}, \Gamma \Rightarrow \Delta$ .

**Example.** If the premiss of rule *ser* is valid in all serial models, then its conclusion is valid in all serial models.

$$\text{ser} \frac{xRy, \mathcal{R}, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} y \text{ fresh}$$

**Lemma (Cut).** The cut rule is admissible in  $\text{labK} \cup X$ :

$$\text{cut} \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x:A \quad x:A, \mathcal{R}', \Gamma' \Rightarrow \Delta'}{\mathcal{R}, \mathcal{R}', \Gamma, \Gamma' \Rightarrow \Delta, \Delta'}$$

For  $\Gamma$  set of formulas and  $x:\Gamma = \{x:G \mid \text{for each } G \in \Gamma\}$ :

**Theorem (Syntactic Completeness).** If  $\Gamma \vdash_{K \cup X} A$  then  $\vdash_{\text{labK} \cup X} x:\Gamma \Rightarrow x:A$ .







- **Systems of rules** [Negri, 2016], to capture theories / logics characterized by generalized geometric implications:

$$GA_0 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \cdots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

$$GA_1 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_0 \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_0 \right) \right) \right)$$

$$GA_{n+1} = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_{k_1} \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_{k_m} \right) \right) \right)$$

for  $k_1, \dots, k_m \geq n$

- **Systems of rules** [Negri, 2016], to capture theories / logics characterized by generalized geometric implications:

$$GA_0 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \cdots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

$$GA_1 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_0 \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_0 \right) \right) \right)$$

$$GA_{n+1} = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_{k_1} \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_{k_m} \right) \right) \right)$$

for  $k_1, \dots, k_m \geq n$

Systems of rules cover all systems of normal modal logics axiomatised by Sahlqvist formulas.

- ▶ **Systems of rules** [Negri, 2016], to capture theories / logics characterized by generalized geometric implications:

$$GA_0 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \cdots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

$$GA_1 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_0 \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_0 \right) \right) \right)$$

$$GA_{n+1} = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_{k_1} \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_{k_m} \right) \right) \right)$$

for  $k_1, \dots, k_m \geq n$

Systems of rules cover all systems of normal modal logics axiomatised by Sahlqvist formulas.

- ▶ Gödel-Löb provability logic (GL):

Transitivity:  $R$  is transitive

Converse well-foundedness: there are no infinite  $R$ -chains

- **Systems of rules** [Negri, 2016], to capture theories / logics characterized by generalized geometric implications:

$$GA_0 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 (Q_1) \vee \cdots \vee \exists \vec{y}_m (Q_m) \right) \right)$$

$$GA_1 = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_0 \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_0 \right) \right) \right)$$

$$GA_{n+1} = \forall \vec{x} \left( P \rightarrow \left( \exists \vec{y}_1 \left( \bigwedge GA_{k_1} \right) \vee \cdots \vee \exists \vec{y}_m \left( \bigwedge GA_{k_m} \right) \right) \right)$$

for  $k_1, \dots, k_m \geq n$

Systems of rules cover all systems of normal modal logics axiomatised by Sahlqvist formulas.

- Gödel-Löb provability logic (GL):

Transitivity:  $R$  is transitive

Converse well-foundedness: there are no infinite  $R$ -chains

[Negri, 2005]: labelled proof system for GL!

- ▶ Derive axiom 4, that is,  $\Box A \rightarrow \Box \Box A$ , in  $\text{labK} \cup \{t, 5\}$ . Then, show that rule tr is derivable in  $\text{labK} \cup \{t, 5\} \cup \{wk_L, wk_R\}$ .
- ▶ Derive axiom 5, that is,  $\Diamond A \rightarrow \Box \Diamond A$ , in  $\text{labK} \cup \{b, 4\}$ . Then, show that rule euc is derivable in  $\text{labK} \cup \{b, 4\} \cup \{wk_L, wk_R\}$ .
- ▶ Write down the labelled rule corresponding to the frame condition of confluence:

$$\forall x, y, z \left( (xRy \wedge xRz) \rightarrow \exists k (yRk \wedge zRk) \right)$$

- ▶ Write down the sequent calculus rules corresponding to the axioms of Robinson Arithmetic. Can we use the results from [Negri, 2003] to prove consistency of Robinson Arithmetic? If yes, how?