

Proof Theory of Modal Logic

Lecture : Nested Sequents

Tiziano Dalmonte, Marianna Girlando

Free University of Bozen-Bolzano, University of Amsterdam

ESSLLI 2024

Leuven, 5-9 August 2024

Recap

Nested sequents for the S5-cube:
Soundness

Independently introduced in:

- ▶ [Bull, 1992]; [Kashima, 1994] \rightsquigarrow *nested seuquents*
- ▶ [Brünnler, 2006], [Brünnler, 2009] \rightsquigarrow *deep sequents*
- ▶ [Poggiolesi, 2008], [Poggiolesi, 2010] \rightsquigarrow *tree-hypersequents*

Main references for this lecture:

- ▶ [Lellmann & Poggiolesi, 2022 (arXiv)]
- ▶ [Brünnler, 2009], [Brünnler, 2010 (arXiv)]
- ▶ [Marin & Straßburger, 2014]

Sequent

$\Gamma \Rightarrow \Delta$

Γ, Δ multisets of formulas

Sequent

$$\Gamma \Rightarrow \Delta$$

Γ, Δ multisets of formulas

One-sided sequent

$$\Gamma$$

Γ multiset of formulas

Sequent

$$\Gamma \Rightarrow \Delta$$

Γ, Δ multisets of formulas

One-sided sequent

$$\Gamma$$

Γ multiset of formulas

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B$$

Sequent

$$\Gamma \Rightarrow \Delta$$

Γ, Δ multisets of formulas

One-sided sequent

$$\Gamma$$

Γ multiset of formulas

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B}$$

Sequent

$$\Gamma \Rightarrow \Delta$$

Γ, Δ multisets of formulas

One-sided sequent

$$\Gamma$$

Γ multiset of formulas

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Sequent	$\Gamma \Rightarrow \Delta$	Γ, Δ multisets of formulas
One-sided sequent	Γ	Γ multiset of formulas

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Rules of $G3cp^{one}$

$$\text{init} \frac{}{\Gamma, p, \bar{p}} \quad \wedge \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \quad \vee \frac{\Gamma, A, B}{\Gamma, A \vee B}$$

Sequent	$\Gamma \Rightarrow \Delta$	Γ, Δ multisets of formulas
One-sided sequent	Γ	Γ multiset of formulas

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Rules of $G3cp^{one}$

$$\text{init} \frac{}{\Gamma, p, \bar{p}} \quad \wedge \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \quad \vee \frac{\Gamma, A, B}{\Gamma, A \vee B}$$

Exercise. $\vdash_{G3cp} \Gamma \Rightarrow \Delta$ iff $\vdash_{G3cp^{one}} \bar{\Gamma}, \Delta$, where $\bar{\Gamma} = \{\bar{A} \mid A \in \Gamma\}$.

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B} \quad \overline{\Box A} := \Diamond \bar{A} \quad \overline{\Diamond A} := \Box \bar{A}$$

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\begin{aligned} \overline{A \wedge B} &:= \bar{A} \vee \bar{B} & \overline{A \vee B} &:= \bar{A} \wedge \bar{B} & \overline{\Box A} &:= \Diamond \bar{A} & \overline{\Diamond A} &:= \Box \bar{A} \\ A \rightarrow B &:= \bar{A} \vee B & \perp &:= p \wedge \bar{p} \end{aligned}$$

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B} \quad \overline{\Box A} := \Diamond \bar{A} \quad \overline{\Diamond A} := \Box \bar{A}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Nested sequents (denoted Γ, Δ, \dots) are inductively generated as follows:

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B} \quad \overline{\Box A} := \Diamond \bar{A} \quad \overline{\Diamond A} := \Box \bar{A}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Nested sequents (denoted Γ, Δ, \dots) are inductively generated as follows:

- ▶ A multiset of formulas is a nested sequent;

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B} \quad \overline{\Box A} := \Diamond \bar{A} \quad \overline{\Diamond A} := \Box \bar{A}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Nested sequents (denoted Γ, Δ, \dots) are inductively generated as follows:

- ▶ A multiset of formulas is a nested sequent;
- ▶ If Γ and Δ are nested sequents, then Γ, Δ is a nested sequent;

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B} \quad \overline{\Box A} := \Diamond \bar{A} \quad \overline{\Diamond A} := \Box \bar{A}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Nested sequents (denoted Γ, Δ, \dots) are inductively generated as follows:

- ▶ A multiset of formulas is a nested sequent;
- ▶ If Γ and Δ are nested sequents, then Γ, Δ is a nested sequent;
- ▶ If Γ is a nested sequent, then $[\Gamma]$ is a nested sequent.

We call $[\Gamma]$ a **boxed sequent**.

$$A, B ::= p \mid \bar{p} \mid A \wedge B \mid A \vee B \mid \Box A \mid \Diamond A$$

$$\overline{A \wedge B} := \bar{A} \vee \bar{B} \quad \overline{A \vee B} := \bar{A} \wedge \bar{B} \quad \overline{\Box A} := \Diamond \bar{A} \quad \overline{\Diamond A} := \Box \bar{A}$$

$$A \rightarrow B := \bar{A} \vee B \quad \perp := p \wedge \bar{p}$$

Nested sequents (denoted Γ, Δ, \dots) are inductively generated as follows:

- ▶ A multiset of formulas is a nested sequent;
- ▶ If Γ and Δ are nested sequents, then Γ, Δ is a nested sequent;
- ▶ If Γ is a nested sequent, then $[\Gamma]$ is a nested sequent.

We call $[\Gamma]$ a **boxed sequent**.

Nested sequents are multisets of formulas and boxed sequents:

$$A_1, \dots, A_m, [\Delta_1], \dots, [\Delta_n]$$

$$\Gamma = A_1, \dots, A_m, [\Delta_1], \dots, [\Delta_n]$$

To a nested sequent Γ there corresponds the following tree $tr(\Gamma)$, whose nodes γ, δ, \dots are multisets of formulas:

The formula interpretation $i(\Gamma)$ of a nested sequent Γ is defined as:

- ▶ If $m = n = 0$, then $i(\Gamma) := \perp$
- ▶ Otherwise, $i(\Gamma) := A_1 \vee \dots \vee A_m \vee \Box(i(\Delta_1)) \vee \dots \vee \Box(i(\Delta_n))$

A **context** is a nested sequent with one or multiple holes, denoted by $\{\}$, each taking the place of a formula in the nested sequent.

- ▶ Unary context $\Gamma\{\}$
- ▶ Binary context $\Gamma\{\}\{\}$

The **depth** $depth(\Gamma\{\})$ of a unary context $\Gamma\{\}$ is defined as:

- ▶ $depth(\{\}) := 0$;
- ▶ $depth(\Gamma\{\}, \Delta) := depth(\Gamma\{\})$;
- ▶ $depth([\Gamma\{\}]) := depth(\Gamma\{\}) + 1$.

A **context** is a nested sequent with one or multiple holes, denoted by $\{\}$, each taking the place of a formula in the nested sequent.

- ▶ Unary context $\Gamma\{\}$ \rightsquigarrow $\Gamma\{\Delta\}$: filling $\Gamma\{\}$ with a nested sequent Δ
- ▶ Binary context $\Gamma\{\}\{\}$ \rightsquigarrow $\Gamma\{\Delta_1\}\{\Delta_2\}$: filling $\Gamma\{\}\{\}$ with Δ_1, Δ_2

The **depth** $depth(\Gamma\{\})$ of a unary context $\Gamma\{\}$ is defined as:

- ▶ $depth(\{\}) := 0$;
- ▶ $depth(\Gamma\{\}, \Delta) := depth(\Gamma\{\})$;
- ▶ $depth([\Gamma\{\}]) := depth(\Gamma\{\}) + 1$.

$$\begin{array}{c}
 \text{init} \frac{}{\Gamma\{p, \bar{p}\}} \quad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \quad \vee \frac{\Gamma\{A, B\}}{\Gamma\{A \vee B\}} \\
 \\
 \Box \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}} \quad \Diamond \frac{\Gamma\{\Diamond A, [A, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}}
 \end{array}$$

$$\begin{array}{c}
 \text{init} \frac{}{\Gamma\{p, \bar{p}\}} \quad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \quad \vee \frac{\Gamma\{A, B\}}{\Gamma\{A \vee B\}} \\
 \\
 \Box \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}} \quad \Diamond \frac{\Gamma\{\Diamond A, [A, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}}
 \end{array}$$

Example. Proof of $(\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)$ in NK

$$\begin{array}{c}
 \text{init} \frac{}{\Diamond p, [p, \bar{p}, q]} \quad \text{init} \frac{}{\Diamond \bar{q}, [\bar{q}, \bar{p}, q]} \\
 \Diamond \frac{}{\Diamond p, [\bar{p}, q]} \quad \Diamond \frac{}{\Diamond \bar{q}, [\bar{p}, q]} \\
 \wedge \frac{}{\Diamond p \wedge \Diamond \bar{q}, [\bar{p}, q]} \\
 \vee \frac{}{\Diamond p \wedge \Diamond \bar{q}, [\bar{p} \vee q]} \\
 \Box \frac{}{\Diamond p \wedge \Diamond \bar{q}, \Box(\bar{p} \vee q)} \\
 \vee \frac{}{(\Diamond p \wedge \Diamond \bar{q}) \vee \Box(\bar{p} \vee q)}
 \end{array}$$

Roadmap

For a nested sequent Γ and a model $\mathcal{M} = \langle W, R, v \rangle$, an \mathcal{M} -map for Γ is a map $f : tr(\Gamma) \rightarrow W$ such that whenever δ is a child of γ in $tr(\Gamma)$, then $f(\gamma)Rf(\delta)$.

For a nested sequent Γ and a model $\mathcal{M} = \langle W, R, v \rangle$, an **\mathcal{M} -map for Γ** is a map $f : tr(\Gamma) \rightarrow W$ such that whenever δ is a child of γ in $tr(\Gamma)$, then $f(\gamma)Rf(\delta)$.

A nested sequent Γ is **satisfied** by an \mathcal{M} -map for Γ iff

$$\mathcal{M}, f(\delta) \models B, \text{ for some } \delta \in tr(\Gamma), \text{ for some } B \in \delta$$

For a nested sequent Γ and a model $\mathcal{M} = \langle W, R, v \rangle$, an **\mathcal{M} -map for Γ** is a map $f : tr(\Gamma) \rightarrow W$ such that whenever δ is a child of γ in $tr(\Gamma)$, then $f(\gamma)Rf(\delta)$.

A nested sequent Γ is **satisfied** by an \mathcal{M} -map for Γ iff

$$\mathcal{M}, f(\delta) \models B, \text{ for some } \delta \in tr(\Gamma), \text{ for some } B \in \delta$$

A nested sequent Γ is **refuted** by an \mathcal{M} -map for Γ iff

$$\mathcal{M}, f(\delta) \not\models B, \text{ for all } \delta \in tr(\Gamma), \text{ for all } B \in \delta$$

For a nested sequent Γ and a model $\mathcal{M} = \langle W, R, v \rangle$, an **\mathcal{M} -map for Γ** is a map $f : tr(\Gamma) \rightarrow W$ such that whenever δ is a child of γ in $tr(\Gamma)$, then $f(\gamma)Rf(\delta)$.

A nested sequent Γ is **satisfied** by an \mathcal{M} -map for Γ iff

$$\mathcal{M}, f(\delta) \models B, \text{ for some } \delta \in tr(\Gamma), \text{ for some } B \in \delta$$

A nested sequent Γ is **refuted** by an \mathcal{M} -map for Γ iff

$$\mathcal{M}, f(\delta) \not\models B, \text{ for all } \delta \in tr(\Gamma), \text{ for all } B \in \delta$$

For $X \subseteq \{d, t, b, 4, 5\}$, a nested sequent is **X-valid** iff it is satisfied by all \mathcal{M} -map for Γ , for all models \mathcal{M} satisfying the frame conditions in X .

Lemma. If Γ is derivable in NK then $\bigvee \Gamma$ is valid in all Kripke frames.

$$\begin{array}{ccc}
 d^\diamond \frac{\Gamma\{\diamond A, [A]\}}{\Gamma\{\diamond A\}} & t^\diamond \frac{\Gamma\{\diamond A, A\}}{\Gamma\{\diamond A\}} & b^\diamond \frac{\Gamma\{[\Delta, \diamond A], A\}}{\Gamma\{[\Delta, \diamond A]\}} \\
 4^\diamond \frac{\Gamma\{\diamond A, [\diamond A, \Delta]\}}{\Gamma\{\diamond A, [\Delta]\}} & 5^\diamond \frac{\Gamma\{\diamond A\}\{\diamond A\}}{\Gamma\{\diamond A\}\{\emptyset\}} & \text{depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

$$\begin{array}{c}
 d^\diamond \frac{\Gamma\{\diamond A, [A]\}}{\Gamma\{\diamond A\}} \quad t^\diamond \frac{\Gamma\{\diamond A, A\}}{\Gamma\{\diamond A\}} \quad b^\diamond \frac{\Gamma\{[\Delta, \diamond A], A\}}{\Gamma\{[\Delta, \diamond A]\}} \\
 4^\diamond \frac{\Gamma\{\diamond A, [\diamond A, \Delta]\}}{\Gamma\{\diamond A, [\Delta]\}} \quad 5^\diamond \frac{\Gamma\{\diamond A\}\{\diamond A\}}{\Gamma\{\diamond A\}\{\emptyset\}} \text{ } depth(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write X^\diamond for the corresponding subset of $\{d^\diamond, t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\}$. We shall consider the calculi $NK \cup X^\diamond$.

$$\begin{array}{c}
 d^\diamond \frac{\Gamma\{\diamond A, [A]\}}{\Gamma\{\diamond A\}} \quad t^\diamond \frac{\Gamma\{\diamond A, A\}}{\Gamma\{\diamond A\}} \quad b^\diamond \frac{\Gamma\{[\Delta, \diamond A], A\}}{\Gamma\{[\Delta, \diamond A]\}} \\
 4^\diamond \frac{\Gamma\{\diamond A, [\diamond A, \Delta]\}}{\Gamma\{\diamond A, [\Delta]\}} \quad 5^\diamond \frac{\Gamma\{\diamond A\}\{\diamond A\}}{\Gamma\{\diamond A\}\{\emptyset\}} \text{ depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write X^\diamond for the corresponding subset of $\{d^\diamond, t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\}$. We shall consider the calculi $NK \cup X^\diamond$.

Example. Proof of $\Box p \rightarrow \Box\Box p$ in $NK \cup \{t, 4\}$

$$\begin{array}{c}
 \text{init} \frac{}{\diamond \bar{p}, [\diamond \bar{p}, [\diamond \bar{p}, \bar{p}, p]]} \\
 t^\diamond \frac{}{\diamond \bar{p}, [\diamond \bar{p}, [\diamond \bar{p}, p]]} \\
 4^\diamond \frac{}{\diamond \bar{p}, [\diamond \bar{p}, [p]]} \\
 4^\diamond \frac{}{\diamond \bar{p}, [[p]]} \\
 \Box \frac{}{\diamond \bar{p}, [\Box p]} \\
 \Box \frac{}{\diamond \bar{p}, \Box\Box p} \\
 \vee \frac{}{\diamond \bar{p} \vee \Box\Box p}
 \end{array}$$

$$\text{wk} \frac{\Gamma\{\emptyset\}}{\Gamma\{\Delta\}}$$

$$\text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}}$$

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

$$\text{wk} \frac{\Gamma\{\emptyset\}}{\Gamma\{\Delta\}}$$

$$\text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}}$$

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

For $X \subseteq \{d, t, b, 4, 5\}$:

Lemma. The rules wk and ctr are hp-admissible in $NK \cup X^\diamond$.

$$\text{wk} \frac{\Gamma\{\emptyset\}}{\Gamma\{\Delta\}}$$

$$\text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}}$$

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

For $X \subseteq \{d, t, b, 4, 5\}$:

Lemma. The rules wk and ctr are hp-admissible in $NK \cup X^\diamond$.

Lemma. All the rules of $NK \cup X^\diamond$ are hp-invertible.

$$\text{wk} \frac{\Gamma\{\emptyset\}}{\Gamma\{\Delta\}}$$

$$\text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}}$$

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

For $X \subseteq \{d, t, b, 4, 5\}$:

Lemma. The rules wk and ctr are hp-admissible in $\text{NK} \cup X^\diamond$.

Lemma. All the rules of $\text{NK} \cup X^\diamond$ are hp-invertible.

Proposition. Rule 5^\diamond is derivable in $\text{NK} \cup \{5_1^\diamond, 5_2^\diamond, 5_3^\diamond\} \cup \{\text{wk}\}$.

$$5^\diamond \frac{\Gamma\{\Diamond A\}\{\Diamond A\}}{\Gamma\{\Diamond A\}\{\emptyset\}} \text{depth}(\Gamma\{\}\{\emptyset\}) > 0$$

$$5_1^\diamond \frac{\Gamma\{[\Delta, \Diamond A], \Diamond A\}}{\Gamma\{[\Delta, \Diamond A]\}} \quad 5_2^\diamond \frac{\Gamma\{[\Delta, \Diamond A], [\Lambda, \Diamond A]\}}{\Gamma\{[\Delta, \Diamond A], [\Lambda]\}} \quad 5_3^\diamond \frac{[\Delta, \Diamond A, [\Lambda, \Diamond A]]}{\Gamma\{[\Delta, \Diamond A, [\Lambda]]\}}$$

$$\text{wk} \frac{\Gamma\{\emptyset\}}{\Gamma\{\Delta\}}$$

$$\text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}}$$

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

For $X \subseteq \{d, t, b, 4, 5\}$:

Lemma. The rules wk and ctr are hp-admissible in $NK \cup X^\diamond$.

Lemma. All the rules of $NK \cup X^\diamond$ are hp-invertible.

Proposition. Rule 5^\diamond is derivable in $NK \cup \{5_1^\diamond, 5_2^\diamond, 5_3^\diamond\} \cup \{\text{wk}\}$.

$$5^\diamond \frac{\Gamma\{\Diamond A\}\{\Diamond A\}}{\Gamma\{\Diamond A\}\{\emptyset\}} \text{depth}(\Gamma\{\}\{\emptyset\}) > 0$$

$$5_1^\diamond \frac{\Gamma\{[\Delta, \Diamond A], \Diamond A\}}{\Gamma\{[\Delta, \Diamond A]\}} \quad 5_2^\diamond \frac{\Gamma\{[\Delta, \Diamond A], [\wedge, \Diamond A]\}}{\Gamma\{[\Delta, \Diamond A], [\wedge]\}} \quad 5_3^\diamond \frac{[\Delta, \Diamond A, [\wedge, \Diamond A]]}{\Gamma\{[\Delta, \Diamond A, [\wedge]]\}}$$

Lemma. If Γ is derivable in $NK \cup X^\diamond$ then $\forall \Gamma$ is valid in all X-frames.

Rules of NK_{ctr}

$$\begin{array}{c}
 \text{init} \frac{}{\Gamma\{p, \bar{p}\}} \quad \text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}} \quad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \quad \vee \frac{\Gamma\{A, B\}}{\Gamma\{A \vee B\}} \\
 \\
 \square \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}} \quad \diamond_{ctr} \frac{\Gamma\{\cancel{\Diamond A}, [A, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}}
 \end{array}$$

Rules for extensions

$$\begin{array}{c}
 d_{ctr}^{\diamond} \frac{\Gamma\{\cancel{\Diamond A}, [A]\}}{\Gamma\{\Diamond A\}} \quad t_{ctr}^{\diamond} \frac{\Gamma\{\cancel{\Diamond A}, A\}}{\Gamma\{\Diamond A\}} \quad b_{ctr}^{\diamond} \frac{\Gamma\{[\Delta, \cancel{\Diamond A}], A\}}{\Gamma\{[\Delta, \Diamond A]\}} \\
 \\
 4_{ctr}^{\diamond} \frac{\Gamma\{\Diamond A, [\cancel{\Diamond A}, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}} \quad 5_{ctr}^{\diamond} \frac{\Gamma\{\cancel{\Diamond A}\}\{\Diamond A\}}{\Gamma\{\Diamond A\}\{\emptyset\}} \quad \text{depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write X_{ctr}^{\diamond} for the corresponding subset of $\{d_{ctr}^{\diamond}, t_{ctr}^{\diamond}, b_{ctr}^{\diamond}, 4_{ctr}^{\diamond}, 5_{ctr}^{\diamond}\}$.

Rules of NK_{ctr}

$$\begin{array}{c}
 \text{init} \frac{}{\Gamma\{p, \bar{p}\}} \quad \text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}} \quad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \quad \vee \frac{\Gamma\{A, B\}}{\Gamma\{A \vee B\}} \\
 \\
 \Box \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}} \quad \Diamond_{ctr} \frac{\Gamma\{\cancel{\Diamond A}, [A, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}}
 \end{array}$$

Rules for extensions

$$\begin{array}{c}
 d_{ctr}^{\Diamond} \frac{\Gamma\{\cancel{\Diamond A}, [A]\}}{\Gamma\{\Diamond A\}} \quad t_{ctr}^{\Diamond} \frac{\Gamma\{\cancel{\Diamond A}, A\}}{\Gamma\{\Diamond A\}} \quad b_{ctr}^{\Diamond} \frac{\Gamma\{[\Delta, \cancel{\Diamond A}], A\}}{\Gamma\{[\Delta, \Diamond A]\}} \\
 \\
 4_{ctr}^{\Diamond} \frac{\Gamma\{\Diamond A, [\cancel{\Diamond A}, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}} \quad 5_{ctr}^{\Diamond} \frac{\Gamma\{\cancel{\Diamond A}\}\{\Diamond A\}}{\Gamma\{\Diamond A\}\{\emptyset\}} \quad \text{depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write X_{ctr}^{\Diamond} for the corresponding subset of $\{d_{ctr}^{\Diamond}, t_{ctr}^{\Diamond}, b_{ctr}^{\Diamond}, 4_{ctr}^{\Diamond}, 5_{ctr}^{\Diamond}\}$.

Lemma. The rule wk is hp-admissible in $NK \cup X^{\Diamond}$.

Rules of NK_{ctr}

$$\begin{array}{c}
 \text{init} \frac{}{\Gamma\{p, \bar{p}\}} \quad \text{ctr} \frac{\Gamma\{\Delta, \Delta\}}{\Gamma\{\Delta\}} \quad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \quad \vee \frac{\Gamma\{A, B\}}{\Gamma\{A \vee B\}} \\
 \\
 \Box \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}} \quad \Diamond_{ctr} \frac{\Gamma\{\cancel{\Diamond A}, [A, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}}
 \end{array}$$

Rules for extensions

$$\begin{array}{c}
 d_{ctr}^{\Diamond} \frac{\Gamma\{\cancel{\Diamond A}, [A]\}}{\Gamma\{\Diamond A\}} \quad t_{ctr}^{\Diamond} \frac{\Gamma\{\cancel{\Diamond A}, A\}}{\Gamma\{\Diamond A\}} \quad b_{ctr}^{\Diamond} \frac{\Gamma\{[\Delta, \cancel{\Diamond A}], A\}}{\Gamma\{[\Delta, \Diamond A]\}} \\
 \\
 4_{ctr}^{\Diamond} \frac{\Gamma\{\Diamond A, [\cancel{\Diamond A}, \Delta]\}}{\Gamma\{\Diamond A, [\Delta]\}} \quad 5_{ctr}^{\Diamond} \frac{\Gamma\{\cancel{\Diamond A}\}\{\Diamond A\}}{\Gamma\{\Diamond A\}\{\emptyset\}} \quad \text{depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write X_{ctr}^{\Diamond} for the corresponding subset of $\{d_{ctr}^{\Diamond}, t_{ctr}^{\Diamond}, b_{ctr}^{\Diamond}, 4_{ctr}^{\Diamond}, 5_{ctr}^{\Diamond}\}$.

Lemma. The rule wk is hp-admissible in $NK \cup X^{\Diamond}$.

Proposition. Γ is derivable in $NK \cup X^{\Diamond}$ iff Γ is derivable in $NK_{ctr} \cup X_{ctr}^{\Diamond}$.

Nested sequents for the S5-cube:
Completeness

Three problems for completeness

- ▶ Axiom 5, that is, $\Diamond A \rightarrow \Box \Diamond A$, is valid in all $\{b, 4\}$ -frames, but it is **not** derivable in $NK \cup \{b^\Diamond, 4^\Diamond\}$.

- ▶ Axiom 5, that is, $\Diamond A \rightarrow \Box \Diamond A$, is valid in all $\{b, 4\}$ -frames, but it is **not** derivable in $NK \cup \{b^\Diamond, 4^\Diamond\}$.

Failed proof of $\Diamond A \rightarrow \Box \Diamond A$ in $NK \cup \{b^\Diamond, 4^\Diamond\}$

$$\begin{array}{c} b^\Diamond \frac{[\bar{p}], p, [\Diamond p]}{[\bar{p}], [\Diamond p]} \\ \square \frac{[\bar{p}], [\Diamond p]}{\Box \bar{p}, [\Diamond p]} \\ \square \frac{\Box \bar{p}, [\Diamond p]}{\Box \bar{p}, \Box \Diamond p} \\ \vee \frac{\Box \bar{p}, \Box \Diamond p}{\Box \bar{p} \vee \Box \Diamond p} \end{array}$$

- ▶ Axiom 5, that is, $\Diamond A \rightarrow \Box \Diamond A$, is valid in all $\{b, 4\}$ -frames, but it is **not** derivable in $NK \cup \{b^\Diamond, 4^\Diamond\}$.
- ▶ Axiom 4, that is, $A \rightarrow \Box \Box A$, is valid in all $\{t, 5\}$ -frames, but it is **not** derivable in $NK \cup \{t^\Diamond, 5^\Diamond\}$.

Failed proof of $\Diamond A \rightarrow \Box \Diamond A$ in $NK \cup \{b^\Diamond, 4^\Diamond\}$

$$\begin{array}{c} b^\Diamond \frac{[\bar{p}], p, [\Diamond p]}{[\bar{p}], [\Diamond p]} \\ \quad \Box \frac{[\bar{p}], [\Diamond p]}{\Box \bar{p}, [\Diamond p]} \\ \quad \quad \Box \frac{\Box \bar{p}, [\Diamond p]}{\Box \bar{p}, \Box \Diamond p} \\ \quad \quad \vee \frac{\Box \bar{p}, \Box \Diamond p}{\Box \bar{p} \vee \Box \Diamond p} \end{array}$$

- ▶ Axiom 5, that is, $\Diamond A \rightarrow \Box \Diamond A$, is valid in all $\{b, 4\}$ -frames, but it is **not** derivable in $NK \cup \{b^\Diamond, 4^\Diamond\}$.
- ▶ Axiom 4, that is, $A \rightarrow \Box \Box A$, is valid in all $\{t, 5\}$ -frames, but it is **not** derivable in $NK \cup \{t^\Diamond, 5^\Diamond\}$.
- ▶ Axiom 4, that is, $A \rightarrow \Box \Box A$, is valid in all $\{b, 5\}$ -frames, but it is **not** derivable in $NK \cup \{b^\Diamond, 5^\Diamond\}$.

Failed proof of $\Diamond A \rightarrow \Box \Diamond A$ in $NK \cup \{b^\Diamond, 4^\Diamond\}$

$$\begin{array}{c} b^\Diamond \frac{[\bar{p}], p, [\Diamond p]}{[\bar{p}], [\Diamond p]} \\ \Box \frac{[\bar{p}], [\Diamond p]}{\Box \bar{p}, [\Diamond p]} \\ \Box \frac{\Box \bar{p}, [\Diamond p]}{\Box \bar{p}, \Box \Diamond p} \\ \vee \frac{\Box \bar{p}, \Box \Diamond p}{\Box \bar{p} \vee \Box \Diamond p} \end{array}$$

For each set of frames characterised by the 5-axioms, there is at least one combination of modal rules which is complete.

For each set of frames characterised by the 5-axioms, there is at least one combination of modal rules which is complete.

For $X \subseteq \{d, t, b, 4, 5\}$, the **45-closure** of X is defined as:

$$\hat{X} = \begin{cases} X \cup \{4\} & \text{if } \{b, 5\} \subseteq X \text{ or } \{t, 5\} \subseteq X \\ X \cup \{5\} & \text{if } \{b, 4\} \subseteq X \\ X & \text{otherwise} \end{cases}$$

We say that X is **45-closed** if $X = \hat{X}$.

For each set of frames characterised by the 5-axioms, there is at least one combination of modal rules which is complete.

For $X \subseteq \{d, t, b, 4, 5\}$, the **45-closure** of X is defined as:

$$\hat{X} = \begin{cases} X \cup \{4\} & \text{if } \{b, 5\} \subseteq X \text{ or } \{t, 5\} \subseteq X \\ X \cup \{5\} & \text{if } \{b, 4\} \subseteq X \\ X & \text{otherwise} \end{cases}$$

We say that X is **45-closed** if $X = \hat{X}$.

Proposition. For $X \subseteq \{d, t, b, 4, 5\}$ X is 45-closed iff, for $\rho \in \{4, 5\}$, it holds that if ρ is valid in all X -frames, then $\rho \in X$.

For each set of frames characterised by the 5-axioms, there is at least one combination of modal rules which is complete.

For $X \subseteq \{d, t, b, 4, 5\}$, the **45-closure** of X is defined as:

$$\hat{X} = \begin{cases} X \cup \{4\} & \text{if } \{b, 5\} \subseteq X \text{ or } \{t, 5\} \subseteq X \\ X \cup \{5\} & \text{if } \{b, 4\} \subseteq X \\ X & \text{otherwise} \end{cases}$$

We say that X is **45-closed** if $X = \hat{X}$.

Proposition. For $X \subseteq \{d, t, b, 4, 5\}$ X is 45-closed iff, for $\rho \in \{4, 5\}$, it holds that if ρ is valid in all X -frames, then $\rho \in X$.

To prove:

Theorem (Completeness). For $X \subseteq \{d, t, b, 4, 5\}$, if Γ is X -valid, then Γ is derivable in $NK \cup \hat{X}^\diamond$.

Theorem (Completeness). For $X \subseteq \{d, t, b, 4, 5\}$, if Γ is X -valid, then Γ is derivable in $NK \cup \hat{X}^\diamond$.

Theorem (Cut-elimination). For $X \subseteq \{d, t, b, 4, 5\}$ 45-closed, if Γ is derivable in $NK \cup X^\diamond \cup \{\text{cut}\}$, then it is derivable in $NK \cup X^\diamond$.

Theorem (Cut-elimination). For $X \subseteq \{d, t, b, 4, 5\}$ 45-closed, if Γ is derivable in $NK \cup X^\diamond \cup \{\text{cut}\}$, then it is derivable in $NK \cup X^\diamond$.

The proof uses:

- ▶ A generalised version of cut (eliminable)

$$\text{cut} \frac{\frac{\square \frac{\Gamma\{[A], [\Delta]\}}{\Gamma\{\square A, [\Delta]\}} \quad \text{tr}^\diamond \frac{\Gamma\{\diamond \bar{A}, [\diamond \bar{A}, \Delta]\}}{\Gamma\{\diamond \bar{A}, [\Delta]\}}}{\Gamma\{[\Delta]\}}}$$

- ▶ Additional structural modal rules (admissible)

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

$$\text{Y-cut} \frac{\Gamma\{\Box A\}\{\emptyset\}^n \quad \Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n}{\Gamma\{\emptyset\}\{\emptyset\}^n}$$

In the Y-cut:

- ▶ $\{\Delta\}^n$ denotes $\overbrace{\{\Delta\} \dots \{\Delta\}}^{n \text{ times}}$;
- ▶ $n \geq 0$;
- ▶ $Y \subseteq \{4, 5\}$;
- ▶ there is a derivation of $\Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n$ to $\Gamma\{\Diamond \bar{A}\}\{\emptyset\}^n$ in system Y^\Diamond .

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

$$\text{Y-cut} \frac{\Gamma\{\Box A\}\{\emptyset\}^n \quad \Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n}{\Gamma\{\emptyset\}\{\emptyset\}^n}$$

In the Y-cut:

- ▶ $\{\Delta\}^n$ denotes $\overbrace{\{\Delta\} \dots \{\Delta\}}^{n \text{ times}}$;
- ▶ $n \geq 0$;
- ▶ $Y \subseteq \{4, 5\}$;
- ▶ there is a derivation of $\Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n$ to $\Gamma\{\Diamond \bar{A}\}\{\emptyset\}^n$ in system Y^\Diamond .

The rank of the cut formula A is defined as the complexity of A , plus one.
 The **cut rank** of a derivation is the maximum of the ranks of its cuts.

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}}$$

$$\text{Y-cut} \frac{\Gamma\{\Box A\}\{\emptyset\}^n \quad \Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n}{\Gamma\{\emptyset\}\{\emptyset\}^n}$$

In the Y-cut:

- ▶ $\{\Delta\}^n$ denotes $\overbrace{\{\Delta\} \dots \{\Delta\}}^{n \text{ times}}$;
- ▶ $n \geq 0$;
- ▶ $Y \subseteq \{4, 5\}$;
- ▶ there is a derivation of $\Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n$ to $\Gamma\{\Diamond \bar{A}\}\{\emptyset\}^n$ in system Y^\Diamond .

The rank of the cut formula A is defined as the complexity of A , plus one.
The **cut rank** of a derivation is the maximum of the ranks of its cuts.

The notions of cut rank-preserving admissible rule and cut rank-preserving invertible rule are defined analogously to the notions of hp admissible rule and hp invertible rule.

$$\text{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\emptyset\}} \qquad \text{Y-cut} \frac{\Gamma\{\Box A\}\{\emptyset\}^n \quad \Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n}{\Gamma\{\emptyset\}\{\emptyset\}^n}$$

If $Y = \{4\}$, then $\Gamma\{\}\{\}^n$ is of the form $\Gamma_1\{\}, \Gamma_2\{\}^n$:

$$4\text{-cut} \frac{\Gamma_1\{\Box A\}, \Gamma_2\{\emptyset\}^n \quad \Gamma_1\{\Diamond A\}, \Gamma_2\{\Diamond A\}^n}{\Gamma_1\{\emptyset\}, \Gamma_2\{\emptyset\}^n}$$

$$\begin{array}{c} \Box \\ \text{cut} \end{array} \frac{\Gamma\{[A], [\Delta]\}}{\Gamma\{\Box A, [\Delta]\}} \quad 4^\Diamond \frac{\Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, \Delta]\}}{\Gamma\{\Diamond \bar{A}, [\Delta]\}} \quad \rightsquigarrow \quad \begin{array}{c} \Box \\ 4\text{-cut} \end{array} \frac{\Gamma\{[A], [\Delta]\}}{\Gamma\{\Box A, [\Delta]\}} \quad \Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, \Delta]\}}{\Gamma\{[\Delta]\}}$$

$$\begin{array}{ccc}
 d^{[]} \frac{\Gamma\{\emptyset\}}{\Gamma\{\emptyset\}} & t^{[]} \frac{\Gamma\{[\Delta]\}}{\Gamma\{\Delta\}} & b^{[]} \frac{\Gamma\{[\Sigma, [\Delta]]\}}{\Gamma\{\Delta, [\Sigma]\}} \\
 4^{[]} \frac{\Gamma\{[\Delta], [\Sigma]\}}{\Gamma\{[[\Delta], \Sigma]\}} & 5^{[]} \frac{\Gamma\{\Delta\}\{\emptyset\}}{\Gamma\{\emptyset\}\{\Delta\}} & \text{depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write $X^{[]}$ for the corresponding subset of $\{d^{[]}, t^{[]}, b^{[]}, 4^{[]}, 5^{[]}\}$.

$$\begin{array}{c}
 d^{[]} \frac{\Gamma\{\emptyset\}}{\Gamma\{\emptyset\}} \quad t^{[]} \frac{\Gamma\{[\Delta]\}}{\Gamma\{\Delta\}} \quad b^{[]} \frac{\Gamma\{[\Sigma, [\Delta]]\}}{\Gamma\{\Delta, [\Sigma]\}} \\
 4^{[]} \frac{\Gamma\{[\Delta], [\Sigma]\}}{\Gamma\{[[\Delta], \Sigma]\}} \quad 5^{[]} \frac{\Gamma\{\Delta\}\{\emptyset\}}{\Gamma\{\emptyset\}\{\Delta\}} \text{ depth}(\Gamma\{\}\{\emptyset\}) > 0
 \end{array}$$

For $X \subseteq \{d, t, b, 4, 5\}$, we write $X^{[]}$ for the corresponding subset of $\{d^{[]}, t^{[]}, b^{[]}, 4^{[]}, 5^{[]}\}$.

Example. Proof of $\Diamond A \rightarrow \Box \Diamond A$ in $NK \cup \{b^{[]}, 4^{[]}\}$

$$\begin{array}{c}
 \text{init} \frac{}{[[[\bar{p}, p], \Diamond p]]} \\
 \Diamond \frac{}{[[[\bar{p}], \Diamond p]]} \\
 4^{[]} \frac{}{[[[\bar{p}]], \Diamond p]} \\
 b^{[]} \frac{}{[\bar{p}], [\Diamond p]} \\
 \Box \frac{}{\Box \bar{p}, [\Diamond p]} \\
 \Box \frac{}{\Box \bar{p}, \Box \Diamond p} \\
 \vee \frac{}{\Box \bar{p} \vee \Box \Diamond p}
 \end{array}$$

Problem: Rule $d^{[]}$ is not admissible in the presence of cut.

Problem: Rule $d^{[]}$ is not admissible in the presence of cut.

Solution:

- ▶ Show how derivations in $NK \cup \{t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\} \cup \{d^{[]} \} \cup \{\text{cut}\}$ can be transformed into derivations in $NK \cup \{t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\} \cup \{d^{[]} \}$;
- ▶ Show that $d^{[]} is admissible in $NK \cup X^\diamond$.$

Problem: Rule $d^{[]}$ is not admissible in the presence of cut.

Solution:

- ▶ Show how derivations in $NK \cup \{t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\} \cup \{d^{[]} \} \cup \{\text{cut}\}$ can be transformed into derivations in $NK \cup \{t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\} \cup \{d^{[]} \}$;
- ▶ Show that $d^{[]} is admissible in $NK \cup X^\diamond$.$

For $X \subseteq \{d, t, b, 4, 5\}$:

Lemma (Weakening, Contraction). The rules wk and ctr are height- and cut-rank preserving admissible in $NK \cup X^\diamond \cup \{d^{[]} \} \cup \{\text{cut}\}$.

Problem: Rule $d^{[]}$ is not admissible in the presence of cut.

Solution:

- ▶ Show how derivations in $NK \cup \{t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\} \cup \{d^{[]} \} \cup \{\text{cut}\}$ can be transformed into derivations in $NK \cup \{t^\diamond, b^\diamond, 4^\diamond, 5^\diamond\} \cup \{d^{[]} \}$;
- ▶ Show that $d^{[]} is admissible in $NK \cup X^\diamond$.$

For $X \subseteq \{d, t, b, 4, 5\}$:

Lemma (Weakening, Contraction). The rules wk and ctr are height- and cut-rank preserving admissible in $NK \cup X^\diamond \cup \{d^{[]} \} \cup \{\text{cut}\}$.

Lemma (Invertibility). All the rules of $NK \cup X^\diamond \cup \{d^{[]} \} \cup \{\text{cut}\}$ are height- and cut-rank preserving invertible.

Lemma (Admissibility of structural modal rules).

- (i) Let $X \subseteq \{t, b, 4, 5\}$ be 45-closed, and let $\rho \in X$. Then rule $\rho^{[]}$ is cut-rank preserving admissible in $NK \cup X^\diamond \cup \{\text{cut}\}$ and in $NK \cup X^\diamond \cup \{\text{cut}\} \cup \{d^{[]} \}$.
- (ii) Let $X \subseteq \{d, t, b, 4, 5\}$ be 45-closed, and let $d \in X$. Then rule $d^{[]} is admissible in $NK \cup X^\diamond$.$

Proof. Case $b^{[]} is admissible in $NK \cup \{b^\diamond, 4^\diamond, 5^\diamond\} \cup \{\text{cut}\} \cup \{d^{[]} \}$.$

$$\begin{array}{c}
 \frac{\Gamma\{\Sigma, [\Delta]\}}{b^{[]} \frac{\Gamma\{\Delta, [\Sigma]\}}{\Gamma\{\Delta, [\Sigma]\}}} \\
 \\
 \frac{4^\diamond \frac{\Gamma\{[\diamond A, \Sigma, [\diamond A, \Delta]]\}}{b^{[]} \frac{\Gamma\{[\diamond A, \Sigma, [\Delta]]\}}{\Gamma\{\Delta, [\diamond A, \Sigma]\}}} \quad \rightsquigarrow \quad \frac{b^{[]} \frac{[\diamond A, \Sigma, [\diamond A, \Delta]]}{\Gamma\{\Delta, \diamond A, [\diamond A, \Sigma]\}}}{5^\diamond \frac{\Gamma\{\Delta, \diamond A, [\diamond A, \Sigma]\}}{\Gamma\{\Delta, [\diamond A, \Sigma]\}}}
 \end{array}$$

Let $X \subseteq \{t, b, 4, 5\}$, and let Y be a subset of $\{4, 5\} \cap X$. Then:

- ▶ Let \mathcal{D} be a proof in $\text{NK} \cup X^\diamond \cup \{\text{cut}\}$ (or in $\text{NK} \cup X^\diamond \cup \{\text{cut}\} \cup \{\text{ser}^{[1]}\}$) as displayed below, with $cr(\mathcal{D}_1) = cr(\mathcal{D}_2) = p = c(A)$. Then, we can construct the proof \mathcal{D}^* below in the same system, with $cr(\mathcal{D}^*) = p$.

$$\mathcal{D} = \frac{\frac{\mathcal{D}_1}{\Gamma\{A\}} \quad \frac{\mathcal{D}_2}{\Gamma\{\bar{A}\}}}{\text{cut} \quad \Gamma\{\emptyset\}} \rightsquigarrow \frac{\mathcal{D}^*}{\Gamma\{\emptyset\}}$$

- ▶ Let \mathcal{D} be a proof in $\text{NK} \cup X^\diamond \cup \{\text{cut}\}$ (or in $\text{NK} \cup X^\diamond \cup \{\text{cut}\} \cup \{\text{ser}^{[1]}\}$) as displayed below, with $cr(\mathcal{D}_1) = cr(\mathcal{D}_2) = p = c(A)$. Then, we can construct the proof \mathcal{D}^* below in the same system, with $cr(\mathcal{D}^*) = p$.

$$\mathcal{D} = \frac{\frac{\mathcal{D}_1}{\Gamma\{\Box A\}\{\emptyset\}^n} \quad \frac{\mathcal{D}_2}{\Gamma\{\Diamond \bar{A}\}\{\Diamond \bar{A}\}^n}}{Y\text{-cut} \quad \Gamma\{\emptyset\}\{\emptyset\}^n} \rightsquigarrow \frac{\mathcal{D}^*}{\Gamma\{\emptyset\}}$$

Proof: By induction on the sum of heights of \mathcal{D}_1 and \mathcal{D}_2 .

Two cases of the Reduction Lemma

$$\frac{\frac{\frac{\Gamma\{[A], [\Delta]\}}{\Gamma\{\Box A, [\Delta]\}} \quad \frac{\Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, \Delta]\}}{\Gamma\{\Diamond \bar{A}, [\Delta]\}}}{\Gamma\{[\Delta]\}} \quad \rightsquigarrow \quad \frac{\frac{\Gamma\{[A], [\Delta]\}}{\Gamma\{\Box A, [\Delta]\}} \quad \Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, \Delta]\}}{\Gamma\{[\Delta]\}} \quad \text{4-cut}$$

$$\frac{\frac{\frac{\Gamma\{[A], [[\Sigma]]\}}{\Gamma\{\Box A, [[\Sigma]]\}} \quad \frac{\Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, [\bar{A}, \Sigma]]\}}{\Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, [\Sigma]]\}}}{\Gamma\{[[\Sigma]]\}} \quad \rightsquigarrow$$

$$\rightsquigarrow \frac{\frac{\frac{\Gamma\{[A], [[\Sigma]]\}}{\Gamma\{[[A], [\Sigma]]\}} \quad \frac{\frac{\Gamma\{[A], [[\Sigma]]\}}{\Gamma\{\Box A, [[\Sigma]]\}} \quad \frac{\Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, [\bar{A}, \Sigma]]\}}{\Gamma\{\Diamond \bar{A}, [\Diamond \bar{A}, [\Sigma]]\}}}{\Gamma\{[[\bar{A}, \Sigma]]\}} \quad \text{4-cut} \quad \frac{\Gamma\{[[A], [\Sigma]]\}}{\Gamma\{[[A, \Sigma]]\}} \quad \text{4-l} \quad \frac{\Gamma\{[[A, \Sigma]]\}}{\Gamma\{[[\Sigma]]\}} \quad \text{cut}$$

Theorem (Cut-elimination). For $X \subseteq \{d, t, b, 4, 5\}$ 45-closed, if Γ is derivable in $NK \cup X^\diamond \cup \{\text{cut}\}$, then it is derivable in $NK \cup X^\diamond$.

Can we get rid of the 45-closure condition?

Can we get rid of the 45-closure condition?

YES: by adding to NK both the propagation rules X^\diamond and the structural rules $X^[]$. The price to pay is that contraction is no longer admissible.

Theorem. For $X = \{d, t, b, 4, 5\}$, and Γ a set of formulas, it holds that Γ is derivable in $NK_{ctr} \cup X_{ctr}^\diamond \cup X^[]$ iff Γ is X-valid.

Can we get rid of the 45-closure condition?

YES: by adding to NK both the propagation rules X^\diamond and the structural rules $X^[]$. The price to pay is that contraction is no longer admissible.

Theorem. For $X = \{d, t, b, 4, 5\}$, and Γ a set of formulas, it holds that Γ is derivable in $NK_{ctr} \cup X_{ctr}^\diamond \cup X^[]$ iff Γ is X-valid.

Can we get rid of the propagation rules, and use $NK_{ctr} \cup X^[]$?

Can we get rid of the 45-closure condition?

YES: by adding to NK both the propagation rules X^\diamond and the structural rules $X^[]$. The price to pay is that contraction is no longer admissible.

Theorem. For $X = \{d, t, b, 4, 5\}$, and Γ a set of formulas, it holds that Γ is derivable in $NK_{ctr} \cup X_{ctr}^\diamond \cup X^[]$ iff Γ is X-valid.

Can we get rid of the propagation rules, and use $NK_{ctr} \cup X^[]$?

NO, some combinations are incomplete, and one example is given in [Marin & Straßburger, 2014].

Roadmap

Conclusions

Comparison

Within the S5-cube ($X \subseteq \{d, t, b, 4, 5\}$):

	fml. interpr.	invertible rules	analyti- city	termination proof search	counterterm. constr.	modu- larity
$\text{labK} \cup X$	no	yes	yes	yes, for most	yes, easy!	yes
HS5	yes	yes	yes	yes	yes, easy!	no
$\text{NK} \cup X^\diamond$	yes	yes	yes	yes	yes	45-clause

Comparison

Within the S5-cube ($X \subseteq \{d, t, b, 4, 5\}$):

	fml. interpr.	invertible rules	analyti- city	termination proof search	counterterm. constr.	modu- larity
$\text{labK} \cup X$	no	yes	yes	yes, for most	yes, easy!	yes
HS5	yes	yes	yes	yes	yes, easy!	no
$\text{NK} \cup X^\diamond$	yes	yes	yes	yes	yes	45-clause

And beyond the S5-cube?

A few words in conclusion

- ▶ There is no good or bad calculus, rather there are different calculi with different properties. The “right” calculus to consider (if there is any) depends on your aim

A few words in conclusion

- ▶ There is no good or bad calculus, rather there are different calculi with different properties. The “right” calculus to consider (if there is any) depends on your aim
- ▶ Labelled and structured calculi are different but not necessarily opposite or incompatible approaches
 - ▶ In some cases, mutual translations between labelled and structured sequents, labelled and structured derivations
 - ▶ Possibility to combine labels and structure in the same calculus

A few words in conclusion

- ▶ There is no good or bad calculus, rather there are different calculi with different properties. The “right” calculus to consider (if there is any) depends on your aim
- ▶ Labelled and structured calculi are different but not necessarily opposite or incompatible approaches
 - ▶ In some cases, mutual translations between labelled and structured sequents, labelled and structured derivations
 - ▶ Possibility to combine labels and structure in the same calculus
- ▶ We have presented the most standard (and possibly simplest) extensions of the sequent calculus. However, once established that one can extend the language or the structure, there is no limit to imagination: 2-sequents, display calculus, sequents with histories, linear nested sequents, grafted hypersequents, etc.

A few words in conclusion

- ▶ There is no good or bad calculus, rather there are different calculi with different properties. The “right” calculus to consider (if there is any) depends on your aim
- ▶ Labelled and structured calculi are different but not necessarily opposite or incompatible approaches
 - ▶ In some cases, mutual translations between labelled and structured sequents, labelled and structured derivations
 - ▶ Possibility to combine labels and structure in the same calculus
- ▶ We have presented the most standard (and possibly simplest) extensions of the sequent calculus. However, once established that one can extend the language or the structure, there is no limit to imagination: 2-sequents, display calculus, sequents with histories, linear nested sequents, grafted hypersequents, etc.
- ▶ We have presented labelled and structured calculi for the S5 cube of normal modal logics because it is a well-known family of modal logics, and it is the context where this solutions have been initially developed. However, the same or similar solutions have been applied to many other kinds of logics: non-normal modal logics, intuitionistic modal logics, conditional logics, temporal logics, intermediate logics, etc.

Questions?

	fml. interpr.	invertible rules	analyti- city	termination proof search	counterm. constr.	modu- larity
labK \cup X	no	yes	yes	yes, for most	yes, easy!	yes
HS5	yes	yes	yes	yes	yes, easy!	no
NK \cup X $^\diamond$	yes	yes	yes	yes	yes	45-clause

- ▶ Questions, suggestions, discussion etc. are very welcome
“m.girlando at uva dot nl” “tiziano.dalmonte at unibz dot it”
- ▶ Thank you for attending, we hope you enjoyed the course