

PSY9511: Seminar 5

Unsupervised learning

Esten H. Leonardsen

24.10.24



**UNIVERSITETET
I OSLO**

1. Overview of unsupervised learning
2. Clustering
 - K-means
 - Hierarchical
3. Dimensionality reduction
 - Principal component analysis (PCA)
 - Independent component analysis (ICA)

Unsupervised learning




UNIVERSITY
OF OSLO

Supervised learning: Find $\hat{y} = f(X)$

Unsupervised learning: Motivation

Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y




Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y
- Predictive: Predict y given new X .

Unsupervised learning: Motivation


Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y
- Predictive: Predict y given new X .
 - Because the predictions are useful



Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y
- Predictive: Predict y given new X .
 - Because the predictions are useful
 - Because we want to know if it is possible



Unsupervised learning: Motivation

Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y
- Predictive: Predict y given new X .
 - Because the predictions are useful
 - Because we want to know if it is possible

Unsupervised learning: Are there some interesting patterns in X ?



Unsupervised learning: Motivation

Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y
- Predictive: Predict y given new X .
 - Because the predictions are useful
 - Because we want to know if it is possible

Unsupervised learning: Are there some interesting patterns in X ?

- Can we find subgroups or interesting axes of variability?



Unsupervised learning: Motivation

Supervised learning: Find $\hat{y} = f(X)$

- Descriptive: Understand the relationship between X and y
- Predictive: Predict y given new X .
 - Because the predictions are useful
 - Because we want to know if it is possible

Unsupervised learning: Are there some interesting patterns in X ?

- Can we find subgroups or interesting axes of variability?
- Visualization



Clustering



UNIVERSITY
OF OSLO

Are there some (naturally occurring) subgroups in our dataset?

Clustering: Background


Are there some (naturally occurring) subgroups in our dataset?

x_1	x_2
0.20	-0.26
0.15	-0.33
0.03	0.07
-0.07	-0.01
-0.06	0.00
0.28	-0.24
0.21	-0.35
0.20	-0.32
0.30	0.25
0.00	-0.12

Clustering: Background

Are there some (naturally occurring) subgroups in our dataset?


x_1	x_2
0.20	-0.26
0.15	-0.33
0.03	0.07
-0.07	-0.01
-0.06	0.00
0.28	-0.24
0.21	-0.35
0.20	-0.32
0.30	0.25
0.00	-0.12



Clustering: Background

Are there some (naturally occurring) subgroups in our dataset?


x_1	x_2
0.20	-0.26
0.15	-0.33
0.03	0.07
-0.07	-0.01
-0.06	0.00
0.28	-0.24
0.21	-0.35
0.20	-0.32
0.30	0.25
0.00	-0.12



Clustering: Background

Are there some (naturally occurring) subgroups in our dataset?

x_1	x_2
0.20	-0.26
0.15	-0.33
0.03	0.07
-0.07	-0.01
-0.06	0.00
0.28	-0.24
0.21	-0.35
0.20	-0.32
0.30	0.25
0.00	-0.12



K-means clustering

K-means clustering: Find k clusters in the data to minimize the *within-cluster variance*:

$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sum_{j=1}^p (x_{ij} - x_{i'j})^2 \right)$$

K-means clustering


K-means clustering: Find k clusters in the data to minimize the *within-cluster variance*:

$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$



K-means clustering


x_1	x_2
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25



$$\underset{C_1, \dots, C_K}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$


K-means clustering

x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering



$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$


K-means clustering



$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering


x_1	x_2
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25



$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering

x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering

↪


x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25



$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering


x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25



$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering

x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25




$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering

← j →

x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25



$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering

x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25

$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$

K-means clustering

x ₁	x ₂
1	1
1.5	1
1.25	1.25
-1	1
-1.5	1
-1.25	1.25
0.25	0
-0.25	0
0	-0.25

$$\underset{C_1, \dots, C_k}{\text{minimize}} \left(\sum_{k=1}^K \frac{1}{|C_k|} \sum_{i, i' \in C_k} \sqrt{\sum_{j=1}^p (x_{ij} - x_{i'j})^2} \right)$$