

The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders

Esten H. Leonardsen

26.10.23



**UNIVERSITETET
I OSLO**

Overview

1. Background: Defining the scope of the lecture.
2. State-of-the-art: How is neuroimaging beyond T1-weighted MRI currently being used with respect to neuropsychiatric disorders.
3. The future: Challenges and opportunities in using neuroimaging for predicting neuropsychiatric disorders moving forward.



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Background

- The role of **neuroimaging** beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Background

- The role of **neuroimaging** beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3




Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3



Sample from the MNE library




Background


- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3



Sample from the MNE library



Sample from Tremlay et al., 2016

Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic interneurons in the neocortex: from cellular properties to circuits. *Neuron*, 91(2), 260-292




Background


- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3



Sample from the MNE library



Sample from Tremblay et al., 2016



Meta Quest Pro

Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic interneurons in the neocortex: from cellular properties to circuits. *Neuron*, 91(2), 260-292



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3




Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3



Arbabshirani et al., 2017




Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3




Ebrahimighahnaveh, M. A., Luo, S., & Chiong, R. (2020). Deep learning to detect Alzheimer's disease from neuroimaging: A systematic literature review. Computer methods and programs in biomedicine, 187, 105242




Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Bert from FreeSurfer 7.3



Quaak et al., 2020

Quaak, M., van de Mortel, L., Thomas, R. M., & van Wingen, G. (2021). Deep learning applications for the classification of psychiatric disorders using neuroimaging data: systematic review and meta-analysis. *NeuroImage: Clinical*, 30, 102584



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders

Alzheimer's disease and other
causes of dementia

Multiple Sclerosis

Parkinson's Disease



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders

Alzheimer's disease and other
causes of dementia

Multiple Sclerosis

Parkinson's Disease


Bipolar Disorder
Schizophrenia

Depressive disorders




Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders




Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Background

- The role of neuroimaging beyond T1-weighted MRI in the diagnosis and prediction of neuropsychiatric disorders



Vogel & Black (2024)



The current roles of neuroimaging modalities




UNIVERSITETET
I OSLO


Other structural MRI modalities




T1



Other structural MRI modalities




Other structural MRI modalities




Adapted from Shoeibi et al., 2021

Shoeibi, A., Khodatars, M., Jafari, M., Mordian, P., Rezaei, M., Alizadehsani, R., ... & Acharya, U. R. (2021). Applications of deep learning techniques for automated multiple sclerosis detection using magnetic resonance imaging: A review. *Computers in Biology and Medicine*, 136, 104697.


Other structural MRI modalities




Adapted from Shoeibi et al., 2021



Other structural MRI modalities



Adapted from Shoeibi et al., 2021




Other structural MRI modalities



The image displays three axial MRI slices of a human brain side-by-side. The left slice is labeled 'T1' at the top, the middle slice is labeled 'T2' at the top, and the right slice is labeled 'FLAIR' at the top. Each slice shows internal brain structures in grayscale.

Adapted from Shoeibi et al., 2021




De Angelis et al., 2019



De Angelis, F., Brownlee, W. J., Chard, D. T., & Trip, S. A. (2019). New MS diagnostic criteria in practice. *Practical Neurology*, 19(1), 64-67.

Other structural MRI modalities




Adapted from Shoeibi et al., 2021

T2 in PD



Other structural MRI modalities




Adapted from Shoeibi et al., 2021


T2 in AD



Diffusion Tensor Imaging (DTI)



Positron Emission Tomography (PET)



Functional Magnetic Resonance Imaging (fMRI)



Magnetoencephalography (MEG)



Alzheimer's disease

- Pathological changes are apparent in a variety of neuroimaging modalities, to a level which can support individual diagnostics.
-

Method	Sensitivity	Specificity	Accuracy	Pros	Cons	Refs.
MRI	80–95%	55–98%	89–97%	Spatial resolution	Temporal resolution, MR exposure	[36,105–108]
FDG-PET	43–100%	57–100%	50–100%	Clear image	FDG injection	[44,47,57,63,109]
fNIRS	82–94%	72–88%	50–90%	High speed, portability	Spatial resolution	[110,111]
fMRI	84–94%	68–91%	75–93%	Spatial resolution	Not portable/Expose MR	[78,80,112]
EEG	35–88%	82–100%	62–92%	High speed, portability	Spatial resolution	[99,101,103,113,114]



Multiple sclerosis

- Lesions apparent in T2-weighted and FLAIR images are used clinically to support diagnostics.
- DTI reveals group-wise differences also in apparently healthy white matter.
- A stereotypical trajectory that has been observed in patients using fMRI is heightened activation up until a point where it is no longer possible to compensate for structural damage, after which the activation generally resides.




Parkinson's disease


- DTI has been shown useful for classifying patient subtypes and correlates with motor and cognitive function in PD patients.
- fMRI has been shown useful for differentiating patient subtypes, supporting the notion of PD subtypes as network models.
- PET imaging can be used to reveal subtypes



Bipolar disorder



Bipolar disorder



Future opportunities and challenges




UNIVERSITETET
I OSLO

Multimodality and data fusion



The role of prediction



Prediction versus interpretability

