| | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|-------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Total deliveries ^a | 8,877 | 8,915 | 8,760 | 8,485 | 8,682 | 8,209 | 8,335 | 8,289 | 7,958 | 8,083 |
| GH-no sig proteinuria | 6.0% | 6.0% | 4.9% | 5.0% | 5.1% | 5.5% | 5.2% | 4.3% | 5.1% | 5.4% |
| GH-sig proteinuria | 1.9% | 1.5% | 1.9% | 2.3% | 2.0% | 2.1% | 2.2% | 2.1% | 2.5% | 2.5% |
| GH-total | 7.8% | 7.5% | 6.8% | 7.3% | 7.1% | 7.6% | 7.4% | 6.5% | 7.6% | 7.9% |

^a Among women without pre-existing hypertension.

Note: Gestational hypertension is hypertension that is first detected after the 20th week of gestation. Gestational hypertension with significant proteinuria includes those cases denoted as such; severe pre-eclampsia; HELLP syndrome (Hemolysis, Elevated Liver Enzymes, Low Platelets); and eclampsia.