



Surveillance of Congenital Anomalies

NAVIGATING BIRTH DEFECTS IN NOVA SCOTIA A TEN YEAR PERSPECTIVE 2013-2022



2024

Reproductive Care Program of Nova Scotia

Navigating Birth Defects in Nova Scotia

***A Ten Year Perspective
2012-2022***



Contents

Background	v
SCAN-NS Activity and Report Summary	vii
Introduction	ix
1 Methodology	1
1.1 Case Definitions	1
1.2 Case Ascertainment	2
1.3 Quality Control Measures and Data Linkage	3
1.4 Confidentiality and Release of Data	3
2 Patterns of Selected Congenital Anomalies in Nova Scotia	9
2.1 Selected Anomalies	9
2.1.1 Selected Anomaly Definitions	9
2.1.1.1 Neural tube defects	9
2.1.1.2 Selected central nervous system defects	9
2.1.1.3 Selected sense organ defects	10
2.1.1.4 Selected congenital heart defects	11
2.1.1.5 Oro-facial clefts	12
2.1.1.6 Selected gastrointestinal defects	12
2.1.1.7 Selected genital anomalies	13
2.1.1.8 Selected urinary tract defects	13
2.1.1.9 Hip dysplasia (Q65)	14
2.1.1.10 Limb deficiency defects (Q71-Q73)	14
2.1.1.11 Selected abdominal wall defects	14
2.1.1.12 Selected chromosomal defects	15

3	Anomalies Overview	17
3.1	Neural tube defects	17
3.2	Selected central nervous system defects	20
3.3	Selected sense organ defects	21
3.4	Selected congenital heart defects	22
	Appendix	23
3.5	Appendix 1 - Linear trend analysis and p-values, 2013-2022 . .	23
3.6	Appendix 2 - Total birth, number of cases, anomalies and anomalies per case, 2013-2022	26
3.7	Appendix 3 - Prevalence rate per 10,000 total births, aggregated prevalence rate, 2013-2022	26

Background

Gaps exist in the nation's capacity to conduct surveillance for congenital anomalies. In 2002, the Canadian Congenital Anomalies Surveillance Network (CCASN) was established

To support the development and maintenance of high quality population-based surveillance systems of congenital anomalies that will provide information to improve the health of Canadian children and their families.¹

The CCASN produced a series of documents regarding the establishment of provincial/territorial surveillance systems for congenital anomalies², coding of congenital anomalies³, and delineating data elements required for minimal and enhanced surveillance for congenital anomalies⁴.

In Nova Scotia, data regarding congenital anomalies diagnosed at birth or during the birth admission have been collected since 1987 in the Nova Scotia Atlee Perinatal Database (NSAPD) managed by the Reproductive Care Program of Nova Scotia (RCP). The NSAPD collects data about all live born infants and stillbirths for all fetuses delivered ≥ 20 weeks gestation, or ≥ 500 grams, or live born and all co-multiples of the aforementioned; the NSAPD also captures information about the mothers of the infants/fetuses mentioned above. The NSAPD does not capture data for congenital anomaly cases identified during admissions that occur after 28 days of age. The NSAPD also does not capture data for pregnancies ending in termination.

Since 1992, data regarding congenital anomalies diagnosed antenatally, or postnatally up to the time of discharge from the IWK Health Centre, or on the basis of an autopsy have been captured through the Fetal Anomaly Database (FADB) housed at the Division of Maternal and Fetal Medicine, IWK Health Centre. The IWK is the only obstetrics/paediatric tertiary care centre in the

¹Public Health Agency of Canada. What is Canadian Congenital Anomalies Surveillance Network (CCASN)

²Canadian Congenital Anomalies Surveillance Network. Recommendations concerning the establishment of congenital anomalies surveillance in Canadian jurisdictions.

³Canadian Congenital Anomalies Surveillance Network. Coding of fetal anomalies.

⁴Canadian Congenital Anomalies Surveillance Network. Recommended data variables for congenital anomalies surveillance in Canadian provinces & territories.

Maritimes; as such, the FADB captures most congenital anomaly cases in Nova Scotia resulting in live births, still births, and terminations. The FADB does not capture congenital anomaly cases diagnosed after the birth admission, many of which are present in other data sources within the province. The FADB also uses an in-house coding system to classify anomalies, a practice that predates the CCASN recommendation to use the Royal College of Paediatrics and Child Health codes (adapted from the [International Statistical Classification of Diseases and Related Health Problems, 10th Revision – ICD-10](#)).

SCAN-NS Activity and Report Summary

1. This is the first provincial report on the prevalence of congenital anomalies in Nova Scotia generated by the Surveillance of Congenital Anomalies in Nova Scotia project (SCA-NS).
 2. The International Classification of Diseases - 10th Edition ([ICD-10-CA](#)) has been adopted by the IWK Health Centre as the reporting classification system and SCA-NS uses the Royal College of Paediatrics and Child Health adaptation of ICD-10 as well. The anomalies outlined in this report reflect the “sentinel anomalies list” adopted by the Nova Scotia Congenital Anomaly Advisory Group. Other anomaly data is collected and available upon request.
 3. Nova Scotia continues to participate in the Canadian Congenital Anomalies Surveillance System (CCASS), administered by the Maternal and Infant Health Section of PHAC.
 4. The numerator data include live births, stillbirths, and also fetal losses < 20 weeks gestation with congenital anomalies. Denominator data include live births and stillbirths only. By including fetal losses in the numerator, the reported rates should be more representative of true congenital anomaly rates. Fetal losses have been ascertained since 1992.
 5. Data provided in this report is a compilation of several data Nova Scotia databases.
- Nova Scotia Atlee Perinatal Database (NSAPD).
 - Fetal Anomaly Database (FADB).
 - Canadian Institute for Health Information Discharge Abstract Database.
 - Vital Statistics Database; Service Nova Scotia.
 - Medical Services Insurance (MSI) Claims.



Introduction

The purpose of SCA-NS is to improve Congenital Anomaly case ascertainment, cluster detection, local and national data comparison and reporting capability. Program objectives - Provide baseline data on occurrence. - Identify populations at increased risk. - Monitor changes in occurrence. - Investigate clusters. - Refer affected children to services. - Evaluate prevention programs. - Create research opportunities.

SCA-NS includes all Nova Scotia residents, with ascertainment from other Maritime Provinces for those cases where the mother delivered elsewhere. Information on cases includes the prenatal period up to one year of age including live births, stillbirths and terminations of pregnancies in which a fetal anomaly has been detected.



1

Methodology

1.1 Case Definitions

A congenital anomaly is an abnormality that is present at birth, even if not diagnosed until months or years later. Most congenital anomalies are present long before the time of birth, some in the embryonic period (up to the end of the seventh week of gestation) and others in the fetal period (eighth week to term). The term “anomaly” covers all the major classes of abnormalities of development, of which there are four major categories as follows:

- **Malformation** - a morphologic defect of an organ, part of an organ or a larger region of the body resulting from an intrinsically abnormal developmental process (e.g., spina bifida, cleft lip and palate).
- **Deformation** - an abnormal form, shape or position of a part of the body caused by mechanical forces (e.g., extrinsic force such as intrauterine constraint causing some forms of clubfoot).
- **Disruption** - a morphologic defect of an organ, part of an organ or a larger region of the body resulting from the extrinsic breakdown of, or an interference with, an originally normal developmental process (e.g., an infection such as rubella or a teratogen such as thalidomide).
- **Dysplasia** - the abnormal organization of cells into tissues and its morphologic result (e.g., Marfan Syndrome, osteogenesis imperfecta).

Other definitions related to pregnancy outcomes for the purposes of this report are as follows:

- **Live birth** - a complete expulsion or extraction from the mother, irrespective of the duration of the pregnancy, of a fetus in which, after expulsion or extraction, there is breathing, beating of the heart, pulsation of the umbilical cord or definite movement of voluntary muscle (Alberta Vital Statistics Annual review, 2000).

- **Stillbirth** - a complete expulsion or extraction from the mother, after at 20 weeks of pregnancy or more or after attaining a weight of 500 grams or more, of a fetus in which, after the expulsion or extraction, there is no breathing, beating of the heart, pulsation of the umbilical cord or unmistakable movement of voluntary muscle (Alberta Vital Statistics Annual review¹).
- **Gestation** - completed weeks of pregnancy at delivery.
- **Preterm birth (a.k.a premature)** - a birth before 37 weeks of gestation (< 37 weeks).
- **Termination of Pregnancy (ToP)** - for our purposes, includes any pregnancy loss before 20 weeks gestation (< 20 weeks). Most cases are therapeutic terminations for congenital anomalies but spontaneous abortions or intrauterine fetal deaths with fetal anomalies could also be included.

Anomaly definitions are based, for the most part, on those provided by the International Clearinghouse for Birth Defects Surveillance and Research (ICBDSR) and National Birth Defects Prevention Network (NBDPN).

1.2 Case Ascertainment

An infant can be ascertained at any time up to the first birthday. Multiple ascertainment of the same infant can occur and is encouraged, as this frequently improves the quality and reliability of the data.

As several malformations may occur in the same infant, it is advantageous to allow each to be reported so that groups of associated malformations may be studied. This, however, leads to difficulties since the final tabulations may be reported as total malformations (anomaly rates) or as the total number of malformed infants (case rates). The tables in [Appendix A3] report anomaly rates, which in most cases are similar to case rates (e.g. cleft palate, hypospadias, microcephaly). Whereas with limb anomalies, there can be multiple different limb anomalies in the same infant.

SCA-NS obtains information about infants with congenital anomalies from a variety of independent sources. Acquisition of additional reporting agencies is always a priority since the use of multiple sources of information improves not only the ease but also completeness of ascertainment as well as for verification of the diagnostic data.

¹Alberta Vital Statistics Annual review

1.3 Quality Control Measures and Data Linkage

Data quality is ensured through both a coded linking process and by manual chart audits of unmatched anomaly records. The first step in data quality control involves the linking of disparate fetal/infant records between databases. Linking is based on the matching of several unique variables in each record.

- If two records “pass”: this variable matching process, the records are considered link and therefore report anomalies on the same fetus/infant. This allows SCA-NS to create a record that reports anomalies from several databases for the same fetus/infant.
- If there are records that “fail” the automated matching process they are reviewed manually to determine if that information was only found in one data source or if information was entered incorrectly.

1.4 Confidentiality and Release of Data

Authority for the Collection, Use, and Disclosure of Personal Information

In Nova Scotia, personal health information is protected by legislation, policies, and professional codes of ethics of healthcare providers. SCA-NS will follow the provisions outlined in the IWK’s Privacy of Personal Health Information policy and the RCP Privacy Policy² and the Data Access for Research and Planning Policy and Procedures³

Data Requests

Data requests will be reviewed by a sub-committee of the Joint Perinatal Epidemiology Research Unit-Population Health Research Unit-RCP Data Access Committee (JDAC). Clinical and technical staff will participate, as will representatives from all stakeholders currently involved in providing data, i.e. Maternal-Fetal Medicine, BIAP, both with a privacy and a technical representative, and other groups such as Cardiology, as the database develops.

²[RCP Privacy Policy](#)

³[Data Access for Research and Planning Policy and Procedures](#)

Disclosure for research purposes also requires approval by a recognized Research Ethics Board and, if approved, a Data Use Agreement. SCA-NS will only consider disclosing individual level data (with or without personal identifiers) following a technical review and approval from a specially-constituted committee.

Limitations of Data and Analysis

One of the major limitations of any surveillance system is that on its own, the information provided does not allow etiology to be determined. If trends indicate a potential problem, then separate investigative studies need to be done. However, with appropriate approvals in place, it would be possible to conduct linkage studies with other data sources to explore potential causes of specific birth defects. SCA-NS data are collected passively from a variety of sources and the completeness and accuracy of data are largely dependent on reporting and coding of several different organizations. SCA-NS reviews anomalies that have been entered into the database on a regular basis. Detailed studies of some individual anomalies or anomaly groups aid in the assessment and maintenance of the data quality. With intensive review, some cases might be reassigned, re-coded or discarded altogether from the database. This continuing review might explain some discrepancies in the data from earlier reports.

Epidemiological and Statistical Measures

- Unless otherwise stated, all the formulas used to compute prevalence and confidence intervals are calculated following the [National Birth Defects Prevention Network \(NBDPN\) guidelines, chapter 8](#):

$$\text{Prevalence at birth} = 1,000 \times \frac{\text{Number of cases with birth defect A in an area and time period}}{\text{Number of live births and stillbirths in that area and time period}}$$

- **Pregnancy outcomes included** - the cases in the **numerator** are derived from all pregnancy outcomes collected by the program and may include:
 - Live births.
 - Stillbirths.
 - Termination of pregnancy (ToP): all fetal losses < 20 weeks gestation with congenital anomalies.

For the **denominator**, we use the total number of live births and stillbirths in the same area and time period from which the cases were ascertained.

TABLE 1.1

Number of Cases	Lower Bound	Upper Bound
0	0.000	3.689
1	0.025	5.572
2	0.242	7.225
3	0.619	8.767
4	1.090	10.242
5	1.623	11.668

Confidence intervals

(95%) confidence intervals are also included because the rate obtained is actually only a point estimate of the unknown, true population rate. The confidence interval provides information about the precision of the estimate. Thus, the confidence intervals are an estimated range of values within which there is a 95% probability that the true population rate will fall.

- ***For a prevalence based on a small number of cases*** - For small numbers of cases (arbitrarily defined here as < 30 cases), we use the Poisson distribution since birth defects are considered to be rare events. The easiest way to use the Poisson distribution is to refer to a table that provides the upper and lower 95% confidence limits for an observed number of cases. For that, we make use of the relationship between the Poisson and chi-square distribution functions. If X is a single observation from a Poisson distribution with mean μ . Then the “exact” 95% confidence limits for μ are given by

$$\left[\frac{qchisq(0.025, 2X)}{2}; \frac{qchisq(0.975, 2(X+1))}{2} \right],$$

where $qchisq()$ is the quantile function for the chi-squared (χ^2) distribution with $2X$, and $2(X+1)$ degrees of freedom. These limits can be computed using any statistical software or taken from a chi-square table (reproduced below for up to 5 cases).

1. Calculate prevalence

$$1,000 \times \frac{\text{Number of cases with birth defect A in an area and time period}}{\text{Number of live births and stillbirths in that area and time period}}$$

2. Look up the lower and upper 95% confidence limit for the number of cases with birth defect A. Using this new number in the numerator calculate the lower and upper 95% confidence limit for prevalence.

$$\begin{aligned} \text{Lower 95\% CL for prevalence} &= 1,000 \times \frac{\text{Lower 95\% CL for cases with birth defect A in an area and time period}}{\text{Number of live births and stillbirths in that area and time period}} \\ \text{Upper 95\% CL for prevalence} &= 1,000 \times \frac{\text{Upper 95\% CL for cases with birth defect A in an area and time period}}{\text{Number of live births and stillbirths in that area and time period}} \end{aligned}$$

Example - Considering that for a birth defect A in an area and time period we had 5 cases with a number of total births of 12,500 we have

$$\begin{aligned} \text{Prevalence} &= 1,000 \times \frac{5}{12,500} = 0.4 \\ \text{Lower 95\% CL for prevalence} &= 1,000 \times \frac{1.623}{12,500} = 0.13 \\ \text{Upper 95\% CL for prevalence} &= 1,000 \times \frac{11.668}{12,500} = 0.93 \end{aligned}$$

- **For a prevalence based on a large number of cases -** For a large number of cases (arbitrarily defined here as ≥ 30 cases), use the Normal distribution because as the number of cases grows larger, the Poisson distribution approximates the Normal distribution. Let c = number of cases in an area and time period and b = number of live births and stillbirths in an area and time period

1. Calculate the lower and upper confidence limit using the following:

$$\begin{aligned} \text{Lower 95\% CL for prevalence} &= 1,000 \times \frac{\left(1 - \frac{1}{9c} - \frac{1.96}{3} \sqrt{\frac{1}{c}}\right)^3}{b} \times c \\ \text{Upper 95\% CL for prevalence} &= 1,000 \times \frac{\left(1 - \frac{1}{9(c+1)} + \frac{1.96}{3} \sqrt{\frac{1}{(c+1)}}\right)^3}{b} \times (c+1) \end{aligned}$$

2. To determine different $(1 - \alpha)$ % confidence limits, replace 1.96 with the corresponding value from the Normal distribution.
3. To obtain confidence limits for the number of cases instead of the prevalence, apply the formulae but do not divide by b = number of live births and stillbirths in an area and time period or multiply by 1,000.

Trend analysis

When appropriate, **trend analysis** are performed. Since our response variable of interest is the number of cases, there is no fixed upper bound n for it. Since number of cases must take a non-negative integer value, its distribution should place its mass on that range.

The simplest such distribution is the **Poisson**. Its probabilities depend on a single parameter, the mean μ . The Poisson distribution is used for counts of events that occur randomly over time or space, when outcomes in disjoint periods or regions are independent. It satisfies $E(X) = Var(X) = \mu$. It approaches normality as μ increases.

In the context of prevalence, the response count X_i has an index t_i such that its expected value is proportional to t_i . Our approach here, is to consider the total births as the index, since this value varies over time. Then, our sample rate is x_i/t_i , with expected value μ_i/t_i . A log-linear model for the expected rate has form

$$\begin{aligned} \log(\mu_i/t_i) &= \beta_0 + \beta_1(Year) \\ \log(\mu_i) - \log(t_i) &= \beta_0 + \beta_1(Year) \\ \log(\mu_i) &= \beta_0 + \beta_1(Year) + \log(t_i) \end{aligned}$$

The adjustment term $-\log(t_i)$, to the log link of the mean is called an *offset*. Another option is to run this model using the identity link. The model is then

$$\begin{aligned} \mu_i/t_i &= \beta_0 + \beta_1(Year) \\ \mu_i &= \beta_0 t_i + \beta_1(Year)t_i \end{aligned}$$

which does not require an offset. It corresponds to an ordinary Poisson GLM using the identity link with no intercept and explanatory variables t_i , $(Year)t_i$ and its estimates are equal to an OLS model without an intercept.

Since we are working with the observed rate per 1,000 total births, adjusting the yearly count by total births is equivalent to have $t_i = \text{total births}/1,000$.

Sometimes, one can note that the count data could show great variability. Under a Poisson model, we would expect the means and variances of the response to be about the same in various groups. A severe limitation of Poisson models is that $E(X) = Var(X) = \mu$. Hence, at a fixed mean the variance cannot decrease as additional predictors enter the model. The greater variability than predicted by the GLM random component reflects **overdispersion**. A common cause of overdispersion is subject heterogeneity. Without adjusting for overdispersion, we use incorrect, artificially small standard errors leading

to artificially small p-values for model coefficients which may lead to wrong inferences and conclusions.

We can take overdispersion into account in several different ways. The simplest is to use an estimated dispersion factor to inflate standard errors. Another way is to use a **negative-binomial** regression model. In this case, $E(X) = \mu$, and $Var(X) = \mu + \gamma \cdot \mu^2$. The index $\gamma > 0$ is a type of dispersion parameter. As $\gamma \rightarrow 0$, $Var(X) \rightarrow \mu$ and the negative binomial distribution converges to the Poisson.

For the aforementioned model,

$$\mu_i = \beta_0 t_i + \beta_1 (Year) t_i$$

significance tests focus on $H_0 : \beta_1 = 0$. The Wald test uses the log-likelihood at $\hat{\beta}_1$, with test statistic $z = \hat{\beta}_1 / SE(\hat{\beta}_1)$ or its square. Under H_0 , z^2 is asymptotically χ_1^2 . At a 5% level, any p-value lower than 5% would provide evidences against H_0 , meaning that at a 5% level they would be considered statistically significant.

2

Patterns of Selected Congenital Anomalies in Nova Scotia

2.1 Selected Anomalies

2.1.1 Selected Anomaly Definitions

All the birth defects descriptions and respective ICD10 codes were adapted from the [National Birth Defects Prevention Network \(NBDPN\) guidelines, chapter 3](#)

2.1.1.1 Neural tube defects

- **Anencephaly and similar malformations (Q00)**

Partial or complete absence of the brain and skull.

- **Encephalocele (Q01)**

Herniation of brain tissue and/or meninges through a defect in the skull. The hernia sac is usually covered by skin.

- **Spina bifida (Q05)**

Incomplete closure of the vertebral spine (usually posteriorly) through which spinal cord tissue and/or the membranes covering the spine (meninges) herniate.

2.1.1.2 Selected central nervous system defects

- **Microcephaly (Q02)**

Microcephaly, or microcephalus, is the clinical finding of a small head when compared with infants of the same sex and age. The head circumference (HC), also known as the occipitofrontal circumference (OFC), is considered a reliable assessment of the volume of the underlying brain. Microcephaly itself is not a malformation but a sign that the brain is abnormally small.

- **Congenital hydrocephalus (Q03)**

An increase in the amount of cerebrospinal fluid within the brain resulting in enlargement of the cerebral ventricles and increased intracranial pressure.

- **Arhinencephaly/Holoprosencephaly (Q04.1, Q04.2)**

Arhinencephaly is an older term for holoprosencephaly which refers more specifically to structural defects of the olfactory system or nose. Holoprosencephaly results from variable degrees of incomplete division of the brain into right and left cerebral hemispheres. There are four types which vary in severity: alobar, semi-lobar, lobar, and middle interhemispheric (MIHV). The condition can also affect development of the face and eyes. The most severely affected have one central eye (cyclopia) and a single tubular-shaped nose located above the eye (proboscis).

2.1.1.3 Selected sense organ defects

- **Anophthalmos/Micropthalmos (Q11.0, Q11.1, Q11.2)**

Anophthalmia is the total absence of eye tissue or apparent absence of the globe of the eye in an otherwise normal orbit. Micropthalmia is the reduced volume of the eye. The corneal diameter is usually less than 10 millimeters, or the anteroposterior globe diameter is less than 20 millimeters. Anophthalmia or micropthalmia may affect one or both eyes, or there may be anophthalmia of one eye and micropthalmia of the other.

- **Anotia/Microtia (Q16.0, Q17.2)**

Anotia is the total absence of the external ear and canal. Microtia is the malformation or hypoplasia of the external ear (auricle, pinna).

- **Choanal atresia (Q30.0)**

Congenital obstruction of the opening of the nasal cavity into the nasopharynx on either side. This prevents communication of the nasal cavity with the pharynx.

2.1.1.4 Selected congenital heart defects

- **Common truncus (Q20.0)**

Failure of separation of the aorta and the pulmonary artery during development, resulting in a single common arterial trunk carrying blood from the heart to both the body and lungs.

- **Transposition of great vessels (Q20.1, Q20.3, Q20.5)**

Transposition of the aorta and the pulmonary artery such that the aorta arises from the right ventricle (instead of the left) and the pulmonary artery arises from the left ventricle (instead of the right).

- **Atrioventricular septal defect (Q21.2)**

A defect in both the lower portion of the atrial septum and the upper portion of the ventricular septum. In extreme cases, virtually the entire atrial and ventricular septae may be missing. The valves controlling blood flow from the atria to the ventricles, the tricuspid and mitral valves may also be abnormal. They may not form from the endocardial cushions during cardiac development into two separate valves, and thus be a single common atrioventricular valve. Together, these defects producing a large opening (canal) in the central part of the heart.

- **Tetralogy of Fallot (Q21.3)**

Congenital obstruction of the opening of the nasal cavity into the nasopharynx on either side. This prevents communication of the nasal cavity with the pharynx.

- **Hypoplastic left heart syndrome (Q23.4)**

A condition in which the structures on the left side of the heart and the aorta are extremely small, insufficient to support systemic circulation and with normally related great arteries. Classically, this condition includes hypoplasia of the left ventricle, atresia or severe hypoplasia of both the mitral and aortic valves, hypoplasia of the aortic arch, and coarctation of the aorta.

- **Coarctation of aorta (Q25.1)**

Narrowing of the descending aorta, which may obstruct blood flow from the heart to the rest of the body. The most common site of coarctation occurs distal to the origin of the left subclavian artery in the region of the ductus arteriosus. If there is complete loss of communication in this location, it is a form of interruption of the aorta (Type A).

2.1.1.5 Oro-facial clefts

- **Cleft palate (Q35, excluding Q35.7)**

An opening in the roof of the mouth resulting from incomplete fusion of the shelves of the palate. The opening may involve the hard palate only, the soft palate only, or both.

- **Cleft lip (Q36)**

A defect in the upper lip resulting from incomplete fusion of the parts of the lip.

- **Cleft palate with cleft lip (Q37)**

A defect in the upper lip resulting from incomplete fusion of the parts of the lip, with an opening in the roof of the mouth.

2.1.1.6 Selected gastrointestinal defects

- **Oesophageal atresia/stenosis, tracheoesophageal fistula (Q39.0-Q39.4)**

Esophageal atresia is condition in which the esophagus ends in a blind pouch and fails to connect with the stomach. Tracheoesophageal fistula is an abnormal communication between the esophagus and the trachea. This is almost always associated with some form of esophageal atresia.

- **Small intestine absence/atresia/stenosis (Q41)**

Complete or partial occlusion of the lumen of one or more segments of the small intestine. Small intestinal atresias are often assigned a type descriptor in the surgical or autopsy report, depending upon the severity of the atresia (types include I, II, IIIA, IIIB, and VI).

- **Ano-rectal absence/atresia/stenosis (Q42.0-Q42.3)**

Complete or partial occlusion of the lumen of one or more segments of the large intestine and/or rectum.

- **Hirschsprung disease (Q43.1)**

Also called congenital aganglionic megacolon is a condition that affects the large intestine (colon) and causes problems with passing stool.

- **Atresia of bile ducts (Q44.2)**

Congenital absence of the lumen of the extrahepatic bile ducts.

2.1.1.7 Selected genital anomalies

- **Cryptorchidism/undescended testicles (Q53.1, Q53.2, Q53.9)**

A testicle that hasn't moved into its proper position in the bag of skin hanging below the penis (scrotum) before birth.

- **Hypospadias (Q54, excluding Q54.4)**

Displacement of the opening of the urethra (urethral meatus) ventrally and proximally (underneath and closer to the body) in relation to the tip of the glans of the penis.

- **Indeterminate sex (Q56)**

In a baby with ambiguous genitalia, the genitals may be incompletely developed or the baby may have characteristics of both sexes. The external sex organs may not match the internal sex organs or genetic sex. Pseudohermaphroditism - is a condition in which the individual has a single chromosomal and gonadal sex but combines features of both sexes in the external genitalia, causing doubt as to the true sex

- **Epispadias (Q64.0)**

Displacement of the opening of the urethra dorsally and proximally (on the top and closer to the body) in relation to the tip of the glans of the penis.

2.1.1.8 Selected urinary tract defects

- **Renal agenesis (Q60.0-Q60.2)**

Complete absence of the kidney.

- **Cystic kidney (Q61.1-Q61.5, Q61.8, Q61.9)**

Genetic disorder characterized by the formation of fluid-filled sacs (cysts) in the kidneys

- **Bladder and cloacal exstrophy (Q64.1)**

Bladder exstrophy is a defect in the lower abdominal wall and anterior wall of the bladder through which the lining of the bladder is exposed to the outside. Cloacal exstrophy is a congenital persistence of a common cloacal cavity into which gut, urethra, and reproductive tracts open with exstrophy of the cavity: usually accompanied by a low omphalocele, imperforate anus, and a (closed) neural tube defect.

- **Lower urinary tract obstruction (Q64.2, Q64.3)**

Posterior urethral valves (PUV) are tissue folds of the posterior urethra and function as valves obstructing urine outflow. Congenital PUV is an abnormal congenital obstructing membrane that is located within the posterior male urethra; this valve is the most common cause of bladder outlet obstruction in male children. Congenital PUV can also be found in virilized females and rarely in normal females. Obstruction could vary from mild to severe.

2.1.1.9 Hip dysplasia (Q65)

Hip dysplasia occurs when a baby's hip socket (acetabulum) is too shallow to cover the head of the thighbone (femoral head) to fit properly.

2.1.1.10 Limb deficiency defects (Q71-Q73)

Complete or partial absence of the upper arm (humerus), lower arm (radius and/or ulna), wrist (carpals), hand (metacarpals), fingers (phalanges), thigh (femur), lower leg (tibia and/or fibula), ankle (tarsals), foot (metatarsals), or toes (phalanges).

2.1.1.11 Selected abdominal wall defects

- **Diaphragmatic hernia (Q79.0)**

Incomplete formation of the diaphragm through which a portion of the abdominal contents herniate into the thoracic cavity.

- **Omphalocele/Exomphalos (Q79.2)**

A defect in the anterior abdominal wall in which the umbilical ring is widened, allowing herniation of abdominal organs, including the small intestine, part of the large intestine, and occasionally the liver and spleen, into the umbilical cord. The herniating organs are covered by a nearly transparent membranous sac.

- **Gastroschisis (Q79.3)**

A congenital opening or fissure in the anterior abdominal wall lateral to the umbilicus through which the small intestine, part of the large intestine, and occasionally the liver and spleen, may herniate. The opening is separated from the umbilicus by a small bridge of skin, and the herniating organs are not covered by a protective membrane. Gastroschisis usually occurs on the right side of the umbilicus, although it may occur on the left.

2.1.1.12 Selected chromosomal defects

- **Down Syndrome (Q90)**

The presence of three copies of all or a large part of chromosome 21.

- **Trisomy 13 - Patau - (Q91.4-Q91.7)**

The presence of three copies of all or a large part of chromosome 13.

- **Trisomy 18 - Edwards - (Q91.0-Q91.3)**

The presence of three copies of all or a large part of chromosome 18.

- **Turner syndrome (Q96)**

Presence of an absent or structurally abnormal second X chromosome in a phenotypic female.



3

Anomalies Overview

3.1 Neural tube defects

At a 5% level, there is no significant trend for neural tube defects ($\hat{\beta}_1 = 0.040$, p-value = 0.363) for the period 2013 - 2022.

Over the past 11 years (2013-2023), there is no evidence showing any trend in the prevalence rate of neural tube defects. On average, the prevalence rate for each subsequent year is 1.040 times of the previous year's rate: (prevalence rate = $\exp(0.040) = 1.040$ per 10,000 total births; $\hat{\beta}_1 = 0.040$, p-value = 0.363; 95% CI for $\hat{\beta}_1$ and for the prevalence rate are given by $CI(95\%)_{\hat{\beta}_1}:[-0.046; 0.125]$, and $CI(95\%)_{rate}:[0.955; 1.133]$, respectively).




FIGURE 3.1 Neural tube defects - Prevalence - Nova Scotia - 2013-2022

Based on the statistical analysis, at a 5% level, it is possible to observe:

- (Q00) - Anencephaly and similar malformations: there is no significant trend ($\hat{\beta}_1 = -0.056$, p-value = 0.394).
- (Q01) - Encephalocele: there is no significant trend ($\hat{\beta}_1 = -0.196$, p-value = 0.144).
- (Q05) - Spina bifida: there is evidence of an increasing trend ($\hat{\beta}_1 = 0.175$, p-value = 0.021).




FIGURE 3.2 (Q00) - Anencephaly and similar malformations - Prevalence.




FIGURE 3.3 (Q01) - Encephalocele - Prevalence.




FIGURE 3.4 (Q05) - Spina bifida - Prevalence.

3.2 Selected central nervous system defects

3.3 Selected sense organ defects

3.4 Selected congenital heart defects

Appendix

The following table of selected sentinel anomalies indicate the trends in congenital anomaly prevalence in Nova Scotia from 1987 through 2023. Sentinel anomalies are those which the International Clearinghouse of Birth Defects Surveillance and Research (ICBDSR), of which we are a member, watches worldwide with the rationale that they are quite easily identified hence more accurately reported. *See Section 2.1 for all the anomalies listed in the report.*

3.5 Appendix 1 - Linear trend analysis and p-values, 2013-2022

Selected anomaly	Number of cases	Trend direction	Statistic (χ^2_1)	P-value
Neural tube defects	64	No significant change (\rightarrow)	0.83	0.363
(Q00) - Anencephaly and similar malformations	29	No significant change (\rightarrow)	0.73	0.394
(Q01) - Encephalocele	6	No significant change (\rightarrow)	2.14	0.144
(Q05) - Spina bifida	29	Increasing (\uparrow)	5.33	0.021
Selected central nervous system defects	108	No significant change (\rightarrow)	0.15	0.697
(Q02) - Microcephaly	53	No significant change (\rightarrow)	0.02	0.901
(Q03) - Congenital hydrocephalus	50	No significant change (\rightarrow)	1.28	0.257
(Q04.1, Q04.2) - Arhinencephaly / Holoprosencephaly	9	No significant change (\rightarrow)	0.06	0.809
Selected sense organ defects	28	No significant change (\rightarrow)	1.31	0.252
(Q11.0-Q11.2) - Anophtalmos / Microphtalmos	7	No significant change (\rightarrow)	0.60	0.437

(continued)

Selected anomaly	Number of cases	Trend direction	Statistic (χ^2_1)	P-value
(Q16.0, Q17.2) - Anotia / Microtia	14	No significant change (\rightarrow)	0.05	0.819
(Q30.0) - Choanal atresia	9	No significant change (\rightarrow)	1.92	0.166
Selected congenital heart defects	211	No significant change (\rightarrow)	0.33	0.564
(Q20.0) - Common truncus	11	No significant change (\rightarrow)	0.12	0.724
(Q20.1, Q20.3, Q20.5) - Transposition of great vessels	59	No significant change (\rightarrow)	1.60	0.206
(Q21.2) - Atrioventricular septal defect	71	No significant change (\rightarrow)	0.61	0.434
(Q21.3) - Tetralogy of Fallot	33	No significant change (\rightarrow)	0.02	0.901
(Q23.4) - Hypoplastic left heart syndrome	28	No significant change (\rightarrow)	0.71	0.401
(Q25.1) - Coarctation of aorta	54	No significant change (\rightarrow)	0.05	0.820
Oro-facial clefts	102	Increasing (\uparrow)	5.39	0.020
(Q35) - Cleft palate only	67	No significant change (\rightarrow)	1.69	0.194
(Q36) - Cleft lip only	18	No significant change (\rightarrow)	1.41	0.236
(Q37) - Cleft palate with cleft lip	39	No significant change (\rightarrow)	2.84	0.092
Selected gastrointestinal defects	112	No significant change (\rightarrow)	1.83	0.176
(Q39.0-Q39.4) - Oesophageal atresia / stenosis, tracheoesophageal fistula	29	No significant change (\rightarrow)	0.33	0.566
(Q41) - Small intestine absence / atresia / stenosis	32	No significant change (\rightarrow)	1.28	0.257
(Q42.0-Q42.3) - Ano-rectal absence / atresia / stenosis	31	No significant change (\rightarrow)	1.38	0.240
(Q43.1) - Hirschsprung disease	17	No significant change (\rightarrow)	0.41	0.521
(Q44.2) - Atresia of bile ducts	9	No significant change (\rightarrow)	0.16	0.694
Selected genital anomalies	561	No significant change (\rightarrow)	0.03	0.871










(continued)

Selected anomaly	Number of cases	Trend direction	Statistic (χ^2_1)	P-value
(Q53.1, Q53.2, Q53.9) - Cryptorchidism / undescended testicles	262	No significant change (\rightarrow)	0.19	0.663
(Q54, excluding Q54.4) - Hypospadias	276	No significant change (\rightarrow)	2.77	0.096
(Q56) - Indeterminate sex	24	Increasing (\uparrow)	5.01	0.025
(Q64.0) - Epispadias	8	No significant change (\rightarrow)	0.88	0.349
Selected urinary tract defects	163	No significant change (\rightarrow)	1.78	0.182
(Q60.0-Q60.2) - Renal agenesis	74	No significant change (\rightarrow)	3.23	0.072
(Q61.1-Q61.5, Q61.8, Q61.9) - Cystic kidney	79	No significant change (\rightarrow)	0.33	0.568
(Q64.1) - Bladder and cloacal exstrophy	4	No significant change (\rightarrow)	0.00	0.991
(Q64.2, Q64.3) - Lower urinary tract obstruction	20	No significant change (\rightarrow)	0.57	0.451
Hip dysplasia	214	Decreasing (\downarrow)	7.85	0.005
Limb deficiency defects	30	No significant change (\rightarrow)	2.22	0.136
Selected abdominal wall defects	88	No significant change (\rightarrow)	2.06	0.152
(Q79.2) - Omphalocele / Exomphalos	27	No significant change (\rightarrow)	0.19	0.660
(Q79.3) - Gastroschisis	34	No significant change (\rightarrow)	0.38	0.537
Selected chromosomal defects	172	No significant change (\rightarrow)	1.05	0.306
(Q90) - Down Syndrome	121	No significant change (\rightarrow)	0.36	0.546
(Q91.4-Q91.7) - Trisomy 13 - Patau	11	No significant change (\rightarrow)	0.03	0.868
(Q91.0-Q91.3) - Trisomy 18 - Edwards	24	No significant change (\rightarrow)	0.27	0.600
(Q96) - Turner syndrome	17	No significant change (\rightarrow)	0.15	0.696


























3.6 Appendix 2 - Total birth, number of cases, anomalies and anomalies per case, 2013-2022

Year	Total births	Number of cases	Case rate per 10,000 total births	Number of anomalies	Anomaly rate per 10,000 total births	Anomaly rate per case
2013	8443	191	226.22	218	258.20	1.14
2014	8641	191	221.04	225	260.39	1.18
2015	8160	161	197.30	177	216.91	1.10
2016	8304	154	185.45	184	221.58	1.19
2017	8242	157	190.49	185	224.46	1.18
2018	7939	151	190.20	197	248.14	1.30
2019	8068	180	223.10	221	273.92	1.23
2020	7600	163	214.47	183	240.79	1.12
2021	8139	168	206.41	199	244.50	1.18
2022	7856	148	188.39	168	213.85	1.14


















3.7 Appendix 3 - Prevalence rate per 10,000 total births, aggregated prevalence rate, 2013-2022

Selected anomaly	Number of cases	Aggregated prevalence rate per 10,000 total births (2012-2022)	Prevalence rate per 10,000 total births (2012-2022)
Neural tube defects	64	1.226	
(Q00) - Anencephaly and similar malformations	29	1.215	
(Q01) - Encephalocele	6	1.204	
(Q05) - Spina bifida	29	1.243	
Selected central nervous system defects	108	1.228	
(Q02) - Microcephaly	53	1.229	
(Q03) - Congenital hydrocephalus	50	1.225	
(Q04.1, Q04.2) - Arhinencephaly / Holoprosencephaly	9	1.260	
Selected sense organ defects	28	1.214	

(continued)

Selected anomaly	Number of cases	Aggregated prevalence rate per 10,000 total births (2012-2022)	Prevalence rate per 10,000 total births (2012-2022)
(Q11.0-Q11.2) - Anophtalmos / Microphtalmos	7	1.225	
(Q16.0, Q17.2) - Anotia / Microtia	14	1.218	
(Q30.0) - Choanal atresia	9	1.201	
Selected congenital heart defects	211	1.226	
(Q20.0) - Commom truncus	11	1.213	
(Q20.1, Q20.3, Q20.5) - Transposition of great vessels	59	1.228	
(Q21.2) - Atrioventricular septal defect	71	1.228	
(Q21.3) - Tetralogy of Fallot	33	1.224	
(Q23.4) - Hypoplastic left heart syndrome	28	1.218	
(Q25.1) - Coarctation of aorta	54	1.225	
Oro-facial clefts	102	1.235	
(Q35) - Cleft palate only	67	1.231	
(Q36) - Cleft lip only	18	1.258	
(Q37) - Cleft palate with cleft lip	39	1.231	
Selected gastrointestinal defects	112	1.218	
(Q39.0-Q39.4) - Oesophageal atresia / stenosis, tracheoesophageal fistula	29	1.218	
(Q41) - Small intestine absence / atresia / stenosis	32	1.214	
(Q42.0-Q42.3) - Ano-rectal absence / atresia / stenosis	31	1.224	
(Q43.1) - Hirschsprung disease	17	1.218	
(Q44.2) - Atresia of bile ducts	9	1.205	
Selected genital anomalies	561	1.228	
(Q53.1, Q53.2, Q53.9) - Cryptorchidism / undescended testicles	262	1.232	
(Q54, excluding Q54.4) - Hypospadias	276	1.223	
(Q56) - Indeterminate sex	24	1.237	
(Q64.0) - Epispadias	8	1.240	

(continued)

Selected anomaly	Number of cases	Aggregated prevalence rate per 10,000 total births (2012-2022)	Prevalence rate per 10,000 total births (2012-2022)
Selected urinary tract defects	163	1.228	
(Q60.0-Q60.2) - Renal agenesis	74	1.229	
(Q61.1-Q61.5, Q61.8, Q61.9) - Cystic kidney	79	1.227	
(Q64.1) - Bladder and cloacal exstrophy	4	1.224	
(Q64.2, Q64.3) - Lower urinary tract obstruction	20	1.222	
Hip dysplasia	214	1.223	
(Q65) - Hip dysplasia	214	1.223	
Limb deficiency defects	30	1.211	
(Q79.0) - Diaphragmatic hernia	28	1.239	
(Q71.4-Q71.9, Q72.4-Q72.9, Q73.8, excluding Q71.6, Q71.7, Q72.7) - Limb deficiency defects	30	1.211	
Selected abdominal wall defects	88	1.231	
(Q79.2) - Omphalocele / Exomphalos	27	1.230	
(Q79.3) - Gastroschisis	34	1.225	
Selected chromosomal defects	172	1.225	
(Q90) - Down Syndrome	121	1.225	
(Q91.4-Q91.7) - Trisomy 13 - Patau	11	1.236	
(Q91.0-Q91.3) - Trisomy 18 - Edwards	24	1.230	
(Q96) - Turner syndrome	17	1.210	