

Assignment 2

Math 205 Linear Algebra

Muhammad Shahrom Ali
ma03559

Syed Ammar Mahdi
sm03691

October 10, 2019

1 Answers

1. Linear Independence is a concept applicable to a set of vectors. What it says is that in a set of vectors, if none of the vectors can be expressed as a linear combination of the other vectors (excluding itself, of course), then the set of vectors is linearly independent. It can be mathematically written as

$$\forall \vec{v}_i \in \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \dots, \vec{v}_n\}, \vec{v}_i \neq a_1 \vec{v}_1 + a_2 \vec{v}_2 + \dots + a_{n-1} \vec{v}_{n-1}$$

What it means is that each of the vectors in this set of linearly independent vectors, is contributing to the span of the set. i.e. if a vector was removed, the span of the whole set would be reduced. The set is called ‘linearly’ independent. Intuitively speaking Linear implying in a line, and independence from that would mean no two vectors lie on the same line. But since the concept is general to n dimensions, it doesn't just talk about lying on the same line. Since any vectors that are colinear can be represented as a scalar multiple of the other, we can generalize that as being a linear combination. And hence we can conclude that a set of vectors is linearly dependent if at least one vector in the set can be represented as a linear combination of the other vectors.

If we negate the above mathematical statement it looks something like:

$$\exists \vec{v}_i \in \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \dots, \vec{v}_n\}, \vec{v}_i = a_1 \vec{v}_1 + a_2 \vec{v}_2 + \dots + a_{n-1} \vec{v}_{n-1}$$

$$a_1 \vec{v}_1 + a_2 \vec{v}_2 + \dots + a_{n-1} \vec{v}_{n-1} - \vec{v}_i = 0$$

So this brings us to the formal definition of linear dependence of a set. If the linear combination of the vectors in the set is equal to zero apart

from the trivial solution (all scalars = 0), then the set of vectors is linearly dependent. A set containing the zero vector is always linearly dependent. If a subset of a set of vectors is linearly dependent, then the set is also linearly dependent. [1]

The Basis of a vector space **A** is **the linearly independent set of vectors** such that, the vectors in the set span the whole of **A**. Since it is linearly independent, it is the least number of vectors required to span the space and removal of any of the vectors will reduce the span of the vectors. For any \mathbb{R}^n the bases may be infinitely many but the number of vectors in each of them remains the same. And the number of vectors in the Basis is known as the dimension of the vector space. The dimension of a basis of a vector space in \mathbb{R}^n is n . [2]

a) We form a matrix of these vectors:

$$\begin{bmatrix} 1 & 3 & 1 & -14 \\ 4 & 0 & 1 & 13 \\ 5 & -1 & -2 & 7 \\ 2 & 4 & 1 & -19 \end{bmatrix}$$

Perform following row operations:

$$-4R_1 + R_2$$

$$-5R_1 + R_3$$

$$-2R_1 + R_4$$

$$\frac{1}{3}R_2$$

$$-4R_2 + R_3$$

$$-R_2 + 2R_4$$

$$-\frac{1}{3}R_3 + R_4$$

We get the matrix:

$$\begin{bmatrix} 1 & 3 & 1 & -14 \\ 0 & -4 & -1 & 23 \\ 0 & 0 & -3 & -15 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

The last column can be written as a linear combination of the other 3 vectors, it does not have a pivot, and removing it doesn't impact the span of the vectors. Hence, these are linearly dependent.

dent.

- b) No, because there will be a vector in the matrix such that it can be recreated by a linear combination. In a subspace of \mathbb{R}^m there can be at most m independent vectors and since $n > m$, the columns won't be linearly independent.
- c) Yes. In a subspace of \mathbb{R}^m there can be **at most** m linearly independent vectors. A subset of a linearly independent set of vectors is also linearly independent. And since $n < m$, the columns of the tall matrix can be linearly independent.
- d) Only the trivial solution, Where \vec{x} is the zero vector.
- e) We will see non-zero diagonal elements in \mathbf{A} .
- f) No. We require a set of m vectors which are linearly independent as well, and hence allow to span the whole of the subspace in \mathbb{R}^m .

2. Suppose $\mathbf{A} = \begin{bmatrix} 3 & 6 & 9 \\ 6 & 9 & 12 \\ 9 & 12 & 15 \end{bmatrix}$

- Find $N(\mathbf{A})$
- $\mathbf{B} = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$. Find $N(\mathbf{B})$ and $N(\mathbf{B}^2)$
- What can you say about the relationship between $N(\mathbf{C})$ and $N(\mathbf{C}^2)$. Assume for this part that \mathbf{C} can be any arbitrary square matrix

3. A linear system $\mathbf{A}\vec{x} = \vec{b}$ has special solution of the form

$$\vec{x}_n = x_2 \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} \quad (1)$$

- (a) What are the basis for $N(\mathbf{A})$
 - (b) What is the dimension of $N(\mathbf{A})$
 - (c) Write the reduced row echelon form for \mathbf{A}
 - (d) Describe intuitively the geometry of the solution space
4. Determine with reason which of the following are subspaces of 3×3 matrix \mathbf{M}

- (a) all 3×3 matrices \mathbf{A} such that $\mathbf{A}^T = -\mathbf{A}$
- (b) all 3×3 matrices such that the linear system $\mathbf{A}\vec{x} = \vec{0}$ has only trivial solution
5. For which right sides are these systems solvable? Give your reasons

For all \vec{b} in the $C(\mathbf{A})$, there exists a solution for the system

(a)

$$\begin{bmatrix} 5 & 7 & 5 \\ 10 & 14 & 15 \\ 20 & 28 & 25 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

Applying row operations on \mathbf{A} :

$$-2R_1 + R_2$$

$$-4R_1 + R_3$$

$$-R_3 + R_1$$

$$-R_3 + R_2$$

$$\frac{1}{5}R_1$$

$$\frac{1}{5}R_3$$

$$R_2 \leftrightarrow R_3$$

$$\begin{bmatrix} 1 & \frac{7}{5} & 0 \\ 0 & 0 & 5 \\ 0 & 0 & 0 \end{bmatrix}$$

\vec{a}_1 and \vec{a}_3 are linearly independent and their span is the $C(\mathbf{A})$
hence

$\forall \vec{b} \in C(\mathbf{A})$ the system is solvable.

(b)

$$\begin{bmatrix} 1 & 4 \\ 2 & 9 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

Applying row operations on \mathbf{A}

$$R_1 + R_3$$

$$-2R_1 + R_2$$

$$-4R_2 + R_1$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$\forall \vec{b}$ where \vec{b} is the linear combination of the two columns of \mathbf{A} , the system is solvable.

So speaking in terms of the cartesian coordinate system, all vectors lying on the xy plane, with z component zero.

(c)

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

Since all three vectors of this 3×3 matrix are independent, the system is solvable for all vectors \vec{b} in \mathbb{R}^3 .

(d)

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

In this part, \mathbf{A} has only 2 independent vectors, hence similar to part b, the system is solvable for all vectors \vec{b} which are the linear combinations of \vec{a}_1 and \vec{a}_2 .

6. Given that \mathbf{A} is an arbitrary 4×3 matrix, if we add an extra column \vec{a}_4 to a matrix \mathbf{A} . then the column space gets larger unless Give an example where the column space gets larger and an example where it doesn't.

What should be the condition on \vec{b} for $\mathbf{A}\vec{x} = \vec{b}$ to have a solution if the $C(\mathbf{A})$ doesn't get larger?

7. We begin by making a matrix of all of these vectors.

$$[\vec{v}_1 \quad \vec{v}_2 \quad \vec{v}_3 \quad \vec{v}_4 \quad \vec{v}_5 \quad \vec{v}_6]$$

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 0 & 1 & -1 \end{bmatrix}$$

We then perform row operations on this matrix:

$$R_1 + R_2$$

$$R_2 + R_3$$

$$R_3 + R_4$$

$$\frac{1}{2}R_4$$

The Matrix then becomes

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Out of these we can see $\vec{v}_1, \vec{v}_4, \vec{v}_6$, and \vec{v}_5 to be linearly independent.

So we have atmost 4 linearly independent vectors.

8. Find the bases for the $C(\cdot)$ and $N(\cdot)$ associated with **A** and **B**:

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 2 & 4 & 8 \end{bmatrix}, B = \begin{bmatrix} 1 & 2 & 4 \\ 2 & 5 & 8 \end{bmatrix}$$

9. If **V** is the subspace spanned by $(1, 1, 1)$ and $(2, 1, 0)$, find a matrix **A** that has **V** as its column space and a matrix **B** that has **V** as its nullspace.
10. Find the complete solution for the following equations and describe the solution space:

$$\begin{aligned} x + 3y + 3z &= 0 \\ 2x + 6y + 9z &= 0 \\ -x - 3y + 3z &= 0. \end{aligned} \tag{2}$$

References

- [1] Linear Independence

<https://textbooks.math.gatech.edu/ila/linear-independence.html>

- [2] Bases and Dimensions

<https://textbooks.math.gatech.edu/ila/dimension.html>