

Mineria de Datos

Estudiante UNED

August 2019

2.8

$$\begin{aligned} E_y [E_x [x|y]] &= \int_y \left(\int_x x p(x|y) dx \right) p(y) dy = \int_y \int_x x p(x|y) p(y) dy dx = \int_y \int_x x p(x, y) dy dx = \int_x x \left(\int_y p(x, y) dy \right) dx = \\ &= \int_x p(x) dx = E[x] \\ E_y [var_x [x|y]] + var_y [E_x [x|y]] &= \int_y \left(\int_x (x - E_x [x|y])^2 p(x|y) dx \right) p(y) dy + \int_y (E_y [E_x [x|y]] - E_x [x|y])^2 p(y) dy = \\ &= \int_y \int_x (x^2 - 2xE_x [x|y] + E_x [x|y]^2) p(x|y) p(y) dy dx + \int_y (E_y [E_x [x|y]]^2 - 2E_x [x|y] E_y [E_x [x|y]] + E_y [E_x [x|y]]^2) p(y) dy = \\ &= \int_y \int_x x^2 p(x|y) p(y) dy dx - 2 \int_y \int_x x E_x [x|y] p(x|y) p(y) dy dx + \int_y \int_x E_x [x|y]^2 p(x|y) p(y) dy dx + \int_y E_y [E_x [x|y]]^2 p(y) dy - \\ &= 2 \int_y E_x [x|y] E_y [E_x [x|y]] p(y) dy + \int_y E_y [E_x [x|y]]^2 p(y) dy \end{aligned}$$

4.8

Using (4.57) and (4.58), derive the result (4.65) for the posterior class probability in the two-class generative model with Gaussian densities, and verify the results (4.66) and (4.67) for the parameters w and w_0 .

$$\begin{aligned} p(c_1|x) &= \frac{p(x|c_1)p(c_1)}{p(x|c_1)p(c_1) + p(x|c_2)p(c_2)} = \frac{1}{1 + \exp(-a)} = \sigma(a) \\ a &= \ln \frac{p(x|c_1)p(c_1)}{p(x|c_2)p(c_2)} \\ p(c_1|x) &= \sigma(w^T x + w_0) \\ w &= \Sigma^{-1}(\mu_1 - \mu_2) \\ w_0 &= -\frac{1}{2}\mu_1^T \Sigma^{-1} \mu_1 + \frac{1}{2}\mu_2^T \Sigma^{-1} \mu_2 + \ln \frac{c_1}{c_2} \end{aligned}$$

Entonces

$$\ln \frac{p(x|c_1)p(c_1)}{p(x|c_2)p(c_2)} = w^T x + w_0$$

Teniendo en cuenta que:

$$p(x|c_k) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma|^{1/2}} \exp \left\{ -\frac{1}{2} (x - \mu_k)^T \Sigma^{-1} (x - \mu_k) \right\}$$

$$\ln \frac{p(x|c_1)p(c_1)}{p(x|c_2)p(c_2)} =$$

$$\begin{aligned}
& \ln \left\{ \frac{\frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma|^{1/2}} \exp \left\{ -\frac{1}{2} (x - \mu_1)^T \Sigma^{-1} (x - \mu_1) \right\}}{\frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma|^{1/2}} \exp \left\{ -\frac{1}{2} (x - \mu_2)^T \Sigma^{-1} (x - \mu_2) \right\}} \right\} + \ln \frac{p(c_1)}{p(c_2)} = \\
& \ln \left\{ \exp \left\{ -\frac{1}{2} (x - \mu_1)^T \Sigma^{-1} (x - \mu_1) + \frac{1}{2} (x - \mu_2)^T \Sigma^{-1} (x - \mu_2) \right\} \right\} + \ln \frac{p(c_1)}{p(c_2)} = \\
& -\frac{1}{2} \{ x^T \Sigma^{-1} (x - \mu_1) - \mu_1^T \Sigma^{-1} (x - \mu_1) - x^T \Sigma^{-1} (x - \mu_2) + \mu_2^T \Sigma^{-1} (x - \mu_2) \} + \ln \frac{p(c_1)}{p(c_2)} = \\
& -\frac{1}{2} \{ x^T \Sigma^{-1} x - x^T \Sigma^{-1} \mu_1 - \mu_1^T \Sigma^{-1} x + \mu_1^T \Sigma^{-1} \mu_1 - x^T \Sigma^{-1} x + x^T \Sigma^{-1} \mu_2 + \mu_2^T \Sigma^{-1} x - \mu_2^T \Sigma^{-1} \mu_2 \} + \ln \frac{p(c_1)}{p(c_2)} = \\
& -\frac{1}{2} \{ -x^T \Sigma^{-1} (\mu_1 - \mu_2) - (\mu_1 - \mu_2)^T \Sigma^{-1} x + \mu_1^T \Sigma^{-1} \mu_1 - \mu_2^T \Sigma^{-1} \mu_2 \} + \ln \frac{p(c_1)}{p(c_2)} = \\
& -\frac{1}{2} \{ -2 (\mu_1 - \mu_2)^T \Sigma^{-1} x + \mu_1^T \Sigma^{-1} \mu_1 - \mu_2^T \Sigma^{-1} \mu_2 \} + \ln \frac{p(c_1)}{p(c_2)} = (\mu_1 - \mu_2)^T \Sigma^{-1} x - \frac{1}{2} \mu_1^T \Sigma^{-1} \mu_1 + \frac{1}{2} \mu_2^T \Sigma^{-1} \mu_2 + \\
& \ln \frac{p(c_1)}{p(c_2)} = \\
& \underbrace{(\Sigma^{-1} (\mu_1 - \mu_2))^T x}_{w^T} - \underbrace{\frac{1}{2} \mu_1^T \Sigma^{-1} \mu_1 + \frac{1}{2} \mu_2^T \Sigma^{-1} \mu_2 + \ln \frac{p(c_1)}{p(c_2)}}_{w_0} = \\
& w^T x + w_0
\end{aligned}$$

4.10

Consider the classification model of Exercise 4.9 and now suppose that the class-conditional densities are given by Gaussian distributions with a shared covariance matrix, so that

$$p(\phi_n | C_k) = \mathcal{N}(\phi_n | \mu, \Sigma)$$

Show that the maximum likelihood solution for the mean of the Gaussian distribution for class C_k is given by

$$\mu_k = \frac{1}{N_k} \sum_{n=1}^N t_{n,k} \phi_n$$

which represents the mean of those feature vectors assigned to class C_k . Similarly, show that the maximum likelihood solution for the shared covariance matrix is given by

$$\Sigma = \sum_{k=1}^K \frac{N_k}{N} S_k$$

where

$$S_k = \frac{1}{N_k} \sum_{n=1}^N t_{n,k} (\phi_n - \mu_k) (\phi_n - \mu_k)^T$$

Thus Σ is given by a weighted average of the covariances of the data associated with each class, in which the weighting coefficients are given by the prior probabilities of the classes.

$$p(\{\phi_n, t_n\} | \{\pi_k\}) = \prod_{n=1}^N \prod_{k=1}^K \{p(\phi_n | C_k) p(C_k)\}^{t_{n,k}}$$

$$p(C_k) = \pi_k$$

$$p(\phi_n | C_k) = \mathcal{N}(\phi_n | \mu, \Sigma)$$

$$\ln p(\{\phi_n, t_n\} | \{\pi_k\}) = \ln \prod_{n=1}^N \prod_{k=1}^K \{\mathcal{N}(\phi_n | \mu, \Sigma) \pi_k\}^{t_{n,k}}$$

$$\ln \prod_{m=1}^M a_m = \ln \sum_{m=1}^M a_m$$

$$\ln a^n = n \ln a$$

$$\ln p(\{\phi_n, t_n\} | \{\pi_k\}) = \sum_{n=1}^N \sum_{k=1}^K t_{n,k} \ln \{\mathcal{N}(\phi_n | \mu, \Sigma) \pi_k\}$$

$$\ln \mathcal{N}(\phi_n | \mu, \Sigma) = \ln (2\pi |\Sigma|)^{-1/2} - \frac{1}{2} (\phi_n - \mu_k)^T \Sigma^{-1} (\phi_n - \mu_k)$$

$$\ln p(\{\phi_n, t_n\} | \{\pi_k\}) = \sum_{n=1}^N \sum_{k=1}^K t_{n,k} \left\{ -\frac{1}{2} \ln(|\Sigma|) - \frac{1}{2} (\phi_n - \mu_k)^T \Sigma^{-1} (\phi_n - \mu_k) - \frac{1}{2} \ln(2\pi) + \ln \pi_k \right\}$$

$$\frac{\partial}{\partial \mu_m} \left\{ \sum_{m=1}^m F(x_m) \right\} = \frac{\partial F(x_m)}{\partial \mu_m}$$

$$\frac{\partial}{\partial \mu_k} \{\ln p(\{\phi_n, t_n\} | \{\pi_k\})\} = \frac{\partial}{\partial \mu_k} \left\{ \sum_{n=1}^N t_{n,k} \left\{ -\frac{1}{2} (\phi_n - \mu_k)^T \Sigma^{-1} (\phi_n - \mu_k) \right\} \right\} = 0$$

$$\frac{\partial}{\partial s} (x-s)^T W (x-s) = 2W (x-s)$$

$$\sum_{n=1}^N t_{n,k} \{-\Sigma^{-1}(\phi_n - \mu_k)\} = 0 \implies -\sum_{n=1}^N t_{n,k} \Sigma^{-1} \phi_n - t_{n,k} \Sigma^{-1} \mu_k = 0$$

$$\sum a - b = \sum a - \sum b$$

$$\sum_{n=1}^N t_{n,k} \Sigma^{-1} \mu_k - \sum_{n=1}^N t_{n,k} \Sigma^{-1} \phi_n = 0 \implies \sum_{n=1}^N t_{n,k} \Sigma^{-1} \mu_k = \sum_{n=1}^N t_{n,k} \Sigma^{-1} \phi_n$$

$$\Sigma \cdot \Sigma^{-1} = I$$

$$\Sigma^{-1} \mu_k \sum_{n=1}^N t_{n,k} = \Sigma^{-1} \sum_{n=1}^N t_{n,k} \phi_n \implies \Sigma \left\{ \Sigma^{-1} \mu_k \sum_{n=1}^N t_{n,k} \right\} = \Sigma \left\{ \Sigma^{-1} \sum_{n=1}^N t_{n,k} \phi_n \right\}$$

$$N_k = \sum_{n=1}^N t_{n,k}$$

$$\mu_k N_k = \sum_{n=1}^N t_{n,k} \phi_n \implies \mu_k = \frac{1}{N_k} \sum_{n=1}^N t_{n,k} \phi_n$$

$$\frac{\partial}{\partial \Sigma} \{\ln p(\{\phi_n, t_n\} | \{\pi_k\})\} = \frac{\partial}{\partial \Sigma} \left\{ \sum_{n=1}^N \sum_{k=1}^K t_{n,k} \left\{ -\frac{1}{2} \ln(|\Sigma|) - \frac{1}{2} (\phi_n - \mu_k)^T \Sigma^{-1} (\phi_n - \mu_k) \right\} \right\}$$

$$\begin{aligned} \sum a + b &= \sum a + \sum b \\ D(a + b) &= Da + Db \end{aligned}$$

$$-\frac{1}{2} \frac{\partial}{\partial \Sigma} \left\{ \sum_{n=1}^N \sum_{k=1}^K t_{n,k} \ln(|\Sigma|) \right\} - \frac{1}{2} \frac{\partial}{\partial \Sigma} \left\{ \sum_{n=1}^N \sum_{k=1}^K t_{n,k} (\phi_n - \mu_k)^T \Sigma^{-1} (\phi_n - \mu_k) \right\} = 0$$

$$S_k = \frac{1}{N_k} \sum_{n=1}^N t_{n,k} (\phi_n - \mu_k) (\phi_n - \mu_k)^T$$

$$-\frac{1}{2} \left\{ \sum_{n=1}^N \sum_{k=1}^K t_{n,k} \frac{\partial \ln(|\Sigma|)}{\partial \Sigma} \right\} = \frac{1}{2} \left\{ \sum_{k=1}^K \frac{\partial}{\partial \Sigma} \underbrace{\sum_{n=1}^N t_{n,k} (\phi_n - \mu_k)^T \Sigma^{-1} (\phi_n - \mu_k)}_{N_k \text{ Tr}(\Sigma^{-1} S_k)} \right\}$$

$$\frac{\partial l \ n(|A|)}{\partial \Sigma} = (A^{-1})^T$$

$$-\sum_{n=1}^N \sum_{k=1}^K t_{n,k} \underbrace{(\Sigma^{-1})^T}_{\Sigma^{-1}} = \sum_{k=1}^K \frac{\partial \{ N_k \text{ Tr}(\Sigma^{-1} S_k) \}}{\partial \Sigma}$$

$$\frac{\partial \{ \text{Tr}(AX^{-1}B) \}}{\partial X} = (X^{-1}B \ A \ X^{-1})^T$$

$$-\sum_{n=1}^N \sum_{k=1}^K t_{n,k} \Sigma^{-1} = -\sum_{k=1}^K N_k (\Sigma^{-1} S_k \Sigma^{-1})^T$$

$$(A \ B \ C)^T = C^T B^T A^T \ N = \sum_{n=1}^N \sum_{k=1}^K t_{n,k}$$

$$\Sigma^{-1} \underbrace{\sum_{n=1}^N \sum_{k=1}^K t_{n,k}}_N = \sum_{k=1}^K N_k (\Sigma^{-1} S_k \Sigma^{-1})^T$$

Multiplicando por Σ tanto por la izquierda como por la derecha ambos miembros de la ecuaci3n

$$\Sigma (\Sigma^{-1} N) \Sigma = \Sigma \left(\sum_{k=1}^K N_k \Sigma^{-1} S_k \Sigma^{-1} \right) \Sigma \ N \ \Sigma = \sum_{k=1}^K N_k S_k \implies \Sigma = \sum_{k=1}^K \frac{N_k}{N} S_k$$