



FOR THE REVIEWER

Show in-app crisis support resources

Push notification to admin / care team

Route post or comment

The system is designed to be aggressive on the side of safety

What we need from you:

CLINICIAN SIGN-OFF



Role / credentials

How the System Works

The detection pipeline that wraps every keyword in this document

Date

DETECTION PIPELINE (6 STAGES)

1. **Normalize.** Lowercase, strip leet-speak (h3r0in → heroin), collapse repeated chars (suuuuicide → suicide), remove obfuscation punctuation (k.i.l.l → kill).
2. **Emergency check.** Phrases like "I need help" or "call 988" are treated as *help-seeking*, not harm — they surface crisis resources but don't penalize the user.
3. **Keyword match.** Compare the normalized text against every phrase in this document. Word-boundary aware for single-word matches.
4. **Regex patterns.** Multi-word grammar like "want / wanna / going to + harm verb + self".
5. **Negation downgrade.** "I do NOT want to kill myself" → downgrade by one tier (e.g. high → medium).
6. **Time-immediacy elevation (NEW).** "Tonight I'm going to..." → upgrade by one tier (e.g. medium → high). See last section.

RISK TIERS

Tier	Triggers
High	Push to admin team with elevated priority. Post is held for moderator review (not auto-published). User sees crisis resources sheet immediately.
Medium	Push to admin team for review. Post is held for moderator approval. User sees crisis resources sheet.
Low	Logged for admin awareness. Post may auto-publish (per author's track record). User does NOT see crisis sheet — just awareness flag in admin dashboard.

CATEGORIES

Six categories total. The original four (self-harm, drugs, alcohol, violence) are well-established. **Eating disorder and domestic violence are new in this version** based on the clinical reality that both heavily co-occur with substance use disorder. Please review the new categories with particular care.

Privacy note: Raw user text never leaves the device. The detection happens locally. Only metadata (category, risk level, count of matches) is sent to the server — never the user's actual words.

Self-Harm / Suicide

Direct suicidal ideation, imminent signals, methods of harm, and distress signals across three tiers. Drawn from C-SSRS, SAMHSA crisis indicators, and common young-adult digital-communication patterns.

kill myself	killing myself
kill my self	kms
kys	end my life
ending my life	take my own life
taking my life	commit suicide
committing suicide	suicide attempt
attempted suicide	want to die
wanna die	wanting to die
wish i was dead	wish i were dead
wish i could die	better off dead
better off not existing	don't want to live
no reason to live	nothing to live for
not worth living	life isn't worth it
i'm done	had enough
i've had enough	can't do this anymore
i want to disappear	wish i could disappear
leaving this world	leaving everyone
won't be around	won't be here much longer
world without me	want to sleep forever
never wake up	make it stop
i want it to end	ending my pain
stop the pain	going to overdose on purpose
od on purpose	intentional overdose
take all my pills	swallow all my pills
pills and alcohol	jump off
jumping off	step in front of
in front of a train	hang myself
tie a noose	make a noose
shoot myself	loaded gun
my gun is loaded	cut my wrists
slit my wrists	drink bleach
swallow bleach	carbon monoxide
gas myself	said my goodbyes
saying goodbye	wrote a note

left a note
my last message
last post ever
final post
tell my family i love them
giving my things away
off my meds
haven't eaten in days

won't be here tomorrow
my last post
goodbye everyone
fp (final post)
give my stuff away
stopped taking my meds
haven't slept in days

☐ Approve all ☐ Remove some (see notes) ☐ Add more (see notes)

MEDIUM RISK

~25 phrases

thinking about killing myself
considered suicide
can't go on anymore
don't want to be here anymore
tired of living
tired of being a burden
no escape
nobody would miss me
see no future
thinking about self harm
hurting myself
cutting
started cutting
self harm / self-harm / selfharm
burning myself

thought about ending it
considering suicide
can't keep going
tired of being alive
tired of being here
no way out
everyone would be better without me
what's the point of living
no future for me
thinking about cutting
hurt myself
cutting myself
cut myself
sh (self-harm abbrev)
razor on my skin

☐ Approve all ☐ Remove some (see notes) ☐ Add more (see notes)

feeling hopeless	completely hopeless
hate myself	hating myself
worthless	i am worthless
burden to everyone	i'm a burden
feel like a burden	no one cares
nobody cares about me	no one would notice
all alone	completely alone
so alone	can't take it anymore
don't care anymore	stopped caring
depressed	deeply depressed
severely depressed	empty inside
feel empty	feel nothing
numb	giving up
i give up	ready to give up
suicidal thoughts	intrusive thoughts about dying
dark thoughts	really dark place
dissociating	feel dissociated
having flashbacks	flashbacks
can't feel my body	out of my body

☐ Approve all
 ☐ Remove some (see notes)
 ☐ Add more (see notes)

Phrases the Milton clinical team would add:

⚡ Drugs / Substance Use

Active substance use, relapse confessions, cravings, named substances and slang. The system flags both consumption ("I used") and intent ("about to use") differently.

heroin	dope
smack	junk
china white	fentanyl
fent	fetty
boy (heroin slang)	tar
black tar	shooting up
shoot up	shot up
mainlining	mainline
using a needle	needle drugs
track marks	needle marks
nodding	nodding out
chasing the dragon	speedball
using again	back on drugs
started using again	picked up again
scored	copped
plugged	got my fix
getting my fix	going to use
gonna use	about to use
broke my streak	broke my sober streak
lost my clean time	went back out
overdosed	od'd / oded / od'ed
took too much	too many pills
almost overdosed	nearly od'd
narcan	got narcan'd
shooting meth	smoking meth
crystal meth	tweaking / tweakin
ice (meth slang)	crack cocaine
smoking crack	freebasing
lean	purple drank
sizzurp	robotripping
candy flipping	

⚠ MEDIUM RISK

~40 phrases

craving	cravings
thinking about using	want to use
want to get high	wanna get high
feel like using	miss getting high
miss being high	looking for stuff
trying to score	want to score
triggered to use	triggered today
relapsed	relapsing
had a relapse	fell off
fell off the wagon	slipped
had a slip	went back to using
tempted to use	meth / methamphetamine
crystal	cocaine / coke / blow / snow
bumps	rails
8-ball / eight ball	crack
percocet / percs	oxycontin / oxy
hydrocodone / vicodin	vikes / norco
xanax / bars / benzos	klonopin
adderall misuse	abusing adderall
ecstasy / molly / mdma	rolling
ketamine / k-hole / ket	dmt / ayahuasca
buying drugs	dealer / drug deal
spice / k2	kratom

☐ Approve all ☐ Remove some ☐ Add more**i LOW RISK**

~20 phrases

weed / marijuana / cannabis	pot / bud
high / stoned / blazed	dabs / dabbing / wax
pills	prescription / prescription drugs
plug (dealer slang)	withdrawal / withdrawals
detox	clean time at risk
trigger / triggered	haven't been sober

☐ Approve all ☐ Remove some ☐ Add more

Milton-specific drug slang or patterns to add:



Alcohol

Active drinking confessions, cravings, named beverages, and pre-drinking patterns.



HIGH RISK

~25 phrases

drinking again	started drinking again
had a drink	had drinks
got drunk	drunk
wasted	hammered
trashed	blacked out
black out	blackout
relapsed on alcohol	drank alcohol
drank too much	couldn't stop drinking
back on the bottle	back on the sauce
broke my sobriety	broke sobriety
broke my clean time	not sober anymore
no longer sober	fell off the wagon
dui / dwi	drunk driving

☐ Approve all ☐ Remove some ☐ Add more



MEDIUM RISK

~14 phrases

craving alcohol	craving a drink
want a drink	want to drink
wanted a drink	thinking about drinking
miss drinking	miss alcohol
almost drank	nearly drank
tempted to drink	buzzed
have a buzz	feeling buzzed
drinking problem	pre-game / pregaming
happy hour	going to the bar

☐ Approve all ☐ Remove some ☐ Add more

wine

beer

liquor

spirits

whiskey / vodka

tequila / rum

gin / bourbon / scotch

champagne

bar / bars

shots

going out for drinks

hangover / hangover

☐ Approve all ☐ Remove some ☐ Add more

Violence / Outgoing Threats

Threats made BY the user against others. Distinct from domestic violence (where user reports being hurt by another).

HIGH RISK

~16 phrases

going to kill / gonna kill	want to kill
i'll kill	going to hurt / gonna hurt
going to attack / gonna attack	going to stab / gonna stab
going to shoot / gonna shoot	shoot up the place
shoot up the school	shoot up the office
shoot up work	have a gun
have a knife	armed
make them pay	they will pay
threaten to hurt	

☐ Approve all ☐ Remove some ☐ Add more

MEDIUM RISK

~14 phrases

want to hurt someone	feel like hurting someone
angry enough to hurt	rage / in a rage
blackout rage	violent thoughts
thoughts of violence	can't control my anger
going to lose it	about to snap
going to snap / gonna snap	seeing red
see red	going to do something stupid
threatening	made threats

☐ Approve all ☐ Remove some ☐ Add more

LOW RISK

~6 phrases

angry / furious / enraged	hate them
i hate	want to scream

☐ Approve all ☐ Remove some ☐ Add more



Eating Disorder — NEW

New category. Eating disorders are heavily co-occurring with SUD recovery. Patterns: restriction (anorexia-leaning), binge/purge (bulimia-leaning), explicit body-image distress.



HIGH RISK

~14 phrases

made myself throw up	make myself throw up
making myself vomit	made myself vomit
purging	purged
purge after eating	throwing up after meals
taking laxatives to lose weight	abusing laxatives
using ipecac	haven't eaten in days
not eating	stopped eating
fasting for days	lost too much weight
passed out from not eating	feel like passing out from hunger

☐ Approve all ☐ Remove some ☐ Add more



MEDIUM RISK

~14 phrases

restricting	restricting again
back to restricting	skipping meals
skipping breakfast	fasting again
intermittent fasting too much	counting calories obsessively
obsessed with calories	won't eat
throw up my food	throwing up my food
binge eating / binged	ate too much
binge then purge	binging and purging
starving myself	

☐ Approve all ☐ Remove some ☐ Add more

 LOW RISK

~10 phrases

feel fat / feeling fat

hate my body

disgusted with my body

need to lose weight

weighing myself constantly

anorexia

bulimia

ana (pro-ED slang)

mia (pro-ED slang)

thinspo (pro-ED content)

body checking

☐ Approve all ☐ Remove some ☐ Add more

Clinical question: Should "ana" / "mia" / "thinspo" be elevated from LOW to MEDIUM? They're code words used in active pro-eating-disorder online communities. Their appearance may signal a user engaging with harmful content. Defer to clinical judgment.



Domestic Violence — Incoming — NEW

New category. For users reporting violence done TO them by partner, family, or household. Triggers care-team outreach and safe-call escalation.



HIGH RISK

~25 phrases

he hit me

they hit me

she beat me

he punched me

he choked me

he strangled me

he broke

broke my nose

threw me against

she kicked me

said he'll kill me

threatened to kill me

pulled a gun on me

held a knife to

said he'll hurt me

not safe in my house

scared for my life

can't leave him

he locked me in

she hit me

he beat me

got beat up

she punched me

she choked me

strangled me

broke my arm

he threw me

he kicked me

going to kill me

said she'll kill me

threatened me with a gun

threatened me with a knife

going to hurt me

not safe at home

afraid for my life

trapped here

can't leave her

she locked me in

☐ Approve all ☐ Remove some ☐ Add more

⚠️ MEDIUM RISK

~18 phrases

abusive partner	abusive husband
abusive wife	abusive boyfriend
abusive girlfriend	he hurts me
she hurts me	afraid of him
afraid of her	scared of him
scared of her	he yells at me every day
controlling everything i do	won't let me leave
won't let me see	takes my phone away
won't let me have money	stalking me
he stalks me	she stalks me
following me everywhere	he pushed me
she pushed me	he shoved me
she shoved me	grabbed me hard

☐ Approve all ☐ Remove some ☐ Add more

📁 LOW RISK

~8 phrases

afraid to go home	walking on eggshells
verbally abusive	name-calling
calls me names	controlling
he won't talk to me	she won't talk to me
isolating me from friends	isolating me from family

☐ Approve all ☐ Remove some ☐ Add more

Clinical question: When this fires, what should admin do? Recommend: trained-staff outreach within 1 hour, NEVER contact via SMS/email (could be intercepted), use the in-app care-team chat which is encrypted. Should the system include the National DV Hotline (1-800-799-7233) in the in-app crisis resources sheet when this category fires?



Emergency Help-Seeking + Time-Immediacy

Two special-purpose lists that modify how the system responds.

EMERGENCY HELP-SEEKING PHRASES (NEVER FLAG AS HARM)

When any of these appears, the system **shows the crisis resources sheet** (988, Crisis Text Line, SAMHSA, Milton FL/OH, 911) and notifies admins for follow-up — but does NOT penalize the user, hold their post, or flag it as harm. The user is reaching out for help; we want to surface help.



HELP-SEEKING

~16 phrases

call 988	988
suicide hotline	crisis line
crisis hotline	call 911
call the police	need help now
please help me	i need help
help me please	i'm in crisis
im in crisis	crisis text line
samhsa	national helpline
going to call for help	reaching out for help
texting a crisis line	

☐ Approve all ☐ Add more (e.g. Milton-specific safe words)

TIME-IMMEDIACY MARKERS (RISK ELEVATION)

When any of these appears anywhere in the same text as a risk match, the system **elevates that match's risk by one tier** (e.g. medium → high). Caps at high risk. Rationale: "I'm cutting tonight" is materially different from "I've thought about cutting before" — the time-boundedness signals imminent action.



TIME ELEVATION

~24 markers

tonight	today
right now	right this moment
in an hour	in a few hours
in 30 minutes	in 15 minutes
this morning	this afternoon
this evening	after work
after school	when i get home
this weekend	tomorrow morning
as soon as	any minute now
minutes away	hours away
i'm doing it / im doing it	doing it now
about to	

☐ Approve all ☐ Adjust (some too aggressive?) ☐ Add more

Example: A user writes "*I'm cutting tonight.*"

Without time elevation: matches "cutting" → MEDIUM risk → admin sees notification within 1 hour.

With time elevation: matches "cutting" + sees "tonight" → upgrades to HIGH risk → push to admin with elevated priority, post held for review, crisis resources surfaced.

✓ Final Sign-Off

Clinical review summary and approval

SUMMARY OF CHANGES FROM CLINICAL REVIEW

Phrases added:

Phrases removed:

Risk tier adjustments (which phrase moved to which tier):

Comments / clinical notes:

CLINICIAN APPROVAL

Reviewer name (printed)

Role / credentials

Signature

Date

ONCE SIGNED

Return this document to Ezra at **media@miltonjefferson.com** or **ebarish@miltonhealthgroup.com**. We'll then update the keyword list in the production Supabase database (no app rebuild required — changes take effect within 4 hours on every user's device).