

Overview

Systems-level approach to develop a body schema and agency:

- The **basal ganglia** will select an action based on the desired state (see Baladron and Hamker, 2020).
- The **central pattern generator** will be execute the action (see Nassour et al., 2020).
- The **cerebellum** will learn to predict the sensory consequences of the motor action in all modalities (vision, touch, proprioception) i.e. the body schema (see Schmid et al., 2019).
- The **prediction error** will be used to improve the prediction and to train action selection in the **basal ganglia**.




Figure 1: babbas

Recurrent Basis Functions¹

Hier was zu dieser Veröffentlichung Pouget et al., 2002.




Figure 2: babbas

Learning a Body Schema¹

Basal Ganglia²

The 3-factor learning principles are primarily determined by presynaptic and postsynaptic neuron activity, as well as the dopamine signal.

The labels high and low indicate whether the pre- and post-activity is more than or less than a given threshold (for example, mean population activity).




Figure 3: Modeling of segregated basal ganglia pathways

DA+ and DA- labels indicate if SNc activity is above or below a given threshold. A sign specifies the weight changes in the relevant projections for each combination of these criteria (+, no sign). Table S8 contains the real mathematical explanation of plasticity, in addition to the general description offered.

		Dopamine				
		DA +		DA -		
		Post-activity				
		High	Low	High	Low	
Pre-activity	High	+		-		Cortex-D1
	Low	-				
	High	-		+		Cortex-D2
	Low		-			
	High	+		-		Cortex-STN
	Low	-				
	High	-	+		-	D1-SNr
	Low					
	High		-	-	+	D2-GPe
	Low					
	High	+	-		+	STN-SNr
	Low					

Table 1: +: LTP; -: LTD; no sign: no weight change

Motor Learning with the Basal Ganglia²





Figure 4: Right: Development of motor learning in the different pathways.



The 3-factor learning principles are primarily determined by presynaptic and postsynaptic neuron activity, as well as the dopamine signal. The labels high and low indicate whether the pre- and post-activity is

more than or less than a given threshold (for example, mean population activity). DA+ and DA- labels indicate if SNc activity is above or below a given threshold.

References

- Baladron, J., & Hamker, F. H. (2020). Habit learning in hierarchical cortex–basal ganglia loops. *European Journal of Neuroscience*, 52(12), 4613–4638. <https://doi.org/10.1111/ejn.14730>
- Nassour, J., Duy Hoa, T., Atoofi, P., & Hamker, F. (2020). Concrete Action Representation Model: From Neuroscience to Robotics. *IEEE Transactions on Cognitive and Developmental Systems*, 12(2), 272–284. <https://doi.org/10.1109/TCDS.2019.2896300>
- Pouget, A., Deneve, S., & Duhamel, J.-R. (2002). A computational perspective on the neural basis of multisensory spatial representations. *Nature Reviews Neuroscience*, 3(9), 741–747. <https://doi.org/10.1038/nrn914>
- Schmid, K., Vitay, J., & Hamker, F. H. (2019). Forward Models in the Cerebellum using Reservoirs and Perturbation Learning. *2019 Conference on Cognitive Computational Neuroscience*. <https://doi.org/10.32470/CCN.2019.1139-0>