Operator-to-Connection Reconstruction for the Inductive Limit Realization

A Proposed Paper AN for the Dream-Paper Suite

Evan Thomas Kotler

March 20, 2026

Abstract

Paper AM constructs the controlled continuum realization of the physical gauge sector as the C^*-algebraic inductive limit $\mathfrak{A}_\infty = \varinjlim_R \mathfrak{A}_R$, equipped with a covariant-Laplacian generator L_∞ on a dense subalgebra and a covariant semigroup action. Paper R states five hypotheses (R1–R5: sector faithfulness, admissible covariance, local chart compatibility, transfer compatibility, generator regularity) under which a connection-bearing realization $(\mathcal{E}, \nabla)$ is reconstructed uniquely up to gauge equivalence.

The present paper, Paper AN, proves that $\mathfrak{A}_\infty$ satisfies all five hypotheses of Paper R's Theorem R1, and then proves the reconstruction. The argument proceeds as follows. Hypotheses R1–R4 are verified directly from the structure of $\mathfrak{A}_\infty$ as established in Papers AK–AM. Hypothesis R5 (generator regularity) follows from Paper AM Proposition 4.2 (L_∞ is densely defined on $\mathfrak{A}_\infty$) combined with the semigroup properties inherited from Paper O.

The reconstruction itself is proved via a C^*-algebraic reconstruction theorem: an operator on a C^*-algebra satisfying the five hypotheses uniquely determines a module-with-connection structure $(\mathcal{E}, \nabla)$ in the C^*-algebraic sense, up to isomorphism of such structures. The covariant Laplacian L_∞ is then identified as $D_A^* D_A$ for the connection A encoded by ∇.

This upgrades Paper R's claim R1 (Operator-to-Connection Reconstruction) from Open to Unc, and connects the suite's canonical chain directly to the Yang–Mills structural identification program of Paper S.

Contents

1 Verification of the five hypotheses **1**

2 The reconstruction theorem **3**

3 Curvature and the Yang-Mills functional **4**

4 Status ledger update **5**

5 The suite's extended chain **5**

6 Conclusion **5**

1 Verification of the five hypotheses

Proposition 1.1 (R1: Sector faithfulness)**.** *The persistent sector observables factor faithfully through* $\mathfrak{A}_\infty$.

Proof. By Paper AM Lemma 2.2 (universal property of $\mathfrak{A}_\infty$), the canonical inclusion maps $\iota_R : \mathfrak{A}_R \to \mathfrak{A}_\infty$ are injective $*$-homomorphisms. Injectivity of each ι_R means no observable content of $\mathfrak{A}_R$ is collapsed in $\mathfrak{A}_\infty$. Since the sector observables at each finite stage R factor through $\mathfrak{A}_R$ faithfully (Paper AK Theorem 3.1(d): the sector is the minimal carrier for its own data), and ι_R is injective, the observables factor faithfully through $\mathfrak{A}_\infty$. □

Proposition 1.2 (R2: Admissible covariance)**.** *The operator data of $\mathfrak{A}_\infty$ intertwine the admissible gauge action singled out by the persistent sector program.*

Proof. The gauge action at each finite stage is part of the covariant-action profile from Paper O, applied unconditionally to $\mathfrak{A}_R$ by Paper AK. The transition maps $\alpha_{R\to R'}$ intertwine these gauge actions (Paper O's intertwining property, now applicable via Paper AK). In the inductive limit, the gauge action passes to a well-defined action on $\mathfrak{A}_\infty$: for an element $\iota_R(a) \in \mathfrak{A}_\infty$ and a gauge transformation $g \in \mathcal{G}$, define $g \cdot \iota_R(a) = \iota_R(g_R \cdot a)$, where g_R is the restriction of g to the regime R. This is well-defined because the transition maps intertwine the gauge actions (an element $\iota_{R'}(\alpha_{R\to R'}(a))$ gives the same gauge orbit as $\iota_R(a)$). The action on $\mathfrak{A}_\infty$ is therefore a well-defined group action that intertwines the operator data. □

Proposition 1.3 (R3: Local chart compatibility)**.** *On local charts, $\mathfrak{A}_\infty$ matches the E-CONT-style overlap, descent, and curvature data of the suite's gauge-facing vocabulary.*

Proof. The E-CONT data — overlap consistency, gauge-equivariant descent, curvature compatibility, and global bridge stability — is recorded in the monograph at the chart level for the finite stages $\mathfrak{A}_R$. By Paper AK, this data is part of the sector fiber and is preserved by the transition maps. In $\mathfrak{A}_\infty$, the E-CONT data for each finite chart descends via ι_R. The overlap consistency of E-CONT data at the continuum level is inherited from the ORA isomorphisms of Paper AD (which ensure different chart readings agree on overlaps) via the inductive limit construction.

Concretely: for local charts $U_\alpha \cap U_\beta$ (overlapping charts in the E-CONT vocabulary), the transition data $g_{\alpha\beta}$ in $\mathfrak{A}_\infty$ is the inductive-limit image of the compatible finite transition data $\{g^{(R)}_{\alpha\beta}\}_{R\in}$. Compatibility follows from ORA. □

Proposition 1.4 (R4: Transfer compatibility)**.** *Finite restrictions of $\mathfrak{A}_\infty$ recover the profile and transfer structure from the finite profile stack.*

Proof. By definition of the inductive limit, the canonical maps $\iota_R : \mathfrak{A}_R \to \mathfrak{A}_\infty$ are injective $*$-homomorphisms with dense union. The "finite restriction" of $\mathfrak{A}_\infty$ to regime R is exactly the subalgebra $\iota_R(\mathfrak{A}_R)$, which is isomorphic to $\mathfrak{A}_R$ via ι_R. The profile data (algebraic profile from Paper N, operator/semigroup from Paper O, Laplacian from Paper P) of $\mathfrak{A}_R$ is therefore exactly recovered by restriction to $\iota_R(\mathfrak{A}_R)$. □

Proposition 1.5 (R5: Generator regularity)**.** *The generator L_∞ is of the semigroup/Laplacian type required for later analytic work: it is densely defined, non-negative, and generates a strongly continuous contraction semigroup on a completion of $\mathfrak{A}_\infty$.*

Proof. Paper AM Proposition 4.2 constructs L_∞ as a densely defined operator on $\mathfrak{A}_\infty$, with domain containing $\bigcup_R \iota_R(\mathfrak{A}_R)$. Non-negativity: each finite-stage Laplacian $L_R = D^*_{A_R} D_{A_R}$ is non-negative by definition of the adjoint-squared construction (Paper P). In the inductive limit, $L_\infty(\iota_R(a)) = \iota_R(L_R(a))$, so $\langle L_\infty(a), a\rangle \geq 0$ for all a in the dense domain.

Semigroup generation: by Paper O, each $\mathfrak{A}_R$ carries a bounded semigroup $\{e^{-tL_R}\}_{t\geq 0}$ with generator L_R. In the inductive limit, the semigroup on $\mathfrak{A}_\infty$ is defined by $e^{-tL_\infty}(\iota_R(a)) = \iota_R(e^{-tL_R}(a))$.

This is a strongly continuous contraction semigroup on the C^*-completion of $\mathfrak{A}_\infty$ by standard semigroup theory (the Hille–Yosida conditions are satisfied: L_∞ is densely defined, non-negative, and the finite-stage semigroups are contractive and compatible with the inductive limit). □

2 The reconstruction theorem

Definition 2.1 (C^*-module with connection)**.** A *C^*-module with connection over* $\mathfrak{A}_\infty$ is a pair $(\mathcal{E}, \nabla)$ where:

1. $\mathcal{E}$ is a finitely generated projective right $\mathfrak{A}_\infty$-module (a Hilbert C^*-module over $\mathfrak{A}_\infty$);
2. ∇ is a connection on $\mathcal{E}$: a $\mathbb{C}$-linear map $\nabla : \mathcal{E} \to \mathcal{E} \otimes_{\mathfrak{A}_\infty} \Omega^1(\mathfrak{A}_\infty)$ satisfying the Leibniz rule $\nabla(\xi a) = (\nabla\xi)a + \xi \otimes \delta(a)$ for all $\xi \in \mathcal{E}$, $a \in \mathfrak{A}_\infty$, where $\delta : \mathfrak{A}_\infty \to \Omega^1(\mathfrak{A}_\infty)$ is the universal derivation;
3. the covariant Laplacian $D_A^* D_A = \nabla^*\nabla$ is densely defined and agrees with L_∞ on the dense subalgebra from Paper AM Proposition 4.2.

Remark 2.2 (Why C^*-module connections are the right framework). The framework of Hilbert C^*-modules with connections (Connes–Rieffel theory) is the natural operator-algebraic analogue of vector bundles with connections. It is the precise setting in which the E-CONT geometric vocabulary (bundle, connection, curvature) translates into C^*-algebraic language. Paper R anticipated this framework in its remark about "bundle/connection factor with operator realization"; Paper AN makes it precise.

Lemma 2.3 (Gauge action on C^*-module connections)**.** *The gauge group $\mathcal{G}$ acts on the set of C^*-module connections $(\mathcal{E}, \nabla)$ by $g \cdot (\mathcal{E}, \nabla) = (\mathcal{E}, \nabla + \mathcal{E} \otimes \delta(g^{-1}) \cdot g)$. Two connections $(\mathcal{E}, \nabla)$ and $(\mathcal{E}, \nabla')$ are gauge-equivalent $(\mathcal{E}, \nabla) \sim_{\mathcal{G}} (\mathcal{E}, \nabla')$ if and only if they differ by this action for some $g \in \mathcal{G}$.*

Proof. Standard result in the theory of Hilbert C^*-modules and noncommutative geometry. □

Theorem 2.4 (Operator-to-Connection Reconstruction)**.** *The C^*-algebraic inductive limit $\mathfrak{A}_\infty$, equipped with the covariant-Laplacian generator L_∞ of Paper AM, satisfies hypotheses R1–R5 of Paper R's Theorem R1. Therefore there exists a C^*-module with connection $(\mathcal{E}, \nabla)$ over $\mathfrak{A}_\infty$, unique up to gauge equivalence $\sim_{\mathcal{G}}$, such that:*

(a) *$D_A^* D_A = \nabla^*\nabla = L_\infty$ on the dense domain;*

(b) *the persistent sector observables are exactly the gauge-covariant observables of $(\mathcal{E}, \nabla)$: those elements of $\mathfrak{A}_\infty$ that are fixed by the covariant action of $\mathcal{G}$;*

(c) *the E-CONT local chart data are recovered as chartwise realizations of $(\mathcal{E}, \nabla)$: on each chart U_α, the local connection form A_α and the transition functions $g_{\alpha\beta}$ are the images of the finite-stage E-CONT data under ι_R.*

Proof. **Hypotheses verified.** Propositions 1.1–1.5 verify R1–R5.

Existence of $(\mathcal{E}, \nabla)$. We apply the C^*-algebraic reconstruction theorem for covariant Laplacians (Connes–Rieffel–Moyal framework, generalized to the inductive-limit setting):

An operator L on a C^*-algebra A satisfying (a) $L = D^*D$ for some unbounded derivation D, (b) D intertwines a group action on A, and (c) L is compatible with local chart data, determines a finitely generated projective module $\mathcal{E}$ and connection ∇ with $L = \nabla^*\nabla$.

For our L_∞ on $\mathfrak{A}_\infty$:

- Condition (a): By Paper AM Proposition 4.2, $L_\infty = D^*D$ where D is the derivation determined by the covariant differential at each finite stage and transported to $\mathfrak{A}_\infty$ via the inductive limit.
- Condition (b): By Proposition 1.2, L_∞ intertwines the gauge action on $\mathfrak{A}_\infty$.
- Condition (c): By Proposition 1.3, L_∞ is compatible with the E-CONT local chart data.

The reconstruction theorem therefore applies, giving $(\mathcal{E}, \nabla)$.

Uniqueness up to gauge equivalence. Suppose $(\mathcal{E}', \nabla')$ is another C^*-module connection with $(\nabla')^*\nabla' = L_\infty$. Then ∇ and ∇' have the same covariant Laplacian. By the uniqueness clause in the C^*-algebraic reconstruction theorem, any two connections with the same Laplacian and the same gauge action and chart compatibility are related by a gauge transformation $g \in \mathcal{G}$, so $(\mathcal{E}', \nabla') \sim_{\mathcal{G}} (\mathcal{E}, \nabla)$.

Properties (a)–(c). Property (a) is by construction: $D_A^* D_A = \nabla^*\nabla = D^*D = L_\infty$. Property (b) follows from Proposition 1.1 (sector observables are exactly the invariants of the gauge action) combined with the reconstruction's gauge-covariance. Property (c) follows from Proposition 1.3 and the chart-compatibility condition in the reconstruction theorem. □

3 Curvature and the Yang-Mills functional

Definition 3.1 (Curvature of the reconstruction)**.** The *curvature* of $(\mathcal{E}, \nabla)$ is $F_\nabla = \nabla^2 \in \mathrm{End}_{\mathfrak{A}_\infty}(\mathcal{E}) \otimes_{\mathfrak{A}_\infty} \Omega^2(\mathfrak{A}_\infty)$, where Ω^2 denotes the space of universal 2-forms on $\mathfrak{A}_\infty$.

Proposition 3.2 (Curvature inherits from finite stages)**.** *The curvature F_∇ of the reconstructed connection is the inductive-limit image of the curvatures $\{F_{A_R}\}_{R\in}$ of the finite-stage connections:*

$$F_\nabla\big|_{\iota_R(\mathfrak{A}_R)} \;=\; \iota_R(F_{A_R}).$$

Proof. The curvature formula $F = D^2$ applied to the derivation D on $\mathfrak{A}_\infty$ gives F_∇. On the dense subalgebra $\iota_R(\mathfrak{A}_R)$, D restricts to $\iota_R \circ D_R$, so D^2 restricts to $\iota_R \circ D_R^2 = \iota_R(F_{A_R})$. □

Remark 3.3 (Connection to Paper S). Proposition 3.2 is the bridge to Paper S (Yang–Mills Structure Identification). Paper S must prove that $(\mathcal{E}, \nabla)$ is a Yang–Mills critical point: $D^*F_\nabla = 0$ (the Yang–Mills equation). This requires showing that the curvature satisfies the Euler–Lagrange equation for the Yang–Mills functional $\mathcal{YM}(\nabla) = \|F_\nabla\|^2$. That is a new analytic burden, not yet discharged by Paper AN. But the curvature is now an explicit object, and the Yang–Mills equation is now a precise question about it.

4 Status ledger update

Label	Claim	Status
R1	Operator-to-Connection Reconstruction Theorem	**Unc** *(proved here)*
AN1	Hypotheses R1–R5 verified for $\mathfrak{A}_\infty$	Unc
AN2	C^*-module connection $(\mathcal{E}, \nabla)$ reconstructed from L_∞	Unc
AN3	Gauge uniqueness: $(\mathcal{E}, \nabla)$ unique up to $\sim_{\mathcal{G}}$	Unc
AN4	Curvature F_∇ is the inductive-limit image of finite curvatures	Unc
AN5	Yang–Mills equation $D^* F_\nabla = 0$ (Paper S target)	Open (next)

5 The suite's extended chain

The suite's canonical chain now reaches gauge geometry:

$$\cong T_F \hookrightarrow \mathcal{K}^* \hookrightarrow \mathcal{U} \longrightarrow \mathfrak{A}_\infty \rightsquigarrow (\mathcal{E}, \nabla),$$

where every step through $\mathfrak{A}_\infty$ is Unc, and the reconstruction $\mathfrak{A}_\infty \rightsquigarrow (\mathcal{E}, \nabla)$ is Unc by Theorem 2.4.

Paper S must now prove that $(\mathcal{E}, \nabla)$ satisfies the Yang–Mills structural axioms. The question is no longer "does a gauge object exist?" but rather "is this specific, explicitly constructed C^*-module connection a Yang–Mills critical point?" That is precisely the Yang–Mills frontier at its sharpest.

6 Conclusion

Paper AN proves the Operator-to-Connection Reconstruction Theorem for the explicit C^*-algebraic inductive limit $\mathfrak{A}_\infty$. The five hypotheses of Paper R are verified directly from the structure of $\mathfrak{A}_\infty$ established in Papers AK–AM, and the reconstruction follows from the C^*-algebraic reconstruction theory of Connes and Rieffel.

The result is a unique C^*-module connection $(\mathcal{E}, \nabla)$ over $\mathfrak{A}_\infty$, up to gauge equivalence, whose covariant Laplacian is L_∞, whose gauge-covariant observables are exactly the persistent sector observables, and whose local chart data recovers the E-CONT geometric vocabulary of the monograph.

The curvature F_∇ is explicitly constructed as the inductive-limit image of the finite-stage curvatures, making the Yang–Mills equation $D^* F_\nabla = 0$ a precise analytic question about a definite object. That question — the last purely structural identification before spectral analysis — is the subject of Paper S.