Percentiles of Normal(0,1) distribution

$\mathbb{P}(Z \leq z)$	0.5	0.75	0.9	0.95	0.975	0.99	0.995
z	0	0.67	1.28	1.64	1.96	2.33	2.58

Percentiles of t distribution

Degrees of freedom:	0.5	0.75	0.9	0.95	0.975	0.99	0.995
1	0	1.00	3.08	6.31	12.71	31.82	63.66
5	0	0.73	1.48	2.02	2.57	3.36	4.03
7	0	0.71	1.41	1.89	2.36	3.00	3.50
8	0	0.71	1.40	1.86	2.31	2.90	3.36
9	0	0.70	1.38	1.83	2.26	2.82	3.25
10	0	0.70	1.37	1.81	2.23	2.76	3.17

Section 1: True or false? Circle the appropriate choice (1 point each).

1. (TRUE / FALSE) In a simple linear regression model, the expected value of Y when X has the value x^* is $\beta_0 + \beta_1 x^*$.

2. (TRUE / FALSE) The value of the correlation between two variables depends on which variable is labeled X and which is labeled Y.

3. (TRUE / FALSE) If the data $\{(x_1, y_1), \ldots, (x_n, y_n)\}$ are a random sample from a population for which the simple linear regression model is true, the least squares estimator of the slope provides an unbiased estimate of the population slope β_1, meaning there is no systematic tendency for $\hat{\beta}_1$ to be too high or too low.

4. (TRUE / FALSE) For a simple linear regression model, if a hypothesis test fails to reject the null hypothesis that the slope equals zero, you can conclude there is no relationship between X and Y.

5. (TRUE / FALSE) For any simple linear regression model we fit to data, the sum of squared residuals $\sum_{i=1}^{n}(y_i - \hat{y}_i)^2$ will always be less than or equal to the total sum of squares $\sum_{i=1}^{n}(y_i - \bar{y}_i)^2$.

6. (TRUE / FALSE) To check the linear regression assumption that the error terms have an expected value of zero, we can compute the mean of the residuals. If the mean of the residuals is not close to zero, this indicates a violation of the model assumptions.

7. (TRUE / FALSE) We are interested in estimating μ, the true average amount of sleep per night among adults in the US. Based on a random sample, we obtain a 95% confidence interval of 6.8 to 7.3 hours. Thus, we can conclude there is a 95% probability that the true average sleep duration is between 6.8 and 7.3 hours.

8. (TRUE / FALSE) One advantage of linear regression is that it allows us to make reliable predictions for Y at X values beyond the range in the observed data.

9. (TRUE / FALSE) When estimating a population proportion p, confidence interval widths are based on the t-distribution rather than the standard normal distribution, to account for the extra uncertainty that comes from not knowing the true standard deviation.

10. (TRUE / FALSE) The Central Limit Theorem provides a basis for statistical inference by guaranteeing that, in sufficiently large random samples, the sample values approximately follow a normal distribution.

Section 2: Circle the appropriate choice (2 points each).

11. Suppose X and Y are random variables, and that the variance of X is 16, and the variance of Y is 9. What is the standard deviation of X-Y?

    (a) 7

    (b) 5

    (c) 6.5

    (d) 25

    (e) cannot be determined with the information given

12. Suppose an internet service provider reports that 30% of its customers are dissatisfied, but we believe the true proportion is higher, and we want to test this. We sample 200 customers, and 40% of them report being dissatisfied. What are the appropriate null and alternative hypotheses for our question?

    (a) $H_0 : p = 0.4,\ H_a : p > 0.4$

    (b) $H_0 : \hat{p} = 0.4,\ H_a : \hat{p} > 0.4$

    (c) $H_0 : \hat{p} = 0.3,\ H_a : \hat{p} > 0.3$

    (d) $H_0 : p = 0.3,\ H_a : p > 0.3$

13. Suppose a hypothesis test yields a p-value of 0.03. Which of the following is the best interpretation?

    (a) There is a 3% chance that the null hypothesis is true.

    (b) There is a 3% chance of observing data this extreme or more extreme if the null hypothesis is true.

    (c) There is a 97% chance the alternative hypothesis is true.

    (d) The alternative hypothesis is 97% more likely to be true than the null hypothesis.

14. An experimenter performed a hypothesis test with a two-sided alternative hypothesis on a set of data and obtained a p-value of 0.44. For the same data, if the experimenter used a one-sided alternative instead, which of the following is true about the possible p-value(s) that the experimenter could have obtained?

    (a) the only possible p-value is 0.22


    (b) the only possible p-value is 0.44

    (c) the only possible p-value is 0.88

    (d) the possible p-values are 0.22 and 0.78

    (e) the possible p-values are 0.22 and 0.88

15. Suppose we fit a linear regression model to the data depicted in the scatter plot below, using X as the predictor and Y as the response, and obtain an estimated slope of $\hat{\beta}_1$.



If we reversed the variables, fitting a linear regression model using Y as the predictor and X as the response, the new estimated slope would be:


(a) equal to $\frac{1}{\hat{\beta}_1}$

(b) less than $\frac{1}{\hat{\beta}_1}$

(c) greater than $\frac{1}{\hat{\beta}_1}$

(d) cannot be determined with the information given

Questions 16-17 are related to the simple linear regression model fit to data depicted in the scatter plot below.



16. Which of the following will be wider:

(a) 99% confidence interval for the average Y value when X=5

(b) 99% prediction interval for an individual Y value when X=5

(c) cannot be determined with the information given

17. Which of the following will be wider:

    (a) 95% confidence interval for the average Y value when X=11

    (b) 95% prediction interval for an individual Y value when X=6

    (c) cannot be determined with the information given

Section 3: Answer all questions in the space provided.

1. Central City Power computes the monthly cost of electricity for a customer as:

$$\text{Cost} = 0.12^{*}\text{usage} + 15,$$

   where cost is measured in dollars, and usage is measured in kilowatt-hours. The average usage each month is 1000, with a standard deviation of 50. What are the mean and standard deviation of the monthly cost?

2. An animal shelter finds that small dogs tend to be adopted more quickly than large dogs. Fitting a linear regression model using weight to predict days to adoption, they obtain a significant positive slope. The shelter director concludes that they should prioritize helping dogs lose weight in order to shorten their time to adoption. From a statistical standpoint, is this a reasonable conclusion? Explain why or why not.

3. Suppose that the height of mature pine trees is normally distributed, with standard deviation of 20 feet. In a random sample of 10 trees, heights had a sample average of 110 feet and sample standard deviation of 14. Give a 90% confidence interval for the true average height of pine trees.

4. The output below shows the results of fitting a linear regression model to predict zinc concentration (in parts per million, "ppm") from lead concentration (in ppm) in fish from a certain river. The analysis is based on data from 80 fish.

```
lm(formula = zinc_ppm ~ lead_ppm)
Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)    16.3     8.62       1.89    0.059
lead_ppm       19.0     3.67       5.18    0.000
R-squared:  0.82
```


   (a) What zinc concentration does the model predict for a fish with a lead concentration of 0.8 ppm?

   (b) What is the residual for an observed fish with a lead concentration of 2 ppm and zinc concentration of 52 ppm?


(c) Find a 99% confidence interval for the slope of the population regression line. Describe the interpretation of the confidence interval.

(d) Describe the interpretation of the R^2 value given above.

5. (8 points) The plots below show the residuals from two different fitted simple linear regression models.



Plot 1



Plot 2

(a) In plot 1, which regression assumptions appear to be violated, if any?

(b) For the violated assumptions identified above (if any), what are the consequences for estimation of the linear model coefficients, and for inference (confidence intervals and hypothesis tests) based on the model?

(c) In plot 2, which regression assumptions appear to be violated, if any?

(d) For the violated assumptions identified above (if any), what are the consequences for estimation of the linear model coefficients, and for inference based on the model?