

CS 480/680

Introduction to Machine Learning

Lecture 7

Support Vector Machines Part II

Soft Margin Classifier

Kathryn Simone

1 October 2024

Interlude: Revising optimization with Lagrangian multipliers

Consider the constrained objective

$$\begin{aligned} \min y(x) &= 0.2x^2 - x + 1 \\ \text{subject to: } &x \geq 5 \end{aligned}$$


This can be transformed into the *dual* problem



$$\begin{aligned} \max \mathcal{L}(x, \lambda) &= 0.2x^2 - x + 1 - \lambda(x - 5) \\ \text{subject to: } &\lambda \geq 0 \end{aligned}$$

This can be transformed into the *dual* problem

$$\min_x \max_{\lambda \geq 0} \mathcal{L}(x, \lambda) = 0.2x^2 - x + 1 - \lambda(x - 5)$$



Interlude: Revising optimization with Lagrangian multipliers

Consider the constrained objective

$$\begin{aligned}\min y(x) &= 0.2x^2 - x + 1 \\ \text{subject to: } &x \geq 5\end{aligned}$$

This can be transformed into the *dual* problem

$$\min_x \max_{\lambda \geq 0} \mathcal{L}(x, \lambda) = 0.2x^2 - x + 1 - \lambda(x - 5)$$

Where λ is a Lagrange multiplier. The dual problem is constructed by first minimizing the Lagrangian with respect to x , and then maximizing the result with respect to $\lambda \geq 0$:


$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial x} &= 2(0.2)x - 1 - \lambda \\ &= 0.4x - 1 - \lambda = 0\end{aligned}$$

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -(x - 5) = 0$$

$$\implies x = 5$$

$$\implies \lambda = 0.4(5) - 1 = 2 - 1 = 1$$


As $\lambda \geq 0$, $x = 5$ is a feasible solution to the dual problem.





Optimal separating hyperplane with support vectors for linearly separable data

$$\hat{w}, \hat{b} = \operatorname{argmin}_{w, b} \frac{1}{2} \|w\|^2$$


subject to: $y_i(w^T x_i + b) \geq 1 \quad \forall i$



Real-world datasets are not usually linearly separable



How can the SVM be generalized to handle harder problems?



Key Questions


- I. How can we relax the hard-margin constraints?
- II. Can we gain any insights deriving the dual?
- III. How do we optimize?

Key Questions

I. How can we relax the hard-margin constraints?

II. Can we gain any insights deriving the dual?

III. How do we optimize?




Soft-Margin SVM turns constraint into a cost

Hard Margin:


$$\hat{w}, \hat{b} = \underset{w, b}{\operatorname{argmin}} \frac{1}{2} \|w\|^2 \quad \text{subject to: } y_i(w^T x_i + b) \geq 1 \quad \forall i$$

Soft Margin:

$$\hat{w}, \hat{b} = \underset{w, b}{\operatorname{argmin}} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))$$



Geometry of points near the Soft-SVM decision boundary



$y_i(w^T x_i + b) < 0$: Incorrectly Classified



$0 \leq y_i(w^T x_i + b) < 1$: Weakly Correctly Classified

$1 \leq y_i(w^T x_i + b)$: Strongly Correctly Classified


Building intuition for the margin/boundary violation penalty “Hinge Loss”

$$\hat{w}, \hat{b} = \underset{w, b}{\operatorname{argmin}} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))$$

$$\hat{y} = w^T x_i + b$$



Hyperparameter C sets the penalty on margin violations



$$\hat{w}, \hat{b} = \operatorname{argmin}_{w, b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))$$

$C = 0$:


Drives to uninteresting solution, $\vec{w} = 0$

$C \rightarrow \infty$:


Reduces to Hard-margin SVM

Playing with hyperparameter C


$C = 1.0$



$C = 2.0$



$C = 10.0$



Soft SVM as regularized regression

Ridge Regression:


$$\begin{aligned}\hat{w}, \hat{b} &= \operatorname{argmin}_{w, b} \left(\frac{1}{n} \sum_{i=1}^n (y_i - (w^\top x_i + b))^2 + \lambda \|w\|^2 \right), \text{ where} \\ l_w(x, y) &= (y_i - (w^\top x_i + b))^2 \\ \implies \hat{w}, \hat{b} &= \operatorname{argmin}_{w, b} (E[l(x, y)] + \lambda \|w\|^2)\end{aligned}$$

Soft SVM:

$$\begin{aligned}\hat{w}, \hat{b} &= \operatorname{argmin}_{w, b} \left(C \sum_{i=1}^n \max(0, 1 - y_i(w^\top x_i + b)) + \frac{1}{2} \|w\|^2 \right), \text{ letting} \\ l_w(x, y) &= \max(0, 1 - y_i(w^\top x_i + b)) \\ \implies \hat{w}, \hat{b} &= \operatorname{argmin}_{w, b} \left(E[l(x, y)] + \frac{1}{2nC} \|w\|^2 \right)\end{aligned}$$


The maximum margin classifier has high-variance

$w = [-8.41 \ -3.55], b = 0.27$




seed = 1

$w = [-5.65 \ -6.96], b = -0.1$




seed = 8

$w = [-13.26 \ -10.88], b = 0.86$



seed = 42

Comparing Hinge loss to other loss functions



Loss Function	$L[y, f(x)]$
Binomial Deviance	$\log[1 + e^{-yf(x)}]$
SVM Hinge Loss	$[1 - yf(x)]_+$
Squared Error	$[y - f(x)]^2 = [1 - yf(x)]^2$

Key Questions

I. How can we relax the hard-margin constraints?

II. Can we gain any insights deriving the dual?

III. How do we optimize?

Dual form of Hard SVM problem yielded substantial insights

Soft Margin: $\hat{w}, \hat{b} = \operatorname{argmin}_{w, b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))$


Hard Margin: $\hat{w}, \hat{b} = \operatorname{argmin}_{w, b} \frac{1}{2} \|w\|^2$ subject to: $y_i(w^T x_i + b) \geq 1 \quad \forall i$

$$\mathcal{L}(w, b, \lambda) = \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$$


$\lambda_i > 0 \implies y_i(w^T x_i + b) = 1$
constraint is active, x_i defines the margin

$y_i(w^T x_i + b) > 1 \implies \lambda_i = 0$
constraint is inactive, x_i is far from the margin

$$w = \sum_{i=1}^n \lambda_i y_i x_i$$



Deriving the dual of the Soft SVM problem



Deriving the dual of the Soft SVM problem: Slack variables


For a dataset of n samples $x_i \in \mathbb{R}^d$, with labels $y_i \in \{\pm 1\}$, linearly separated by a hyperplane parametrized by w and b ; $w, x \in \mathbb{R}^d$, and $b \in \mathbb{R}$, the objective of the Soft SVM problem is given by

$$\hat{w}, \hat{b} = \underset{w, b}{\operatorname{argmin}} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i \hat{y}_i)$$

where $\hat{y}_i = w^T x_i + b$

Towards deriving the dual form, we first introduce a constraint for each point in the dataset, using non-negative slack variables, ξ_i :

$$\begin{aligned} \min_{w, b, \xi} \quad & \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i \\ \text{s.t.} \quad & y_i \hat{y}_i \geq 1 - \xi_i \quad ; \xi_i \geq 0 \quad \forall i \end{aligned}$$



Deriving the dual of the Soft SVM problem: The Lagrangian

We next incorporate the constraints of

$$\min_{w,b,\xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i \quad \text{s.t.} \quad \begin{cases} y_i \hat{y}_i \geq 1 - \xi_i & \forall i \\ \xi_i \geq 0 & \forall i \end{cases}$$

Constraint on each predictions relative to corresponding slack variable

$$\hat{y} = 0$$


directly into the objective by introducing a set of Lagrange multipliers for each set of constraints to obtain the Lagrangian

$$L(w, b, \xi, \lambda, \mu) = \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i [y_i \hat{y}_i - (1 - \xi_i)] - \sum_{i=1}^n \mu_i \xi_i$$

Where $\lambda = \{\lambda_i \geq 0\}$ and $\mu = \{\mu_i \geq 0\}$ are sets of Lagrange multipliers. The corresponding objective is then

$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} L(w, b, \xi, \lambda, \mu)$$

Non-negativity constraint on each slack variable



Expanding the Lagrangian

$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i [y_i \hat{y}_i - (1 - \xi_i)] - \sum_{i=1}^n \mu_i \xi_i \right)$$
$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i y_i \hat{y}_i + \sum_{i=1}^n \lambda_i - \sum_{i=1}^n \lambda_i \xi_i - \sum_{i=1}^n \mu_i \xi_i \right)$$

Substituting $\hat{y}_i = w^T x_i + b$

$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i y_i (w^T x_i + b) + \sum_{i=1}^n \lambda_i - \sum_{i=1}^n \lambda_i \xi_i - \sum_{i=1}^n \mu_i \xi_i \right)$$
$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i y_i w^T x_i - \sum_{i=1}^n \lambda_i y_i b + \sum_{i=1}^n \lambda_i - \sum_{i=1}^n \lambda_i \xi_i - \sum_{i=1}^n \mu_i \xi_i \right)$$
$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 - \sum_{i=1}^n \lambda_i y_i w^T x_i - \sum_{i=1}^n \lambda_i y_i b + \sum_{i=1}^n \lambda_i + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i \xi_i - \sum_{i=1}^n \mu_i \xi_i \right)$$

Deriving the dual of the Soft SVM problem: Stationarity

$$\max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 - \sum_{i=1}^n \lambda_i y_i w^T x_i - \sum_{i=1}^n \lambda_i y_i b + \sum_{i=1}^n \lambda_i + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i \xi_i - \sum_{i=1}^n \mu_i \xi_i \right)$$

$$\frac{\partial L}{\partial w} = w - \sum_{i=1}^n \lambda_i y_i x_i = 0$$

$$\implies w = \sum_{i=1}^n \lambda_i y_i x_i$$

$$\frac{\partial L}{\partial b} = - \sum_{i=1}^n \lambda_i y_i = 0$$

$$\implies \sum_{i=1}^n \lambda_i y_i = 0$$

$$\frac{\partial L}{\partial \xi_i} = C - \lambda_i - \mu_i = 0$$

$$\implies \mu_i = C - \lambda_i$$

Deriving the dual of the Soft SVM problem: Stationarity

$$\mathcal{L}_D = \max_{\lambda, \mu \geq 0} \min_{w, b, \xi} \left(\frac{1}{2} \|w\|^2 - \sum_{i=1}^n \lambda_i y_i w^T x_i - \sum_{i=1}^n \lambda_i y_i b + \sum_{i=1}^n \lambda_i + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \lambda_i \xi_i - \sum_{i=1}^n \mu_i \xi_i \right)$$

subject to: $w = \sum_{i=1}^n \lambda_i y_i x_i$, $\sum_{i=1}^n \lambda_i y_i = 0$, and $\mu_i = C - \lambda_i$


$$= \frac{1}{2} \left(\sum_{i=1}^n \lambda_i y_i x_i \right)^T \left(\sum_{j=1}^n \lambda_j y_j x_j \right) - \sum_{i=1}^n \lambda_i y_i \left(\sum_{j=1}^n \lambda_j y_j x_j \right) x_i - \cancel{\sum_{i=1}^n \lambda_i y_i b} + \sum_{i=1}^n \lambda_i + \sum_{i=1}^n \xi_i (C - \lambda_i - \mu_i)$$

$$= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j - \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j + \sum_{i=1}^n \lambda_i + \sum_{i=1}^n \xi_i (C - \lambda_i - (C - \lambda_i))$$

$$= -\frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j + \sum_{i=1}^n \lambda_i$$

$$= \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$$

Wait... why?



Hard-margin SVM:

$$\mathcal{L}_D = \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$$

Soft-margin SVM:

$$\mathcal{L}_D = \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$$



Soft and Hard Margin SVM have different feasible regions

Hard-margin SVM:

$$\begin{aligned}\mathcal{L}_D &= \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j \\ \sum_{i=1}^n \lambda_i y_i &= 0 \\ \lambda_i &\geq 0\end{aligned}$$

Soft-margin SVM:

$$\begin{aligned}\mathcal{L}_D &= \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j \\ \sum_{i=1}^n \lambda_i y_i &= 0 \\ \lambda_i &\geq 0 \\ \mu_i &\geq 0 \\ \mu_i &= C - \lambda_i\end{aligned}$$

$$\implies C - \lambda_i \geq 0$$

$$\implies \lambda_i \leq C$$

$$\implies 0 \leq \lambda_i \leq C$$

Soft and Hard Margin SVM have different feasible regions

Hard-margin SVM:

$$\begin{aligned}\mathcal{L}_D &= \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j \\ \sum_{i=1}^n \lambda_i y_i &= 0 \\ \lambda_i &\geq 0\end{aligned}$$

Soft-margin SVM:

$$\begin{aligned}\mathcal{L}_D &= \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j \\ \sum_{i=1}^n \lambda_i y_i &= 0 \\ 0 &\leq \lambda_i \leq C\end{aligned}$$

Geometry of support vectors in Soft-SVM

Complementary Slackness:

$$\lambda_i(y_i(w^T x_i + b) - 1 + \xi_i) = 0$$

Either:

$$\lambda_i = 0 :$$

$$\implies y_i(w^T x_i + b) - 1 + \xi_i > 0$$

$$\implies y_i(w^T x_i + b) > 1$$


Or:

$$\lambda_i > 0 :$$

$$\implies y_i(w^T x_i + b) - 1 + \xi_i = 0$$

$$\xi = 0 \implies y_i(w^T x_i + b) = 1 \text{ point is on the margin}$$

$$\xi > 0 \implies y_i(w^T x_i + b) < 1 \text{ margin or boundary violation}$$



The dual exposes a relationship between regularization and the size of the weights

$$w = \sum_{i=1}^n \lambda_i y_i x_i$$

$$0 \leq \lambda_i \leq C$$

Soft Margin:

$$\hat{w}, \hat{b} = \operatorname{argmin}_{w, b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))$$

Key Questions

- I. How can we relax the hard-margin constraints?
- II. Can we gain any insights deriving the dual?
- III. How do we optimize?**

Solve for λ_i

$$\sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$$

$$0 \leq \lambda_i \leq C$$

$$\sum_{i=1}^n \lambda_i y_i = 0$$




λ_i


```
import cvxpy as cp
import numpy as np


# Problem data.
m = 30
n = 20
np.random.seed(1)
A = np.random.randn(m, n)
b = np.random.randn(m)

# Construct the problem.
x = cp.Variable(n)
objective = cp.Minimize(cp.sum_squares(A @ x - b))
constraints = [0 <= x, x <= 1]
prob = cp.Problem(objective, constraints)

# The optimal objective value is returned by `prob.solve()`.
result = prob.solve()
# The optimal value for x is stored in `x.value`.
print(x.value)
# The optimal Lagrange multiplier for a constraint is stored in
# `constraint.dual_value`.
print(constraints[0].dual_value)
```





Can you derive the SGD update on the Primal objective?

Now that we're at the end of the lecture, you should be able to...

- ★ Recognize the **hinge loss** for the soft-SVM problem
- ★ Interpret the **geometric properties of the soft-SVM decision boundary**, including **margin, support vectors**, and **slack variables**.
- ★ Describe the effect of **hyperparameter C** on model performance, generalization, and the norm of the weights.
- ★ Perform hyperparameter tuning using techniques like cross-validation to optimize soft-margin SVM parameter.
- ★ List the strengths and limitations of **algorithms for solving soft-margin SVMs**.

	Hard-Margin	Soft-Margin
Dual Objective	$\mathcal{L}_D = \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$	$\mathcal{L}_D = \sum_{i=1}^n \lambda_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j y_i y_j x_i x_j$
Assumptions	Classes are linearly separable	Classes have some overlap
Misclassification	Not allowed	Allowed
Slack variables	N/A	ξ_i : One for each data point, measures margin violation
Dual variables	λ_i : enforces correct classification with some margin	λ_i : margin violation is no more than allowed slack μ_i : slack variables must be non-negative to enforce a penalty
KKT Conditions		
Stationarity	$w = \sum_{i=1}^n \lambda_i y_i x_i$ $\sum_{i=1}^n \lambda_i y_i = 0$	$w = \sum_{i=1}^n \lambda_i y_i x_i$ $\sum_{i=1}^n \lambda_i y_i = 0$ $\mu_i = C - \lambda_i$
Primal feasibility	$y_i(w^T x_i + b) - 1 \geq 0$	$y_i(w^T x_i + b) - 1 + \xi_i \geq 0$ $\xi_i \geq 0$
Dual feasibility	$\lambda_i \geq 0$	$\mu_i \geq 0$ $\lambda_i \geq 0$
Compl. slackness	$\lambda_i (y_i(w^T x_i + b) - 1) = 0$	$\lambda_i (y_i(w^T x_i + b) - 1 + \xi_i) = 0$
Insights	Either: $\lambda_i > 0$ and point is on the margin, or $\lambda_i = 0$ and point is not on margin, far from the decision boundary and is correctly classified in such a way that the constraint does not directly influence the solution. Only points on the margin “support vectors” contribute to defining the separating hyperplane.	$0 \leq \lambda_i \leq C$

