Policy gradient methods

Lecture 1: from gradient ascent to REINFORCE

Carlos Cotrini

ETH Zurich

20 October 2026



Roman copy of a bronze by Lysippos.
Palazzo Altemps, Rome

Aristotle

384 to 322 BC




Roman copy of a bronze by Lysippos.
Palazzo Altemps, Rome

Aristotle

384 to 322 BC

His lectures on the good life, known as the Nicomachean Ethics




Pieter Bruegel the Elder,
The Tower of Babel (detail), 1563.
Kunsthistorisches Museum, Vienna

How a craft is learned

τὰς δ᾽ ἀρετὰς λαμβάνομεν ἐνεργήσαντες πρότερον, ὥσπερ καὶ ἐπὶ τῶν ἄλλων τεχνῶν· ἃ γὰρ δεῖ μαθόντας ποιεῖν, ταῦτα ποιοῦντες μανθάνομεν, οἷον οἰκοδομοῦντες οἰκοδόμοι γίνονται καὶ κιθαρίζοντες κιθαρισταί·

Aristotle, *Nicomachean Ethics*, II.1, 1103a31 to 34.



Pieter Bruegel the Elder,
The Tower of Babel (detail), 1563.
Kunsthistorisches Museum, Vienna

How a craft is learned

τὰς δ' ἀρετὰς λαμβάνομεν ἐνεργήσαντες πρότερον, ὥσπερ καὶ ἐπὶ τῶν ἄλλων τεχνῶν· ἃ γὰρ δεῖ μαθόντας ποιεῖν, ταῦτα ποιοῦντες μανθάνομεν, οἷον οἰκοδομοῦντες οἰκοδόμοι γίνονται καὶ κιθαρίζοντες κιθαρισταί·

The virtues we acquire by first putting them to work, as with the other crafts: for the things we have to learn before we can do them, we learn by doing them, as people become builders by building and lyre players by playing the lyre.

Aristotle, *Nicomachean Ethics*, II.1, 1103a31 to 34.



John and Elizabeth Gould,
Chimney Swallow.
The Birds of Europe, volume 2, 1837

A complete life

Ἔτι δ' ἐν βίῳ τελείῳ. μία γὰρ χελιδὼν ἔαρ οὐ ποιεῖ, οὐδὲ μία ἡμέρα· οὕτω δὲ οὐδὲ μακάριον καὶ εὐδαίμονα μία ἡμέρα οὐδ' ὀλίγος χρόνος.

Aristotle, *Nicomachean Ethics*, I.7, 1098a18 to 20.



John and Elizabeth Gould,
Chimney Swallow.
The Birds of Europe, volume 2, 1837

A complete life

Ἔτι δ' ἐν βίῳ τελείῳ. μία γὰρ χελιδὼν ἔαρ οὐ ποιεῖ, οὐδὲ μία ἡμέρα· οὕτω δὲ οὐδὲ μακάριον καὶ εὐδαίμονα μία ἡμέρα οὐδ' ὀλίγος χρόνος.

And, further, in a complete life. For one swallow does not make a spring, nor does one day; and so one day, or a short time, does not make anyone blessed and happy.

Aristotle, *Nicomachean Ethics*, I.7, 1098a18 to 20.



Raphael, *The School of Athens* (detail), 1509 to 1511.
Apostolic Palace, Vatican City

The definition of virtue

ἔστιν ἄρα ἡ ἀρετὴ ἕξις προαιρετική, ἐν μεσότητι οὖσα τῇ πρὸς ἡμᾶς, ὡρισμένῃ λόγῳ καὶ ᾧ ἂν ὁ φρόνιμος ὁρίσειεν. μεσότης δὲ δύο κακιῶν, τῆς μὲν καθ᾽ ὑπερβολὴν τῆς δὲ κατ᾽ ἔλλειψιν·

Aristotle, *Nicomachean Ethics*, II.6, 1106b36 to 1107a3.



Raphael, *The School of Athens* (detail), 1509 to 1511.
Apostolic Palace, Vatican City

The definition of virtue

ἔστιν ἄρα ἡ ἀρετὴ ἕξις προαιρετική, ἐν μεσότητι οὖσα τῇ πρὸς ἡμᾶς, ὡρισμένῃ λόγῳ καὶ ᾧ ἂν ὁ φρόνιμος ὁρίσειεν. μεσότης δὲ δύο κακιῶν, τῆς μὲν καθ᾽ ὑπερβολὴν τῆς δὲ κατ᾽ ἔλλειψιν·

Virtue, then, is a disposition to choose, lying in a mean relative to us, determined by reason, and as the person of practical wisdom would determine it; a mean between two vices, one of excess and one of deficiency.

Aristotle, *Nicomachean Ethics*, II.6, 1106b36 to 1107a3.

Where the day goes



ταῦτα ποιοῦντες μανθάνομεν

We learn by doing.

Policy gradient methods I

Where the day goes



ταῦτα ποιοῦντες μανθάνομεν

We learn by doing.

Policy gradient methods I



μία γὰρ χελιδὼν ἔαρ οὐ ποιεῖ

One swallow does not make a spring.

Policy gradient methods II

Where the day goes



ταῦτα ποιοῦντες μανθάνομεν
We learn by doing.

Policy gradient methods I



μία γὰρ χελιδὼν ἔαρ οὐ ποιεῖ
One swallow does not make a spring.

Policy gradient methods II



ἐν μεσότητι οὖσα τῇ πρὸς ἡμᾶς
Virtue lies in a mean.

RLHF

Where the day goes

	ταῦτα ποιοῦντες μανθάνομεν *We learn by doing.*	Policy gradient methods I
	μία γὰρ χελιδὼν ἔαρ οὐ ποιεῖ *One swallow does not make a spring.*	Policy gradient methods II
	ἐν μεσότητι οὖσα τῇ πρὸς ἡμᾶς *Virtue lies in a mean.*	RLHF
	ᾧ ἂν ὁ φρόνιμος ὁρίσειεν *As the person of practical wisdom would decide.*	DPO & GRPO

1

Part 1


Gradient ascent

A grasshopper wants to reach the top of the hill



1 The grasshopper stands on the slope.


A grasshopper wants to reach the top of the hill



1 The grasshopper stands on the slope.

2 The top is the point of highest altitude: the place it wants to reach.

Keep jumping uphill until the ground is flat




Idea

```
while not flat:
```

1 At the top the ground is flat: the grasshopper stops there.

Keep jumping uphill until the ground is flat



Idea


```
while not flat:
    jump in the direction
    that increases height
```

1 At the top the ground is flat: the grasshopper stops there.

2 Everywhere else it jumps uphill.


3 To know which way is uphill, it needs the gradient.

The gradient is the change in altitude per step



1 ε (epsilon) is the small step we probe with. Δ (delta) is the altitude change over it.

The gradient is the change in altitude per step




$$\text{Gradient} \approx \frac{\Delta}{\varepsilon} = \frac{\text{change in altitude}}{\text{step size}}$$

Here the step size is the probe step ε. The jump size α comes later.

$\Delta/\varepsilon \rightarrow$

1. ε (epsilon) is the small step we probe with. Δ (delta) is the altitude change over it.
2. The gradient is Δ/ε. We draw it as an arrow under each slope.

The gradient is the change in altitude per step




$$\text{Gradient} \approx \frac{\Delta}{\varepsilon} = \frac{\text{change in altitude}}{\text{step size}}$$

Here the step size is the probe step ε. The jump size α comes later.

1. ε (epsilon) is the small step we probe with. Δ (delta) is the altitude change over it.
2. The gradient is Δ/ε. We draw it as an arrow under each slope.

The gradient is the change in altitude per step




$$\text{Gradient} \approx \frac{\Delta}{\varepsilon} = \frac{\text{change in altitude}}{\text{step size}}$$

Here the step size is the probe step ε. The jump size α comes later.

1. ε (epsilon) is the small step we probe with. Δ (delta) is the altitude change over it.
2. The gradient is Δ/ε. We draw it as an arrow under each slope.
3. Steeper means a longer arrow. When the altitude drops, the arrow points left.


Python measures the gradient with a small step



function

```
def grad(f, x):
```


Python measures the gradient with a small step



function *location*

```
def grad(f, x):
```

Python measures the gradient with a small step




function *location*

```
def grad(f, x):
    eps = 0.01
```

small

1 `eps` is epsilon, the small step we probe with. The picture draws it large.

Python measures the gradient with a small step




function *location*

```
def grad(f, x):
    eps = 0.01          ← small
    delta = f(x + eps) - f(x)
```

1 `eps` is epsilon, the small step we probe with. The picture draws it large.

2 `delta` is the change in altitude over that step: `f(x + eps)` minus `f(x)`.

Python measures the gradient with a small step



function *location*


```
def grad(f, x):
    eps = 0.01          # small
    delta = f(x + eps) - f(x)
    grad = delta / eps
```

1 `eps` is epsilon, the small step we probe with. The picture draws it large.

2 `delta` is the change in altitude over that step: `f(x + eps)` minus `f(x)`.

3 `grad` is `delta / eps`, the slope of the segment between the two points.

Python measures the gradient with a small step



function *location*


```
def grad(f, x):
    eps = 0.01          # small
    delta = f(x + eps) - f(x)
    grad = delta / eps
    return grad
```

1 `eps` is epsilon, the small step we probe with. The picture draws it large.

2 `delta` is the change in altitude over that step: `f(x + eps)` minus `f(x)`.


3 `grad` is `delta / eps`, the slope of the segment between the two points.

Gradient ascent measures, jumps and repeats





```
def gradient_ascent(f, x, alpha=0.1):
```

Gradient ascent measures, jumps and repeats




```
def gradient_ascent(f, x, alpha=0.1):
    g = grad(f, x)
```


Gradient ascent measures, jumps and repeats




```
def gradient_ascent(f, x, alpha=0.1):
    g = grad(f, x)
    while abs(g) > 0.01:
```

1 g is the gradient at x. Flat means g is close to zero, here `abs(g)` at most 0.01.

Gradient ascent measures, jumps and repeats





```
def gradient_ascent(f, x, alpha=0.1):
    g = grad(f, x)
    while abs(g) > 0.01:
        x = x + alpha * g
```

1 `g` is the gradient at `x`. Flat means `g` is close to zero, here `abs(g)` at most 0.01.

2 `alpha` is how far we jump per unit of gradient: the step size of the climb.

3 `eps` probes the ground, `alpha` sets the jump: they are different numbers.

Gradient ascent measures, jumps and repeats





```
def gradient_ascent(f, x, alpha=0.1):
    g = grad(f, x)
    while abs(g) > 0.01:
        x = x + alpha * g
        g = grad(f, x)
```

1 `g` is the gradient at `x`. Flat means `g` is close to zero, here `abs(g)` at most 0.01.

2 `alpha` is how far we jump per unit of gradient: the step size of the climb.

3 `eps` probes the ground, `alpha` sets the jump: they are different numbers.

Gradient ascent measures, jumps and repeats




```
def gradient_ascent(f, x, alpha=0.1):
    g = grad(f, x)
    while abs(g) > 0.01:
        x = x + alpha * g
        g = grad(f, x)
    return x
```

1 `g` is the gradient at `x`. Flat means `g` is close to zero, here `abs(g)` at most 0.01.

2 `alpha` is how far we jump per unit of gradient: the step size of the climb.


3 `eps` probes the ground, `alpha` sets the jump: they are different numbers.

2

Part 2


Mara’s road

A town of fifteen places joined by roads



1 Fifteen places, joined by roads. A day follows one path down to the coast.

A town of fifteen places joined by roads



1 Fifteen places, joined by roads. A day follows one path down to the coast.

2 Play it yourself: scan the code or type the address.

`https://eth-cscs-hs26.github.io/CSCS-PAGE/viz/part2/courier-road/`


Mara plays her fiddle on the way down to the sea



1 Mara is a street fiddler. In one day she walks from Eagle Pass down to the sea, through three places.

Mara plays her fiddle on the way down to the sea






1 Mara is a street fiddler. In one day she walks from Eagle Pass down to the sea, through three places.

2 At every place the road forks: ↙ southwest or ↘ southeast. She picks one and walks on.

Mara plays her fiddle on the way down to the sea








1 Mara is a street fiddler. In one day she walks from Eagle Pass down to the sea, through three places.

2 At every place the road forks: ↙ southwest or ↘ southeast. She picks one and walks on.

3 In every place she plays and the crowd drops coins in her hat. A place pays its usual coins, give or take one coin, whoever passes by. She wants as many coins as possible, but cannot know what a place will give until she has played there.


A simple strategy: play each path several times



average coins of 5 days on each path

1 Luck makes the days differ. Play 5 days on each of the 8 paths and note the average of each.

A simple strategy: play each path several times




average coins of 5 days on each path

1 Luck makes the days differ. Play 5 days on each of the 8 paths and note the average of each.

2 Keep the best average. It finds the best path in 98% of the tries; one day per path only in 69%.

A simple strategy: play each path several times




average coins of 5 days on each path

1 Luck makes the days differ. Play 5 days on each of the 8 paths and note the average of each.

2 Keep the best average. It finds the best path in 98% of the tries; one day per path only in 69%.


3 Trying every path works here, since there are only 8. One more place on a path doubles that to 16, and each further place doubles it again.

Mara follows a table of chances



1 A table of chances: a column for each place where the road forks, a row for each road. Each column adds up to 1, and every cell starts at 0.5.

Mara follows a table of chances



1 A table of chances: a column for each place where the road forks, a row for each road. Each column adds up to 1, and every cell starts at 0.5.

2 At each place s Mara takes road a with the chance in its cell, written $\pi(a \mid s)$. The table is her policy: her rule for choosing roads.

Mara follows a table of chances

↙	0.5	0.5	0.5	0.5	0.5	0.5	0.5
↘	0.5	0.5	0.5	0.5	0.5	0.5	0.5

1 A table of chances: a column for each place where the road forks, a row for each road. Each column adds up to 1, and every cell starts at 0.5.

2 At each place s Mara takes road a with the chance in its cell, written $\pi(a \mid s)$. The table is her policy: her rule for choosing roads.


3 We use a method inspired by gradient ascent because it works well in practice on a wide variety of problems, and it generalises better than the simple strategy.

G_t is the sum of the coins from step t on



1 Gradient ascent needs an altitude. Ours is made of a day's coins, luck included.


G_t is the sum of the coins from step t on



1 Gradient ascent needs an altitude. Ours is made of a day's coins, luck included.

2 G_t is the sum of the coins from step t to the end of the day.


G_t is the sum of the coins from step t on



1 Gradient ascent needs an altitude. Ours is made of a day's coins, luck included.

2 G_t is the sum of the coins from step t to the end of the day.

G_t is the sum of the coins from step t on



1 Gradient ascent needs an altitude. Ours is made of a day's coins, luck included.

2 G_t is the sum of the coins from step t to the end of the day.

3 G_1 is the whole day: here 15 coins.

The same table gives different days

	step 1	step 2	step 3	G_1
day 1	9	4	2	15
day 2	7	7	3	17
day 3	8	5	1	14
day 4	9	3	3	15
day 5	10	3	6	19

1 One row is one day of the same table: its places and their coins.

The same table gives different days

	step 1	step 2	step 3	G_1
day 1	9	4	2	15
day 2	7	7	3	17
day 3	8	5	1	14
day 4	9	3	3	15
day 5	10	3	6	19

1 One row is one day of the same table: its places and their coins.

2 What changes: the roads, decided by the table's chances, and the luck of each place.

The same table gives different days


	step 1	step 2	step 3	G_1
day 1	9	4	2	15
day 2	7	7	3	17
day 3	8	5	1	14
day 4	9	3	3	15
day 5	10	3	6	19

1 One row is one day of the same table: its places and their coins.

2 What changes: the roads, decided by the table's chances, and the luck of each place.


3 So G_1 is random: a different number every day.

A random number cannot be maximised



1 On many days the untouched table gives G_1 anywhere from 11 to 24 coins, some values more often than others.


A random number cannot be maximised



1 On many days the untouched table gives G_1 anywhere from 11 to 24 coins, some values more often than others.

2 The average of G_1 over all the days it can give is 17.25 coins.

A random number cannot be maximised




1 On many days the untouched table gives G_1 anywhere from 11 to 24 coins, some values more often than others.

2 The average of G_1 over all the days it can give is 17.25 coins.


3 A random number cannot be maximised: asking which table gives the biggest G has no answer.

More days bring the average closer to 17.25



1 Each dot is the average of N days in one of 20 runs. The line is the exact average of all possible days.


More days bring the average closer to 17.25



1 Each dot is the average of N days in one of 20 runs. The line is the exact average of all possible days.

2 The dots miss the line by about 0.89 coins at $N = 10$, 0.28 at $N = 100$ and 0.09 at $N = 1{,}000$: the error shrinks as N grows.

More days bring the average closer to 17.25




1 Each dot is the average of N days in one of 20 runs. The line is the exact average of all possible days.

2 The dots miss the line by about 0.89 coins at $N = 10$, 0.28 at $N = 100$ and 0.09 at $N = 1{,}000$: the error shrinks as N grows.

3 The first run of 100 days averages 17.34: our estimate of the exact average from now on.


Our objective is the long-run average $\mathbb{E}[G_1]$



1 Let π_w be the policy of the table w. Play it on N days: the n-th day pays $G_1^{(n)}$.

Our objective is the long-run average $\mathbb{E}[G_1]$

$$\mathbb{E}[G_1] = \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} G_1^{(n)}$$




1 Let π_w be the policy of the table w. Play it on N days: the n-th day pays $G_1^{(n)}$.

2 The average of $G_1^{(1)}, \ldots, G_1^{(N)}$ settles as N grows, at a number we call $\mathbb{E}[G_1]$.

Our objective is the long-run average $\mathbb{E}[G_1]$

$$\mathbb{E}[G_1] = \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} G_1^{(n)}$$

average coins of the first N days



1 Let π_w be the policy of the table w. Play it on N days: the n-th day pays $G_1^{(n)}$.

2 The average of $G_1^{(1)}, \ldots, G_1^{(N)}$ settles as N grows, at a number we call $\mathbb{E}[G_1]$.


3 $\mathcal{J}(w) = \mathbb{E}[G_1]$ is the altitude of the grasshopper, and our objective is to maximise it.

We pick one cell of the table at random

	0.5	0.5	0.5	0.5	0.5	0.5	0.5
	0.5	0.5	0.5	0.5	0.5	0.5	0.5

1 Pick one cell of the table at random.

We pick one cell of the table at random




1 Pick one cell of the table at random.

2 The chosen cell plays the grasshopper's part: its number is the spot where the grasshopper stands.


A copy of the table has ε added to the cell

policy 1: the table




$0.5 + 0.5 = 1$

A copy of the table has ε added to the cell



1 Policy 2 is the table with a small number $\varepsilon = 0.1$ added to the chosen cell.


A copy of the table has ε added to the cell



1 Policy 2 is the table with a small number $\varepsilon = 0.1$ added to the chosen cell.


2 In the copy that column adds up to 1.1. We fix this in a moment.

The method in one picture



1 The table, and a copy with ε added to the chosen cell.


The method in one picture



1 The table, and a copy with ε added to the chosen cell.

2 Both play N days with the same dice. G_1 is the coins of a day; $\mathcal{J}$ and $\mathcal{J}'$ are the averages.

The method in one picture




1 The table, and a copy with ε added to the chosen cell.

2 Both play N days with the same dice. G_1 is the coins of a day; $\mathcal{J}$ and $\mathcal{J}'$ are the averages.


3 The averages give the measured gradient g. The cell jumps by αg and the loop starts again.

Two averages over N days measure the gradient



1 Play $N = 100$ days with the table. Dice decide the roads and the luck of each place. Their average, $\hat{J} = 17.34$, estimates the altitude $J(w)$.


Two averages over N days measure the gradient



1 Play $N = 100$ days with the table. Dice decide the roads and the luck of each place. Their average, $\mathcal{J} = 17.34$, estimates the altitude $\mathcal{J}(w)$.

2 Play the same days again, with the same dice, on the table plus $\varepsilon = 0.1$ in the chosen cell: $\mathcal{J}' = 17.54$.

Two averages over N days measure the gradient



1 Play $N = 100$ days with the table. Dice decide the roads and the luck of each place. Their average, $\mathcal{J} = 17.34$, estimates the altitude $\mathcal{J}(w)$.

2 Play the same days again, with the same dice, on the table plus $\varepsilon = 0.1$ in the chosen cell: $\mathcal{J}' = 17.54$.

3 The same dice give both policies the same luck, so only the change shows. The measured gradient: $g = (\mathcal{J}' - \mathcal{J})/\varepsilon = 2.0$.


The cell jumps by α times the gradient

$$\text{cell} \leftarrow \text{cell} + \alpha\, g$$



1 If the copy earned more, g is positive and the cell goes up; otherwise it goes down.


The cell jumps by α times the gradient



1 If the copy earned more, g is positive and the cell goes up; otherwise it goes down.

2 α is how far we jump per unit of gradient.

The cell jumps by α times the gradient



1 If the copy earned more, g is positive and the cell goes up; otherwise it goes down.

2 α is how far we jump per unit of gradient.

3 ε and α are two different small numbers.

The whole method is a short loop

```
repeat:
  pick a cell of the table at random
```

1 The chosen cell is the grasshopper.

The whole method is a short loop

```
repeat:
  pick a cell of the table at random
  make a copy, add eps to that cell
  play N days with each, same dice
  J, J' = average coins of each
```

1 The chosen cell is the grasshopper.

2 Two policies, and their two altitudes.

The whole method is a short loop


```
repeat:
  pick a cell of the table at random
  make a copy, add eps to that cell
  play N days with each, same dice
  J, J' = average coins of each
  g = (J' - J) / eps
  cell = cell + alpha * g
```

1 The chosen cell is the grasshopper.

2 Two policies, and their two altitudes.


3 The measured gradient, and the jump.

The method in one picture



1 The table, and a copy with ε added to the chosen cell.


The method in one picture



1 The table, and a copy with ε added to the chosen cell.

2 Both play N days with the same dice. G_1 is the coins of a day; $\mathcal{J}$ and $\mathcal{J}'$ are the averages.

The method in one picture




1 The table, and a copy with ε added to the chosen cell.

2 Both play N days with the same dice. G_1 is the coins of a day; J and J' are the averages.

3 The averages give the measured gradient g. The cell jumps by αg and the loop starts again.

Knobs fix the problem: any number is allowed


chances




1 Adding ε to a chance can break a column: it must add up to 1.

Knobs fix the problem: any number is allowed

knobs w




chances



1 Adding ε to a chance can break a column: it must add up to 1.

2 So every cell holds a knob: any number. All knobs start at 0.

Knobs fix the problem: any number is allowed




1 Adding ε to a chance can break a column: it must add up to 1.

2 So every cell holds a knob: any number. All knobs start at 0.


3 Turn the knobs, then convert each column into chances: the softmax.

A bigger knob favours its road



1 One column of the table: a knob for each road at one place. A knob is any number, positive or negative.


A bigger knob favours its road



1 One column of the table: a knob for each road at one place. A knob is any number, positive or negative.


2 A bigger knob favours its road. Here ↙ is favoured. Two steps turn the knobs into chances.

Step 1: take the exponential of each knob



1 The exponential e^x turns any number into a positive one.


Step 1: take the exponential of each knob



1 The exponential e^x turns any number into a positive one.


2 A large knob becomes a large number. A negative knob becomes a number close to 0.

Step 2: divide each by the sum to get chances



1 Add the two numbers, $20.086 + 0.135 = 20.221$, and divide each by this sum.


Step 2: divide each by the sum to get chances



1 Add the two numbers, $20.086 + 0.135 = 20.221$, and divide each by this sum.

2 The chances add up to 1. Knobs 3 and -2 make the ↙ road nearly certain.

Step 2: divide each by the sum to get chances



1 Add the two numbers, $20.086 + 0.135 = 20.221$, and divide each by this sum.


2 The chances add up to 1. Knobs 3 and −2 make the ↙ road nearly certain.

$$\text{chance}_i = \frac{e^{w_i}}{e^{w_1} + e^{w_2}}$$

the softmax of a column of two knobs


Adding ε to a knob always leaves a valid column

before




Adding ε to a knob always leaves a valid column

before




$+\varepsilon$

after $\varepsilon = 0.1$ is added to the first knob



1 The first road becomes a little more likely, and the second a little less.

Adding ε to a knob always leaves a valid column



1 The first road becomes a little more likely, and the second a little less.

2 Whatever the knobs, a column adds up to 1 and each chance stays between 0 and 1.

The same loop turns the knobs

```
w = knobs, all 0
repeat:
   pick a cell of w at random
```

1 The table of knobs is where the grasshopper stands.

2 One pass of the loop is one hop of the grasshopper.

The same loop turns the knobs

```
w = knobs, all 0
repeat:
   pick a cell of w at random
   w2 = w, add eps to that cell
   play N days with softmax(w)
   play N days with softmax(w2)
   (the same dice for both)
   J, J' = average coins of each
   g = (J' - J) / eps
   w[cell] = w[cell] + alpha * g
```

1 The table of knobs is where the grasshopper stands.

2 One pass of the loop is one hop of the grasshopper.

3 The softmax turns every column of knobs into chances.

3

Part 3

Policy gradient methods

The table of all gradients is called $\nabla \mathcal{J}(w)$

knobs w, for example

	1.1	−0.4	0.3	0.7	1.0	−0.2	0.5
	0.2	0.8	−0.9	0.7	−0.3	0.6	−0.6

1 These knobs w give $\mathcal{J}(w) = 19.03$ coins a day on average.

The table of all gradients is called $\nabla \mathcal{J}(w)$

knobs w, for example

↙	1.1	−0.4	0.3	0.7	1.0	−0.2	0.5
↘	0.2	0.8	−0.9	0.7	−0.3	0.6	−0.6

slopes of $\mathcal{J}$

gradient $\nabla \mathcal{J}(w)$

↙	+0.64	−0.23	+0.14	+0.04	+0.28	−0.24	+0.01
↘	−0.64	+0.23	−0.14	−0.04	−0.28	+0.24	−0.01

1 These knobs w give $\mathcal{J}(w) = 19.03$ coins a day on average.

2 Each knob has a gradient, the slope of $\mathcal{J}$ in that knob. The table of all 14 is called $\nabla \mathcal{J}(w)$.

The table of all gradients is called $\nabla J(w)$

knobs w, for example

↙	1.1	−0.4	0.3	0.7	1.0	−0.2	0.5
↘	0.2	0.8	−0.9	0.7	−0.3	0.6	−0.6

slopes of J

gradient $\nabla J(w)$

↙	+0.64	−0.23	+0.14	+0.04	+0.28	−0.24	+0.01
↘	−0.64	+0.23	−0.14	−0.04	−0.28	+0.24	−0.01

1 These knobs w give $J(w) = 19.03$ coins a day on average.

2 Each knob has a gradient, the slope of J in that knob. The table of all 14 is called $\nabla J(w)$.


3 The loop of the last frames measured one entry at a time.

the entry of road a at place s: $\partial J / \partial w_{a,s}$


A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day




A batch of N days has a row for each day



A batch of N days has a row for each day




A batch of N days has a row for each day




1 Each row is one day: a trajectory of states, actions and rewards.

A batch of N days has a row for each day




1 Each row is one day: a trajectory of states, actions and rewards.

A batch of N days has a row for each day




2 A rectangle shows the rewards that G_t adds, from step t to the end.

A batch of N days has a row for each day




2 A rectangle shows the rewards that G_t adds, from step t to the end.

A batch of N days has a row for each day




2 A rectangle shows the rewards that G_t adds, from step t to the end.

A batch of N days has a row for each day




2 A rectangle shows the rewards that G_t adds, from step t to the end.

A batch of N days has a row for each day




2 A rectangle shows the rewards that G_t adds, from step t to the end.

A batch of N days has a row for each day




2 A rectangle shows the rewards that G_t adds, from step t to the end.

A batch of N days has a row for each day




3 The G_1 of each day is its total coins.
Their average estimates $\mathcal{J}$.

A batch of N days has a row for each day



3 The G_1 of each day is its total coins.
Their average estimates $\mathcal{J}$.

A batch of N days has a row for each day



3 The G_1 of each day is its total coins. Their average estimates $\mathcal{J}$.

$\mathcal{J} = \frac{1}{N} \sum_{n=1}^{N} G_1^{(n)} = 17.34$

for $N = 100$ days. $\mathcal{J} \approx \mathcal{J}(w)$: our estimate of the altitude.

A formula estimates the whole table from N days

$$\nabla \mathcal{J}(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{G_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$


1 Play N days with π_w, each of $T = 3$ steps. Step t starts at place s_t and takes road a_t.

A formula estimates the whole table from N days

$$\nabla \mathcal{J}(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{G_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$


1. Play N days with π_w, each of $T = 3$ steps. Step t starts at place s_t and takes road a_t.
2. With N days it is an estimate. It settles as N grows, like the average coins did.

The score says how to turn the knobs




1 At the pass the knobs make the chances 71% and 29%.

The score says how to turn the knobs




1 At the pass the knobs make the chances 71% and 29%.

2 Mara takes ↙ (71%). Its score says how to turn the knobs to make ↙ likelier.




the other 12 knobs of the table get 0

The score says how to turn the knobs



1 At the pass the knobs make the chances 71% and 29%.


2 Mara takes ↙ (71%). Its score says how to turn the knobs to make ↙ likelier.




the other 12 knobs of the table get 0

3 The entry of road b is $\mathbf{1}[b = a_t] - \pi_w(b \mid s_t)$: the knob of the road taken goes up, the other knob goes down by as much.


A rare road has a large score




1 Mara takes ↙ (71%):
a small score




A rare road has a large score




1 Mara takes ↙ (71%): a small score




2 Mara takes ↘ (29%): a large score




A rare road has a large score



1 Mara takes ↙ (71%): a small score



2 Mara takes ↘ (29%): a large score



3 The rarer the road taken, the larger the score: a rare road moves its knob far.

The gradient averages multiplier times score

$$\nabla \mathcal{J}(w) \approx$$

one entry for each road a and place s

1 $\nabla \mathcal{J}(w)$ says how much to adjust the knobs w to increase $\mathcal{J}$.

The gradient averages multiplier times score

$$\underbrace{\nabla \mathcal{J}(w)}_{\text{one entry for each road } a \text{ and place } s} \approx \overbrace{\frac{1}{N} \sum_{n=1}^{N}}^{\text{average over the } N \text{ days}}$$

1 $\nabla \mathcal{J}(w)$ says how much to adjust the knobs w to increase $\mathcal{J}$.

The gradient averages multiplier times score

$$\underbrace{\nabla \mathcal{J}(w)}_{\text{one entry for each road } a \text{ and place } s} \approx \overbrace{\frac{1}{N} \sum_{n=1}^{N}}^{\text{average over the } N \text{ days}} \underbrace{\sum_{t=1}^{T}}_{\text{sum over the steps}}$$

1 $\nabla \mathcal{J}(w)$ says how much to adjust the knobs w to increase $\mathcal{J}$.

The gradient averages multiplier times score

average over the N days

multiplier: the coins from step t on

$$\underbrace{\nabla \mathcal{J}(w)}_{\text{one entry for each road } a \text{ and place } s} \approx \frac{1}{N} \sum_{n=1}^{N} \underbrace{\sum_{t=1}^{T}}_{\text{sum over the steps}} \boxed{G_t^{(n)}}$$

1 $\nabla \mathcal{J}(w)$ says how much to adjust the knobs w to increase $\mathcal{J}$.

The gradient averages multiplier times score

average over the N days

multiplier: the coins from step t on

$$\underbrace{\nabla \mathcal{J}(w)}_{\text{one entry for each road } a \text{ and place } s} \approx \frac{1}{N} \sum_{n=1}^{N} \underbrace{\sum_{t=1}^{T}}_{\text{sum over the steps}} \boxed{G_t^{(n)}} \underbrace{\nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})}_{\text{score of the road } a_t \text{ taken at place } s_t}$$

1 $\nabla \mathcal{J}(w)$ says how much to adjust the knobs w to increase $\mathcal{J}$.

2 It is an average that grows with G_t and with the score of the road taken.

The multiplier G_t can be replaced by Ψ_t

$$\nabla J(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{G_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

The multiplier G_t can be replaced by Ψ_t

$$\nabla \mathcal{J}(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{G_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

replace G_t by Ψ_t

$$\nabla \mathcal{J}(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{\Psi_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

1 It has been shown that the formula still works with another multiplier, written Ψ_t.

The multiplier G_t can be replaced by Ψ_t

$$\nabla \mathcal{J}(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{G_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

↓ replace G_t by Ψ_t

$$\nabla \mathcal{J}(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \boxed{\Psi_t^{(n)}} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

1 It has been shown that the formula still works with another multiplier, written Ψ_t.

2 There are several options for Ψ_t. The first is the one we have: $\Psi_t = G_t$.

4

Part 4

REINFORCE

$\Psi_t = G_t$ gives REINFORCE

$$\underbrace{w \leftarrow w + \alpha}_{1} \quad \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T} \quad \boxed{\Psi_t} \quad \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

1 Gradient ascent again: move the knobs by α times the gradient.

$\Psi_t = G_t$ gives REINFORCE

$$\underbrace{w \leftarrow w + \alpha}_{1} \quad \underbrace{\frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{T}}_{2} \quad \boxed{\Psi_t} \quad \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

1 Gradient ascent again: move the knobs by α times the gradient.

2 The average over N days and $T = 3$ steps is the estimate of the gradient of the last frames.

$\Psi_t = G_t$ gives REINFORCE


$$w \leftarrow w + \alpha \underbrace{\frac{1}{N}\sum_{n=1}^{N}\sum_{t=1}^{T}}_{2} \underbrace{\boxed{G_t^{(n)}}}_{3} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$$

1 Gradient ascent again: move the knobs by α times the gradient.


2 The average over N days and $T = 3$ steps is the estimate of the gradient of the last frames.

3 The multiplier's box holds G_t, the coins from step t to the end of the day: the first option of the last frame.


One term of the gradient for each cell




1 The same batch, grayed. Each G has a term of the gradient below it. We start from G_3 of the first day.




One term of the gradient for each cell



One term of the gradient for each cell




One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell




$\nabla_w \log \pi_w(a \mid s)$ has the entry $\mathbf{1}[b = a] - \pi_w(b \mid s)$ for road b at place s and 0 elsewhere.

One term of the gradient for each cell



2 Every G of every day gets its term: a table of terms, one for each cell of the batch.

One term of the gradient for each cell



3 Sum each row, average the rows: $\nabla J(w) \approx \frac{1}{N} \sum_{n=1}^{N} \sum_{t=1}^{3} G_t^{(n)} \nabla_w \log \pi_w(a_t^{(n)} \mid s_t^{(n)})$

REINFORCE is a short loop

```
w = knobs, all 0
repeat:
  play N days with softmax(w)
```

1 *N* days with the current knobs. No copy of the table and no ε.

REINFORCE is a short loop

```
w = knobs, all 0
repeat:
  play N days with softmax(w)
  push every road of every day
    as hard as the coins after it
```

1 N days with the current knobs. No copy of the table and no ε.

2 Every road taken is pushed as hard as G_t, the coins that came after it.

REINFORCE is a short loop


```
w = knobs, all 0
repeat:
1   play N days with softmax(w)
    push every road of every day
2     as hard as the coins after it
3   w = w + alpha * average push
```

1 N days with the current knobs. No copy of the table and no ε.

2 Every road taken is pushed as hard as G_t, the coins that came after it.

3 The knobs move by α times the average push.


REINFORCE learns from days alone



Median of 200 runs and their middle half. $\alpha = 0.0316$ is the fastest step at which 99 of 100 other runs close 90% of the gap.

1 REINFORCE (10 days an update, $\alpha = 0.0316$) closes 90% of the gap from 17.25 to 21 coins in a median of 1,950 days.

REINFORCE learns from days alone




Median of 200 runs and their middle half. $\alpha = 0.0316$ is the fastest step at which 99 of 100 other runs close 90% of the gap.

1 REINFORCE (10 days an update, $\alpha = 0.0316$) closes 90% of the gap from 17.25 to 21 coins in a median of 1,950 days.

2 Part 2's method ($N = 100$, $\varepsilon = 0.1$, $\alpha = 0.5$) needs 35,800 days: 200 days played for every move of one cell. Tuned for cost ($N = 5$, $\varepsilon = 1$, $\alpha = 1$) it needs about 900.

REINFORCE learns from days alone




Median of 200 runs and their middle half. $\alpha = 0.0316$ is the fastest step at which 99 of 100 other runs close 90% of the gap.

1 REINFORCE (10 days an update, $\alpha = 0.0316$) closes 90% of the gap from 17.25 to 21 coins in a median of 1,950 days.

2 Part 2's method ($N = 100$, $\varepsilon = 0.1$, $\alpha = 0.5$) needs 35,800 days: 200 days played for every move of one cell. Tuned for cost ($N = 5$, $\varepsilon = 1$, $\alpha = 1$) it needs about 900.


3 One update of REINFORCE moves 13 of the 14 knobs, from 10 days. One update of Part 2's method moves one, from 200 days.

A larger step makes runs fail




1 At $\alpha = 0.0316$, 199 of 200 runs close 90% of the gap. At $\alpha = 0.178$, 5.6 times larger, 170 do.

A larger step makes runs fail





1 At $\alpha = 0.0316$, 199 of 200 runs close 90% of the gap. At $\alpha = 0.178$, 5.6 times larger, 170 do.



2 The other 30 runs end on a day that pays 19 coins. The best day pays 21.

A larger step makes runs fail






1 At $\alpha = 0.0316$, 199 of 200 runs close 90% of the gap. At $\alpha = 0.178$, 5.6 times larger, 170 do.

2 The other 30 runs end on a day that pays 19 coins. The best day pays 21.


3 Every road taken is pushed up, so with a large step the first roads taken can run away: the runs that fail settle on their day after about 120 days, like the others.

Every road taken goes up, whatever it pays



1 Days that start with ↙ pay 19.9 coins on average, days that start with ↘ 16.8.


Every road taken goes up, whatever it pays



1 Days that start with ↙ pay 19.9 coins on average, days that start with ↘ 16.8.

2 Each day pushes the road it took by its coins. Coins are never negative, so both roads go up.

Every road taken goes up, whatever it pays



1 Days that start with ↙ pay 19.9 coins on average, days that start with ↘ 16.8.

2 Each day pushes the road it took by its coins. Coins are never negative, so both roads go up.

3 Only the difference, 3.1 coins, tells the better road apart. It is small next to the pushes, and a batch of days can hide it.

A

Appendix


REINFORCE in PyTorch

G is a running sum from the last step back




1 The rewards of the days: a row for each day, a column for each step. We want the coins from each step to the end of the day.

G is a running sum from the last step back




2 Flip every row, so that the last step comes first.

G is a running sum from the last step back



3 Add up along each row: each cell adds one more reward. The first cell is the last reward alone, the second the last two, and so on.


G is a running sum from the last step back



4 Flip back: the cell of step t is G_t, the sum of the rewards from step t on. The 1 means along each row, the steps.


```
G = rewards.flip(1).cumsum(1).flip(1)
```

$\log \pi$ is the log of the chance of each road taken




1 One day: the places visited and the roads taken.

$\log \pi$ is the log of the chance of each road taken




2 Under each road, the chance the table gave to it at that place. At the start every chance is 0.5.

$\log \pi$ is the log of the chance of each road taken



3 Take the log of each chance. As the knobs move, the chances differ from step to step and so do their logs.

$\log \pi$ is the log of the chance of each road taken




step	1	2	3
day 1	−0.69	−0.69	−0.69
day 2	−0.69	−0.69	−0.69
day 3	−0.69	−0.69	−0.69

```
logp = dist.log_prob(roads)
```


4 The three days make the tensor `logp`, a row for each day and a column for each step.

The loss multiplies G and $\log \pi$ cell by cell



1 G and `logp` have the same shape: a row for each day, a column for each step.


The loss multiplies G and $\log \pi$ cell by cell



first cell: $G_1 \times \log \pi_w(a_1 \mid s_1)$:

2 Follow the first cell: the multiplier G_1 and the log chance of the first road.

The loss multiplies G and $\log \pi$ cell by cell



first cell: $G_1 \times \log \pi_w(a_1 \mid s_1)$: $15 \times (-0.69) = -10.4$

3 The product, cell by cell, is a tensor of the same shape: one number for each step of each day.


Add up the steps, average the days, change the sign

day 1	−10.4	−4.2	−1.4
day 2	−11.8	−6.9	−2.1
day 3	−9.7	−4.2	−0.7

`G * logp`


1 The product tensor of the last frame: a number for each step of each day.

Add up the steps, average the days, change the sign




2 Add up each row, the steps of one day: this is the sum over t of $G_t \log \pi_w(a_t \mid s_t)$, a number for each day.

Add up the steps, average the days, change the sign



3 Average over the days: this is the average over n.

Add up the steps, average the days, change the sign



4 Change the sign: this is the loss. Its gradient is minus the average of G_t times the score, so a step downhill on it is a step uphill on $\mathcal{J}$.

```
loss = -(G * logp).sum(dim=1).mean()
```

A library writes the same update in four lines

```
1 loss = -(G * logp).sum(dim=1).mean()
```

1 `G` and `logp` hold G_t and $\log \pi_w(a_t \mid s_t)$: one row for each of the N days, one column for each of the T steps. The loss sums each row, averages the days and flips the sign.

A library writes the same update in four lines

```
loss = -(G * logp).sum(dim=1).mean()
optimiser.zero_grad()
loss.backward()
```

1 `G` and `logp` hold G_t and $\log \pi_w(a_t \mid s_t)$: one row for each of the N days, one column for each of the T steps. The loss sums each row, averages the days and flips the sign.

2 `backward` gives the gradient of the loss: minus the average of G_t times the score.

A library writes the same update in four lines

```
1 loss = -(G * logp).sum(dim=1).mean()
2 optimiser.zero_grad()
  loss.backward()
3 optimiser.step()
```

1 `G` and `logp` hold G_t and $\log \pi_w(a_t \mid s_t)$: one row for each of the N days, one column for each of the T steps. The loss sums each row, averages the days and flips the sign.

2 `backward` gives the gradient of the loss: minus the average of G_t times the score.

3 A step downhill on the loss is a step uphill on $\mathcal{J}$: Part 4's update.