Collision energy revisited

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use energy as well as momentum to classify a collision.

SUCCESS CRITERIA - I can...

1. Calculate total momentum before and after.
2. Calculate total kinetic energy independently.
3. Account for reduced kinetic energy without losing total energy.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Warm-up: bring back an earlier idea

A spring has k=**300 N/m** and compression **0.05 m**. **Find force magnitude** and **describe its direction**.

Marzano check - circle: ① ② ③

02 Worked example: explain the prediction

Reference: $p_{total,\,i} = p_{total,\,f}$; $K = \frac{1}{2}mv^2$.
Negligible external impulse conserves momentum. Only elastic collisions also conserve total kinetic energy.

Worked reasoning: Two **1 kg** carts at **+2** and **0 m/s** stick. Momentum predicts final speed **1 m/s**. Kinetic energy decreases from **2** to **1 J**; momentum can be conserved while kinetic energy decreases.

Your turn: Draw separate momentum arrows and kinetic-energy bars before and after.

Initial **Transfers** **Final**

Draw bars or transfer arrows; label forms and joules.

Marzano check - circle: ① ② ③

03 Read and explain the diagram or equation

Equal-mass carts approach at equal opposite speeds and stick. **What happens** to total momentum and kinetic energy?

Before contact: add velocity arrows.



Cart A Cart B

Marzano check - circle: ① ② ③

04 Apply the physics

Two **1 kg** carts move at **2 m/s** in opposite directions and stick.
1. **Find total initial K** and final K.
2. **Use the energy** diagram to **show the quantities you compare**; **label units** and any unknowns.

Initial **Transfers** **Final**

Draw bars or transfer arrows; label forms and joules.

Marzano check - circle: ① ② ③

Collision energy revisited

antocciphysics

Continue down each column.

05 Explain your reasoning

Why can a collision be inelastic even when the objects bounce apart?

Marzano check - circle: 1 2 3

06 Review: connect an earlier idea

A cart begins with **24 J** kinetic energy and gains **8 J** net work. **Find final kinetic energy**.
Show the energy changes and transfers; **label units**.

Initial **Transfers** **Final**



Draw bars or transfer arrows; label forms and joules.

Marzano check - circle: 1 2 3

07 Exit check

Total K changes from **10** to **7 J** in isolated collision. Is it elastic? Where can **3 J** go? **Use the energy** diagram to **show the quantities you compare**; **label units** and any unknowns.

Initial **Transfers** **Final**



Draw bars or transfer arrows; label forms and joules.

Marzano check - circle: 1 2 3

08 STRETCH: test the prediction

Two **1 kg** carts approach with velocities **+3** and **-3 m/s** and stick on a low-friction track.
1. **Find the common final velocity** and initial and final kinetic energies.
2. Compare with question **4** and **show the energy transferred out** of motion.

Initial **Transfers** **Final**



Draw bars or transfer arrows; label forms and joules.

Marzano check - circle: 1 2 3