Predict a position and a range

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Predict a future position and explain the limits of the prediction.

SUCCESS CRITERIA - I can...

1. Use matching units and include the starting position.
2. Report a central prediction and its bounds.
3. Distinguish a measurement range from model limitations.

Core: teacher selects from 1-9. **Stretch:** 10 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Warm-up: bring back an earlier idea

East is positive. Start at 6 m, go to -4 m, then finish at 3 m. Sketch the route; calculate distance and displacement. Distinguish both from final position.


Marzano check - circle: ① ② ③

02 Worked example: explain the model

Reference: $x_f = x_i + vt$; $\delta x_f = \delta x_i + t\,\delta v$.
Constant velocity; exact positive t and independent bounded starting position/velocity. δ marks a half-width, not a probability.

Worked reasoning: A cart starts at 1 m and moves right at 2 m/s for 3 s. Central prediction: 1+2×3=7 m. If its speed could be 1 to 3 m/s, the endpoint range is 4 to 10 m.

Your turn: A cart moves along a marked track. Show its start, central predicted endpoint, and full possible endpoint range on the number line. What must remain constant throughout the 3 s?



Position (m); right is positive

Marzano check - circle: ① ② ③

Predict a position and a range

antocciphysics

Continue down each column.


03 Units and a starting position

A train starts at x=5 km and moves at +20 m/s for 2 minutes. Calculate displacement and final position in km.

Marzano check - circle: ① ② ③

04 Independent bounds


A motion sensor reports a cart's starting position as 10 ± 1 m and its constant velocity as 2.0 ± 0.2 m/s. Treat the two bounds independently; the elapsed time is exactly 5.0 s. Find the central final position, interval endpoints, half-width and full width. Label the range model.



Marzano check - circle: ① ② ③

05 A negative velocity


A robot starts at 30 ± 2 m and moves with constant velocity -3.0 ± 0.1 m/s. The time is exactly 10 s. Use the smallest and largest allowed inputs to find its full final-position interval.



Marzano check - circle: ① ② ③

06 Show how the bound grows

For x_0=4±1 m and v=2±0.5 m/s, graph central, lower, and upper x(t) from 0 to 4 s. Label the endpoint range.



Marzano check - circle: ① ② ③

Predict a position and a range

antocciphysics

Continue down each column.

07 Input uncertainty or model error?

A precise speed measurement is used to predict an asteroid position with six months of constant velocity. Explain why tiny measurement bounds do not make the forecast exact.

Marzano check - circle: 1 2 3

08 Review: connect an earlier idea

Velocity is +3 m/s for 5 s, then -2 m/s for 3 s. Find displacement and distance. Explain the reference information needed for an angle of 25°.
Sketch the velocity-time model; label the given values.



Marzano check - circle: 1 2 3


09 Exit check

x_0=-5±0.5 m, v=3.0±0.1 m/s, t=10 s exact. Give central final position, its interval, and the full width. Name an assumption.

Marzano check - circle: 1 2 3

10 STRETCH: test the model

Initial position is 2±0.5 m; constant velocity is 3±0.2 m/s; time is exactly 4 s. Find the central prediction and worst-case endpoint bounds.



Marzano check - circle: 1 2 3