What momentum says about impact

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Connect stopping impulse to average force without claiming a complete damage forecast.

SUCCESS CRITERIA - I can...

1. Compute momentum change during a stop.
2. Compare average force for different stopping times.
3. Name information beyond momentum needed for damage.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Warm-up: bring back an earlier idea

Two 3 kg carts at +5 and -3 m/s stick. Find final velocity and use kinetic energy to explain why this is not elastic.

Marzano check - circle: ① ② ③

02 Worked example: explain the model

Reference: $p = mv$; $|F_{avg}| = |\Delta p|/\Delta t$.
Stopping force depends on stopping time as well as momentum. These quantities alone do not specify damage.

Worked reasoning: Two incoming objects have equal momentum 12 kg·m/s but masses 2 and 4 kg, so their speeds are 6 and 3 m/s. Momentum alone does not identify every possible consequence of stopping them.

Your turn: Draw equal momentum arrows and unequal velocity arrows.




2 kg; 6 m/s 4 kg; 3 m/s

Marzano check - circle: ① ② ③

03 Read and interpret the model

Two equal-momentum objects stop in 0.1 s and 1 s. Which has greater average force magnitude, and by what factor?



Marzano check - circle: ① ② ③

04 Apply the physics

A hypothetical object of mass 10^6 kg at 200 m/s stops in 0.50 s. Find impulse and average force magnitude. Then compare a 5 s stop.

Object / trial	Mass	Velocity	Momentum
A	___	___	___
B	___	___	___

Marzano check - circle: ① ② ③

What momentum says about impact

antocciphysics

Continue down each column.


05 Explain your reasoning

Why is a single stopping-force number not a prediction of buildings damaged by an asteroid?

Marzano check - circle: ① ② ③

06 Review: connect an earlier idea

An ideal circular orbit has GM=576 m³/s² and r=16 m. Find speed, inward acceleration and period; explain how velocity changes.
Label velocity and inward acceleration on the circular model.



Marzano check - circle: ① ② ③

07 Exit check

Objects A and B have equal momentum. A has twice B's mass. Compare speeds and kinetic energies, showing why equal momentum need not mean equal energy.

Object / trial	Mass	Velocity	Momentum
A	____	____	____
B	____	____	____

Marzano check - circle: ① ② ③

08 STRETCH: test the model

A 0.5 kg ball arrives at +10 m/s and either stops or rebounds at -6 m/s in 0.04 s. Compare impulse and average force.

Object / trial	Mass	Velocity	Momentum
A	____	____	____
B	____	____	____

Marzano check - circle: ① ② ③