Temperature and particle motion

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use a particle model to distinguish average motion from total energy.

SUCCESS CRITERIA - I can...

1. Describe temperature in terms of microscopic motion.
2. Use kelvin in proportional temperature comparisons.
3. Avoid treating a single particle speed as the temperature.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

LANGUAGE GOAL Use “kelvin” and “particle” to compare your model or result. Sentence starters are optional.

kelvin: the absolute temperature scale used in thermal ratios

particle: a small constituent, such as an atom or molecule

temperature: a measure of thermal state; it is not an energy amount

01 Warm-up: bring back an earlier idea

Sunlight carries energy across 3 million km of nearly empty space in a simplified example.
Identify the transfer mechanism and explain why matter is unnecessary.

Sentence starter: My claim is … . The diagram/data show …, so … .

Marzano check - circle: ① ② ③

02 Worked example: explain the model


Reference: $K_{avg} \propto T$; $T(K) = T(°C) + 273.15$.
Ideal gas: mean translational kinetic energy per particle scales with absolute temperature. ∝ means proportional to.

Worked reasoning: In a gas model, warmer temperature corresponds to greater average random molecular kinetic energy. Individual particles still have a range of speeds.


Your turn: Draw two equal-size samples using dots and arrows; label arrows as particle velocities and make the warmer sample's average arrow length greater.

Sentence starter: Both … . They differ in … because … .

Sample A



Sample B



Add velocity arrows; use lengths to show a range of speeds.

Marzano check - circle: ① ② ③

Temperature and particle motion

antocciphysics

Continue down each column.

Sentence starters are optional. You may plan in any language.


03 Read and interpret the model

Each box represents a sample of the same ideal gas. One is cooler and one warmer. Add particle-velocity arrows with a range of lengths in each box. Make the warmer sample's average kinetic energy greater, without making every particle identical.


Word help: ideal = using stated simplifying conditions

Sentence starter: Both … . They differ in … because … .

Sample A



Sample B



Add velocity arrows; use lengths to show a range of speeds.

Marzano check - circle: 1 2 3


04 Apply the physics

An ideal-gas simulation warms a sample from 300 K to 600 K. By what factor does average translational kinetic energy per particle change?
Label the two particle models to show the comparison. Do not confuse energy with speed.


Word help: ideal = using stated simplifying conditions

Sentence starter: The … in my model represents … . Its label shows … .

Sample A



Sample B



Add velocity arrows; use lengths to show a range of speeds.

Marzano check - circle: 1 2 3

Temperature and particle motion

antocciphysics

Continue down each column.

Sentence starters are optional. You may plan in any language.

05 Explain your reasoning

A student doubles 20°C to 40°C and claims average particle kinetic energy doubles.
Explain why the scale matters.

Sentence starter: My claim is … . The diagram/data show …, so … .

Marzano check - circle: 1 2 3

06 Review: connect an earlier idea

Two students perform the same lift. One takes 4 times as long as the other.
Compare work and average power.
Use the table to distinguish energy from its transfer rate.

Word help: power = rate of energy transfer

Sentence starter: Both … . They differ in … because … .

Case	Energy (J)	Time (s)	Power (W)
A	____	____	____
B	____	____	____


Marzano check - circle: 1 2 3

07 Exit check


At equal kelvin temperature, does a gas sample with twice as many particles have twice the average energy per particle?

Sentence starter: I notice … . This supports … because … .

Sample A



Sample B



Add velocity arrows; use lengths to show a range of speeds.


Marzano check - circle: 1 2 3

08 STRETCH: test the model


Two gas samples have the same temperature but different volumes. Must the particle speeds differ?
Explain what temperature alone fixes in this model.

Sentence starter: Both … . They differ in … because … .

Sample A



Sample B



Add velocity arrows; use lengths to show a range of speeds.

Marzano check - circle: 1 2 3