Specific heat

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use mass, material and temperature change to calculate heating energy.

SUCCESS CRITERIA - I can...

1. Identify ΔT rather than final temperature alone.
2. Use Q=mcΔT with matching units.
3. Compare materials through specific heat.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Warm-up: bring back an earlier idea

A device receives 50 J and delivers 35 J useful mechanical energy. Find other energy and the useful fraction.

Marzano check - circle: ① ② ③

02 Worked example: explain the model

Reference: $Q = mc\,\Delta T$; $\Delta T = T_f - T_i$.
No phase change. A temperature difference of 1°C equals 1 K.

Worked reasoning: A 2 kg sample with c=100 J/(kg·°C) warms by 3°C. Heat absorbed is Q=mc ΔT=600 J, with no phase change or work transfer in this model.

Your turn: Draw an energy arrow into the sample and show which three factors produce 600 J.

Initial T / phase: _____ Final T / phase: _____

Sample

Add an energy arrow; label mass, c or L as needed.

Marzano check - circle: ① ② ③

03 Read and interpret the model

Two equal masses receive the same energy. Material A has a larger specific heat than B. Which has the smaller temperature rise?



A: larger c



B: smaller c

Each outline represents the labeled sample or container.

Marzano check - circle: ① ② ③

04 Apply the physics

In a heating simulation, a 2 kg sample has specific heat c = 100 J/(kg·°C). Its temperature rises from 20°C to 25°C without a phase change. Label the temperatures and energy arrow. Find ΔT, then the energy absorbed.

Initial T / phase: _____ Final T / phase: _____

Sample

Add an energy arrow; label mass, c or L as needed.

Marzano check - circle: ① ② ③

05 Explain your reasoning

Why is multiplying by the final temperature instead of ΔT a mistake?

Marzano check - circle: 1 2 3

06 Review: connect an earlier idea

A pendulum loses 24 J gravitational energy while descending. Neglect losses. Find its kinetic-energy gain. Show the energy changes and transfers; label units.

Initial	Transfers	Final

Draw bars or transfer arrows; label forms and joules.

Marzano check - circle: 1 2 3

07 Exit check

A 1 kg sample with c=200 J/(kg·°C) warms by 3°C. Find Q. Label the energy arrow and identify what changes in the sample.

Initial T / phase: _____ Final T / phase: _____



Add an energy arrow; label mass, c or L as needed.

Marzano check - circle: 1 2 3

08 STRETCH: test the model

Return to the worked example in problem 2. The same 600 J heats 1 kg and 2 kg samples of this material. Compare temperature rises. Label the energy arrow and identify what changes in the sample.

Initial T / phase: _____ Final T / phase: _____



Add an energy arrow; label mass, c or L as needed.

Marzano check - circle: 1 2 3