What changes an asteroid’s path?

antocciphysics | TEACHER KEY | CPA + MATCHING SEI

Learning target: Explain how gravity can change velocity without contact.

Accept equivalent correct methods and clearly labeled models. Language supports do not change the physics demand. Supplied data are examples; actual lab measurements and teacher checkpoints must be recorded from the lesson.

01. Warm-up: bring back an earlier idea *Review: Unit 1, Day 20.*
$a=5$ m/s² right; $v_f=10$ m/s right. Constant net force and mass throughout.

02. Worked example: explain the model
Result points up and right; an upward force does not instantly erase rightward motion.

03. Read and interpret the model
Velocity is tangent; force points toward the central body. Force changes velocity rather than simply pointing along motion.

04. Apply the physics
4 m/s right and 1 m/s down; diagonal down-right.

05. Explain your reasoning
Space near Earth has gravity. Astronauts and spacecraft fall together, so there is little supporting normal force.

06. Review: connect an earlier idea *Review: Unit 1, Day 11.*
7 m/s²; the mass doubles at the same net force. Keep the net force fixed, including resistance.

07. Exit check
A curved path/turn changes direction, therefore velocity.

08. STRETCH: test the model
Longer action produces a larger upward velocity change, assuming the same mass and constant force.

What changes an asteroid’s path?

antocciphysics | TEACHER KEY | HONORS + MATCHING SEI

Learning target: Explain how gravity can change velocity without contact.

Accept equivalent correct methods and clearly labeled models. Language supports do not change the physics demand. Supplied data are examples; actual lab measurements and teacher checkpoints must be recorded from the lesson.

01. Warm-up: bring back an earlier idea *Review: Unit 1, Day 20.*
x interval [-33, -27] km; y interval [45, 55] km. Simplified geometry, gravity, Earth motion or missing observations limit the model. Bounds are not probabilities.

02. Worked example: explain the model
Result points up and right; an upward force does not instantly erase rightward motion.

03. Read and interpret the model
Velocity is tangent; force points toward the central body. Force changes velocity rather than simply pointing along motion.

04. Apply the physics
v=(4,-2) m/s; speed √20=4.47 m/s, increased from 4. Constant downward a is not uniform circular motion.

05. Explain your reasoning
Space near Earth has gravity. Astronauts and spacecraft fall together, so there is little supporting normal force.

06. Review: connect an earlier idea *Review: Unit 1, Day 11.*
Net force 0.35 N; new a≈0.3889 m/s². Extra resistance reduces net force and acceleration if driving force is unchanged.

07. Exit check
Direction changes; speed stays constant because perpendicular force does no work. Velocity is not constant.

08. STRETCH: test the model
Velocity (8, 6) m/s; speed 10 m/s. Original horizontal component remains.