What changes an asteroid’s path?

antocciphysics

UNIT 2 / DAY 1 / HONORS

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Explain how gravity can change velocity without contact.

SUCCESS CRITERIA - I can...

1. Draw velocity tangent to a path and force toward the attracting body.
2. Distinguish motion from a change in motion.
3. State what a straight-line model leaves out.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

A toy model predicts position (-30, 50) km with bounds ±3 km in x and ±5 km in y. Write a short briefing giving both intervals and two reasons it is not a real asteroid collision forecast.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $a = F_{net}/m$.
Gravity attracts masses and can change speed, direction or both. Our asteroid scenario is fictional.


Worked reasoning: A fictional asteroid has a velocity arrow pointing right. A brief upward force gives it an upward change in velocity. The new velocity combines the original rightward motion with this change.

Your turn: Draw the original velocity, the change, and the resulting velocity tip to tail.

Marzano check - circle: ① ② ③

03 Represent and interpret

Sketch the velocity and gravitational-force arrows at the marked point on the orbit. Why do they point in different directions?



The center is the attracting body; the small dot on the circle is the orbiting object.

Marzano check - circle: ① ② ③

04 Solve and justify

For v=(4, 0) m/s and constant a=(0,-1) m/s² over 2 s, find final velocity and speed. Is speed constant in this particular model?

Marzano check - circle: ① ② ③

What changes an asteroid’s path?

antocciphysics

Continue down each column.

05 Analyze the reasoning

A student says space has no gravity because astronauts float. Explain how orbiting astronauts can be falling.

Marzano check - circle: 1 2 3

06 Choose a method

A 0.5 kg system accelerates at 0.7 m/s². Predict acceleration for 0.9 kg at the same net force. Explain the effect of greater resistance.

Marzano check - circle: 1 2 3

07 Exit check

A force stays perpendicular to velocity during ideal circular motion. Explain what changes and what stays constant.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

An object has velocity (8, 0) m/s and constant acceleration (0, 2) m/s² for 3 s. Find the new velocity and speed. Explain why acceleration direction is not the final velocity direction.

Marzano check - circle: 1 2 3