Kinds of collisions

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Separate momentum conservation from kinetic-energy conservation.

SUCCESS CRITERIA - I can...

1. Identify the system and external-impulse assumption.
2. Recognize sticking as perfectly inelastic.
3. Use kinetic energy to distinguish elastic from inelastic.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

A 15 N net force acts right for 2 s on a 3 kg cart initially at rest. Find impulse and final velocity.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $p_{total,\,i} = p_{total,\,f}$; $K = \frac{1}{2}mv^2$.
Negligible external impulse. Elastic collisions also have $K_{total,\,i} = K_{total,\,f}$; sticking transforms some kinetic energy.

Worked reasoning: A 1 kg cart at 4 m/s joins a 1 kg cart at rest. Initial momentum is 4 kg·m/s. Together, 2 kg move at 2 m/s; sticking is one possible collision outcome.

Your turn: Draw before and after snapshots with masses and velocities.

Marzano check - circle: ① ② ③

03 Look and explain

Sketch before/after for a collision where two carts stick. Label one common final velocity, not two different final velocities.

Before contact: add velocity arrows.



Marzano check - circle: ① ② ③

04 Build the solution

A 2 kg cart at 3 m/s hits and sticks to a 1 kg cart at rest. Total initial momentum=6 kg·m/s. Divide by combined mass to find final speed.

Marzano check - circle: ① ② ③

Kinds of collisions

antocciphysics

Continue down each column.

05 Use the idea

A classmate says energy is destroyed in an inelastic collision. Rewrite the claim accurately.

Marzano check - circle: 1 2 3

06 Choose a method

A ball launched horizontally at 7 m/s stays airborne 5 s. Find its landing point measured horizontally from the platform base.

Marzano check - circle: 1 2 3

07 Exit check

Which description guarantees perfectly inelastic: bouncing apart or sticking together? Which quantity still conserves for an isolated system?

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

Return to the worked example in problem 2. Could the same initial carts finish with the first stopped and the second at 4 m/s? Check momentum and explain why momentum alone cannot select between these outcomes.

Marzano check - circle: 1 2 3