Collision energy revisited

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use energy as well as momentum to classify a collision.

SUCCESS CRITERIA - I can...

1. Calculate total momentum before and after.
2. Calculate total kinetic energy independently.
3. Account for reduced kinetic energy without losing total energy.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

LANGUAGE GOAL Use “elastic” and “inelastic” to explain your model or result. Sentence starters are optional.

elastic: a collision that conserves total kinetic energy

inelastic: a collision that transforms some kinetic energy to other forms

kinetic energy: energy associated with motion

01 Bring back an earlier idea

A spring has k=300 N/m and compression 0.05 m.
Find force magnitude and describe its direction.

Word help: magnitude = size of a quantity without its direction

Sentence starter: I found ... [unit]. This value means ... in this situation.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $p_{total,\,i} = p_{total,\,f}$; $K = \frac{1}{2}mv^2$.
Negligible external impulse conserves momentum. Only elastic collisions also conserve total kinetic energy.

Worked reasoning: Two 1 kg carts at +2 and 0 m/s stick. Momentum predicts final speed 1 m/s. Kinetic energy decreases from 2 to 1 J; momentum can be conserved while kinetic energy decreases.

Your turn: Draw separate momentum arrows and kinetic-energy bars before and after.

Sentence starter: The ... in my model represents ... . Its label shows ... .

Marzano check - circle: ① ② ③

03 Look and explain

Equal-mass carts approach at equal opposite speeds and stick. What happens to total momentum and kinetic energy?

Sentence starter: I notice ... . This supports ... because ... .

Before contact: add velocity arrows.



Marzano check - circle: ① ② ③

04 Build the solution

Two 1 kg carts move at 2 m/s in opposite directions and stick.
Find total initial K and final K.

Sentence starter: I found ... [unit]. This value means ... in this situation.

Marzano check - circle: ① ② ③

Collision energy revisited

antocciphysics

Continue down each column.

Sentence starters are optional. You may plan in any language.

05 Use the idea

Why can a collision be inelastic even when the objects bounce apart?

Sentence starter: My claim is … . The diagram/data show …, so … .

Marzano check - circle: 1 2 3

06 Choose a method

A cart begins with 24 J kinetic energy and gains 8 J net work.
Find final kinetic energy.

Word help: final = at the end of the interval; subscript f

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: 1 2 3

07 Exit check

Total K changes from 10 to 7 J in isolated collision. Is it elastic? Where can 3 Jgo?

Sentence starter: I notice … . This supports … because … .

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

Repeat with speeds +2 and -2 m/s.
Find final velocity and initial kinetic energy. Where does the mechanical energy go when the pair sticks?

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: 1 2 3