Sliding, swinging and rolling

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Choose the energy terms that match the type of motion.

SUCCESS CRITERIA - I can...

1. Distinguish translational and rotational kinetic energy.
2. Use equal height changes to compare ideal energy changes.
3. State when a simple point-mass model is insufficient.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

LANGUAGE GOAL Use “translation” and “rotation” to explain your model or result. Sentence starters are optional.

translation: motion of an object’s center from place to place

rotation: turning about an axis

pendulum: an object that swings about a fixed support

01 Bring back an earlier idea

A 4 kg object descends 6 m near Earth.
Use g=10.
Find ΔU and the work done by gravity.

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $K_{translation} = ½mv^2$; $K_{total} = K_{translation} + K_{rotation}$. A rolling object also rotates. Track all energy forms; an ideal pendulum exchanges gravitational and kinetic energy.

Worked reasoning: An object loses 20 J of gravitational potential energy. If 8 J becomes rotational kinetic energy and losses are negligible, only 12 J becomes translational kinetic energy.

Your turn: Draw three energy bars and explain why assigning all 20 J to forward speed would be wrong.

Sentence starter: The step I would change is … . The evidence is … .

Marzano check - circle: ① ② ③

03 Represent and interpret

A slider and a rolling wheel start at the same height from rest. Why can the wheel reach the bottom with a lower center speed?

Sentence starter: Under the assumption …, the evidence … supports … because … .

Marzano check - circle: ① ② ③

04 Solve and justify

A solid cylinder rolls without slipping from height 1.5 m.
Use $I=½mr^2$ and g=10. Derive v from $mgh=½mv^2+½I(v/r)^2$.

Word help: derive = show the reasoning or algebra that produces a relationship

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: ① ② ③

Read down the left column, then down the right. Use the open space to draw, calculate and explain.

Sliding, swinging and rolling

antocciphysics

Continue down each column.

Sentence starters are optional. You may plan in any language.

05 Analyze the reasoning

A pendulum’s tension is perpendicular to its instantaneous motion in the ideal fixed-length model. What work does tension do, and which interaction changes K?

Word help: ideal = using stated simplifying conditions

Sentence starter: I notice … . This supports … because … .

Marzano check - circle: 1 2 3

06 Choose a method

Keep impulse fixed, multiply target mass by 4 and lead time by 6.
Compare the modeled transverse shift using an equation.

Word help: lead time = time available before the modeled event

Sentence starter: Both … . They differ in … because … .

Marzano check - circle: 1 2 3

07 Exit check

A point-mass pendulum drops .80 m vertically from rest.
Find speed at bottom using g=10 and neglect drag.

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

A solid cylinder rolls without slipping from rest through a height of 1.5 m.
Use g=10 and I=½mr². Derive its final speed.

Word help: derive = show the reasoning or algebra that produces a relationship

Sentence starter: I notice … . This supports … because … .

Marzano check - circle: 1 2 3