Specific heat

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use mass, material and temperature change to calculate heating energy.

SUCCESS CRITERIA - I can...

1. Identify ΔT rather than final temperature alone.
2. Use Q=mcΔT with matching units.
3. Compare materials through specific heat.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

LANGUAGE GOAL Use “specific heat” and “heat” to explain your model or result. Sentence starters are optional.

specific heat: energy per mass needed for a one-degree temperature change

heat: energy transferred because of a temperature difference

temperature: a measure of thermal state; it is not an energy amount

01 Bring back an earlier idea

A device receives 50 J and delivers 35 J useful mechanical energy.
Find other energy and the useful fraction.

Sentence starter: I found ... [unit]. This value means ... in this situation.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $Q = mc\Delta T$; $\Delta T = T_f - T_i$.
No phase change. A temperature difference of 1°C equals 1 K.

Worked reasoning: A 2 kg sample with c=100 J/(kg·°C) warms by 3°C. Heat absorbed is Q=mcΔT=600 J, with no phase change or work transfer in this model.

Your turn: Draw an energy arrow into the sample and show which three factors produce 600 J.

Sentence starter: The ... in my model represents ... . Its label shows ... .

Marzano check - circle: ① ② ③

03 Look and explain

Two equal masses receive the same energy. Material A has a larger specific heat than B. Which has the smaller temperature rise?

Sentence starter: Both ... . They differ in ... because ... .



A: larger c



B: smaller c

Each outline represents the labeled sample or container.

Marzano check - circle: ① ② ③

04 Build the solution

A 2 kg sample has c=100 J/(kg·°C). It warms from 20 to 25°C.
Find ΔT, then Q=2×100×ΔT.

Sentence starter: I found ... [unit]. This value means ... in this situation.

Marzano check - circle: ① ② ③

Specific heat

antocciphysics

Continue down each column.

Sentence starters are optional. You may plan in any language.

05 Use the idea

Why is multiplying by the final temperature instead of ΔT a mistake?

Sentence starter: The step I would change is … . The evidence is … .

Marzano check - circle: ① ② ③

06 Choose a method

A pendulum loses 24 J gravitational energy while descending. Neglect losses.
Find its kinetic-energy gain.

Word help: pendulum = an object that swings about a fixed support

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: ① ② ③

07 Exit check

A 1 kg sample with c=200 J/(kg·°C) warms by 3°C.
Find Q.

Sentence starter: I found … [unit]. This value means … in this situation.

Marzano check - circle: ① ② ③

08 STRETCH: Apply it another way

Return to the worked example in problem 2. The same 600 J heats 1 kg and 2 kg samples of this material.
Compare temperature rises.

Sentence starter: Both … . They differ in … because … .

Marzano check - circle: ① ② ③