Calorimetry lab

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use an energy balance to analyze mixing and heat transfer.

SUCCESS CRITERIA - I can...

1. Identify hot and cold samples and their masses.
2. Balance energy gained and lost in an ideal model.
3. Discuss heat exchange with the container and surroundings.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

A 3 kg sample has c=500 J/(kg·°C) and warms by 7°C without changing phase. Find Q.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $Q_{hot} + Q_{cold} = 0$; $Q = mc(T_f - T_i)$.
Ideal isolated mixing, no phase change. Heat lost is negative; heat gained is positive. Real surroundings may exchange energy.

Worked reasoning: Two equal masses of the same liquid at 20°C and 40°C mix in an insulated container. The final 30°C makes the hot sample lose the same energy the cool sample gains.

Your turn: Draw heat-transfer arrows and show the two equal temperature changes.

Marzano check - circle: ① ② ③

03 Look and explain

Equal masses of the same liquid at 20°C and 60°C are mixed in an ideal insulated container. Predict final temperature and justify.



20°C 60°C

Each outline represents the labeled sample or container.

Marzano check - circle: ① ② ③

04 Build the solution

In a supplied trial, the warm sample loses 300 J. If only the cold sample receives that energy, how much does it gain? What if the cup takes 50 J?

Marzano check - circle: ① ② ③

Read down the left column, then down the right. Use the open space to draw, calculate and explain.

05 Use the idea

Plan a table for masses, initial temperatures, final temperature and a heat-loss observation. Why should the final temperature be measured promptly after mixing?

Marzano check - circle: 1 2 3

06 Choose a method

A spring stores 4 J at a given compression. Within its Hooke’s-law range, find the energy at 6 times that compression.

Marzano check - circle: 1 2 3

07 Exit check

An ideal mixture of 30°C and 70°C samples reaches 80°C with no energy added. Is that plausible? Explain.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

Mix one cup of 20°C water with two equal-size cups of 50°C water, neglecting losses. Predict whether final temperature is above or below 35°C, then calculate it.

Marzano check - circle: 1 2 3