Atmospheric entry

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use energy and geometry to discuss entry heating without claiming a guaranteed outcome.

SUCCESS CRITERIA - I can...

1. Compare kinetic energy with a stated heating requirement.
2. Explain the role of surface area relative to mass.
3. Name missing factors in a survival prediction.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

A system receives 50 J heat and does 30 J work on its surroundings. Find ΔU with $\Delta U = Q - W_{by}$.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $K = \frac{1}{2}mv^2$

$Q_{absorbed}$ = fraction absorbed × energy available.

Entry also involves ablation and fragmentation. An energy comparison alone cannot predict survival.

Worked reasoning: A fictional entering object loses 1000 J of mechanical energy; a model assigns 700 J to heated air and 300 J to the object. Those shares are supplied assumptions, not universal atmospheric-entry fractions.

Your turn: Draw the energy budget and identify one measurement needed to test the shares.

Marzano check - circle: ① ② ③

03 Look and explain

Compare a small and a large sphere of the same material. Which has more surface area per unit mass? What does that suggest about surface interactions per kilogram?



A B

Marzano check - circle: ① ② ③

04 Build the solution

A toy model gives 100 J kinetic energy and 20 J needed for a specified heating change. Find the ratio. Does 5 times the required energy prove all 100 J heats the object?

Marzano check - circle: ① ② ③

Atmospheric entry

antocciphysics

05 Use the idea

Name at least three additional quantities needed before claiming a particular rock completely ablates or reaches the ground.

Marzano check - circle: 1 2 3

06 Choose a method

A toy model requires 24 J to escape, but an object has only 20 J and no later energy input. Does the model predict escape? Explain.

Marzano check - circle: 1 2 3

07 Exit check

Two equal-density spheres have radii r and 2 r. Supplied fact: area grows 4 times and mass 8 times. Compare area per mass.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

Compare two objects with the same 1000 J energy loss but different assumed fractions heating the object: 20% and 60%. Find both heating amounts and explain why survival is still uncertain.

Marzano check - circle: 1 2 3