Specific heat

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use mass, material and temperature change to calculate heating energy.

SUCCESS CRITERIA - I can...

1. Identify ΔT rather than final temperature alone.
2. Use Q=mcΔT with matching units.
3. Compare materials through specific heat.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

A 3 kg cart descends 5 m from rest. Use g=10; losses are 6 J. Find bottom speed.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $Q = mc\Delta T$; $\Delta T = T_f - T_i$.
No phase change. A temperature difference of 1°C equals 1 K.

Worked reasoning: A 2 kg sample with c=100 J/(kg·°C) warms by 3°C. Heat absorbed is Q=mcΔT=600 J, with no phase change or work transfer in this model.

Your turn: Draw an energy arrow into the sample and show which three factors produce 600 J.

Marzano check - circle: ① ② ③

03 Represent and interpret

Two equal masses receive the same energy. Material A has a larger specific heat than B. Which has the smaller temperature rise?



A: larger c



B: smaller c

Each outline represents the labeled sample or container.

Marzano check - circle: ① ② ③

04 Solve and justify

A .20 kg metal sample with c=450 J/(kg·°C) absorbs 2700 J. Find its temperature rise and final temperature if it starts at 18°C.

Marzano check - circle: ① ② ③

Specific heat

antocciphysics

Continue down each column.

05 Analyze the reasoning

Why is multiplying by the final temperature instead of ΔT a mistake?

Marzano check - circle: 1 2 3

06 Choose a method

A cart starts from rest at height 4 m. Can it reach 6 m with no external energy input or losses? Justify using kinetic energy.

Marzano check - circle: 1 2 3

07 Exit check

A .50 kg sample needs 4200 J to warm 4°C. Find c and state a necessary assumption.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

A 0.5 kg material absorbs 900 J and warms 6°C. Find c. How much heat warms 1.2 kg of it by 15°C?

Marzano check - circle: 1 2 3