Temperature and particle motion

antocciphysics

UNIT 5 / DAY 6 / HONORS

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Use a particle model to distinguish average motion from total energy.

SUCCESS CRITERIA - I can...

1. Describe temperature in terms of microscopic motion.
2. Use kelvin in proportional temperature comparisons.
3. Avoid treating a single particle speed as the temperature.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

An object at 350 K sits in a vacuum facing surroundings at 230 K. How can it cool, and are absolute-zero surroundings required?

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $K_{avg} \propto T$; $T(K) = T(°C) + 273.15$.
Ideal gas: mean translational kinetic energy per particle scales with absolute temperature. ∝ means proportional to.

Worked reasoning: In a gas model, warmer temperature corresponds to greater average random molecular kinetic energy. Individual particles still have a range of speeds.

Your turn: Draw two equal-size samples using dots and arrows; label arrows as particle velocities and make the warmer sample's average arrow length greater.

Marzano check - circle: ① ② ③

03 Represent and interpret

Sketch particles in samples at lower and higher temperature. Use arrow lengths to represent a range of speeds, with a larger average for the hotter sample.



A B

Each dot is a particle. Each arrow shows that particle's velocity.

Marzano check - circle: ① ② ③

04 Solve and justify

Use average translational energy=(3/2) kT and $k=1.38\times10^{-23}$ J/K. Calculate it at 300 K and compare with 600 K.

Marzano check - circle: ① ② ③

Temperature and particle motion

antocciphysics

Continue down each column.

05 Analyze the reasoning

A student doubles 20°C to 40°C and claims average particle kinetic energy doubles. Explain why the scale matters.

Marzano check - circle: 1 2 3

06 Choose a method

A constant 40 N forward force acts at steady 6 m/s for 8 s. Find mechanical power and work.

Marzano check - circle: 1 2 3

07 Exit check

For a fixed ideal-gas particle mass, average kinetic energy doubles. By what factor does rms speed change?

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

For the same ideal gas, typical molecular speed scales with√T in kelvin. Compare typical speeds at 300 K and 1200 K. Why is a Celsius ratio inappropriate?

Marzano check - circle: 1 2 3