Thermal effects and energy budgets

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Estimate a temperature change only after stating how much energy enters the material.

SUCCESS CRITERIA - I can...

1. Apply an explicit absorbed-energy fraction.
2. Use mass and specific heat in a temperature estimate.
3. State the no-phase-change and no-loss assumptions.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

Bodies initially at 330 K and 350 K exchange heat in an isolated system. Explain what happens to total energy and to the temperature-difference opportunity for work at equilibrium.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $Q_{absorbed} = mc\Delta T$; $\Delta T = Q_{absorbed}/(mc)$.
First identify the absorbed energy. Use only when there is no phase change and c is approximately constant.


Worked reasoning: A surface receives 500 J. A model accounts for 200 J reflected, 250 J absorbed as heating and 50 J transmitted. The full account sums to 500 J.

Your turn: Draw proportional energy arrows and identify which part can directly raise this surface's temperature.

Marzano check - circle: ① ② ③

03 Represent and interpret

A model releases 100 J. Only 20% enters a sample. Draw the energy partition and calculate energy absorbed.



Bar height represents energy. Read the value and unit above each bar.

Marzano check - circle: ① ② ③

04 Solve and justify

A hypothetical energy release is 2×10^6 J. A .50 kg sample absorbs 1%; c=800 J/(kg·°C). EstimateΔT without phase change or losses.

Marzano check - circle: ① ② ③

Thermal effects and energy budgets

antocciphysics

Continue down each column.

05 Analyze the reasoning

A temperature prediction exceeds a material’s melting point, but the calculation used only mcΔT. What must be added or reconsidered?

Marzano check - circle: 1 2 3

06 Choose a method

A 4 kg cart descends 6 m from rest. Use g=10; losses are 8 J. Find bottom speed.

Marzano check - circle: 1 2 3

07 Exit check

A 2 kg sample with c=500 warms 8°C. Find absorbed energy. If this is 4% of source energy, find source energy.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

A 0.4 kg panel absorbs 1200 J, emits 300 J and does no work. With c=450 J/(kg·K), find internal-energy and temperature changes.

Marzano check - circle: 1 2 3