Intensity, decibels and Doppler

antocciphysics

Name: _____ **Block:** _____ (teacher only) **Mastery Rating:** _____

LEARNING TARGET Distinguish intensity change from frequency shift.

SUCCESS CRITERIA - I can...

1. Use an intensity ratio separately from pitch.
2. Interpret a logarithmic sound-level difference.
3. Predict approach and recession Doppler trends.

Core: teacher selects from 1-7. **Stretch:** 8 after checking your work.

MARZANO SELF-CHECK

① Need help starting

② Can do part; need help

③ Solve and explain with allowed supports

01 Bring back an earlier idea

A 500 Hz tone lasts 0.3 s. Find number of cycles and period.

Marzano check - circle: ① ② ③

02 Explain a worked example

Reference: $\Delta\beta = 10 \log_{10}(I_2/I_1)$.
I is intensity; β is sound level in dB. Tenfold intensity gives +10 dB. Approach generally raises observed frequency; recession lowers it.

Worked reasoning: For the same point source, doubling distance spreads its power over four times the area, so ideal intensity drops to one quarter. This assumes spherical spreading without extra absorption.

Your turn: Draw two concentric wavefronts and indicate why the outer surface shares the power more widely.

Marzano check - circle: ① ② ③

03 Represent and interpret

A moving source approaches a stationary listener, then recedes. Sketch closer wavefront spacing in front and wider spacing behind; compare heard pitch.

Source moves right. Add wavefronts.



Marzano check - circle: ① ② ③

04 Solve and justify

Intensity increases by factor 4. FindΔβ. Then compare intensities for sound levels 60 and 80 dB.

Marzano check - circle: ① ② ③

Intensity, decibels and Doppler

antocciphysics

Continue down each column.

05 Analyze the reasoning

A loud stationary siren and a quiet approaching siren are compared. Does “louder” alone mean “higher frequency”? Explain.

Marzano check - circle: 1 2 3

06 Choose a method

A 4 kg sample with c=500 J/(kg·°C) warms 6°C. Find absorbed energy. If it is 5% of source energy, find source energy.

Marzano check - circle: 1 2 3

07 Exit check

Stationary listener; source approaches at 20 m/s; emits 500 Hz; v_{sound}=340 m/s. Use $f_{obs}=f_s\ v/(v-v_s)$. Find observed frequency.

Marzano check - circle: 1 2 3

08 STRETCH: Apply it another way

Using $\Delta\beta=10\log_{10}(I_2/I_1)$, find the level change when distance triples from an ideal point source. Separately explain the higher observed pitch of an approaching source.

Marzano check - circle: 1 2 3