

# Fundamentals of Linear Algebra and Optimization

## Jean Gallier

### Project 1

September 17, 2015; Due October 6, 2015

Group Members:

Ethan Brooks, Abhishek Ravichandran, Brian Wright

#### Problem P1.

(1) Refer to the MATLAB program `Part1.m`.

(2) Except for the first and last control points, pairs of de Boor control points,  $b_0^{i+1}$  and  $b_3^i$ , have common equations in terms of  $d_{i+1}$ , and often  $d_i$  and/or  $d_{i+2}$ . By taking the coefficients of each  $d$  term and placing them in a matrix,  $A$ , the equation  $Ad = x$  can then be written in a manner which satisfies all of the equations in terms of  $d$ .

$$\begin{aligned}6b_0^2 &= \frac{3}{2}d_1 + \frac{7}{2}d_2 + (1)d_3 \\6b_0^i &= (1)d_{i-1} + (4)d_i + (1)d_{i+1}; (3 \leq i < N-3) \\6b_0^{N-2} &= (1)d_{N-3} + \frac{7}{2}d_{N-2} + \frac{3}{2}d_{N-1}\end{aligned}$$

Since the range of superscript values on the left side of the above equations is 2 to  $N-2$ , and the subscripts of  $d$  have a range 1 to  $N-1$ , a vector can be made for  $d$  which ranges 2 to  $N-2$  and the remaining values of  $d$  (since they each only occur in a single equation) can be included in the  $x$  vector.

#### Problem P2.

Refer to the MATLAB program `Part2_1.m` and `Part2_2.m`.