



Contents lists available at ScienceDirect

## Journal of Cancer Research and Practice

journal homepage: <http://www.journals.elsevier.com/journal-of-cancer-research-and-practice>

## Review Article

## Review of cancer from perspective of molecular



Seyed Hossein Hassanpour\*, Mohammadamin Dehghani

Department of Toxicology, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

## ARTICLE INFO

## Article history:

Received 4 March 2017

Received in revised form

1 July 2017

Accepted 10 July 2017

Available online 10 July 2017

## Keywords:

Cancer

Gene mutations

Carcinogenesis

## ABSTRACT

Cancer in the broader sense refers to more than 277 different types of cancer disease. Scientists have identified different stage of cancers, indicating that several gene mutations are involved in cancer pathogenesis. These gene mutations lead to abnormal cell proliferation. Genetic disorders caused by heritance or inheritance factors have a pivotal role in the increase of cell growth. With the assistance of technological advances in bioinformatics and molecular techniques, additional information has been obtained that can be useful for early diagnosis and proper treatment. The effects of drugs on patients with cancer can predict and even manage some aspects of side effects. In recent years, carcinogenesis mechanisms have been detected by molecular genetic studies. The results of these studies led to an improved understanding of the role of genetic disorders in cancer formation. In this study, our aim was to review molecular aspects of cancer.

© 2017 Taiwan Oncology Society. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Cancer is the second leading cause of mortality worldwide. Overall, the prevalence of cancer has actually increased; just in the United States alone, approximately 1,665,540 people suffered from cancer, and 585,720 of them died due to this disease by 2014<sup>1</sup>. Therefore, cancer is a serious problem affecting the health of all human societies. Unfortunately, it is a variety disease at the tissue level and this variety is a major challenge for its specific diagnosis, followed by efficacy of treatment<sup>2,3</sup>. In men, the highest percentages of cancer types occur in the prostate, lung and bronchus, colon and rectum, and urinary bladder, respectively. In women, cancer prevalence is highest in the breast, lung and bronchus, colon and rectum, uterine corpus and thyroid, respectively. This data indicates that prostate and breast cancer constitute a major portion of cancer in men and women, respectively<sup>4</sup>. For children, the highest percentage types of cancer disease are blood cancer, and cancers related to the brain and lymph nodes, respectively<sup>5,6</sup>. Cancer occurs by a series of successive mutations in genes so that these mutations change cell functions. Chemical compounds have an obvious role of forming gene mutations and cancer cells. In addition, smoking involves several carcinogenic chemical compounds that lead to lung

cancer<sup>7</sup>. Interestingly, environmental chemical substances with carcinogenic properties influence directly or indirectly the cytoplasm and nucleus of cells, and lead to genetic disorders and gene mutations<sup>8–11</sup>. Viruses, bacteria and radiation rays are other carcinogenesis factors, comprising about 7% of all cancers<sup>12</sup>. In general, cancer disrupts cellular relations and results in the dysfunction of vital genes. This disturbance is affective in the cell cycle, and leads to abnormal proliferation<sup>13,14</sup>. Proto-oncogenes are responsible for cell division and growth under normal condition, but become oncogenes during genetic mutation, which are most dangerous for cell existence<sup>15</sup>. In addition, the lack of tumor suppressor genes triggers uncontrolled cells division<sup>16</sup>. Normally, repair genes translate to protein and enzymes that have repairing properties and more than 30 types of detected repair proteins<sup>17</sup>. Removing uracil from DNA bypasses the DNA damage and removes the main DNA lesions induced by ultraviolet light, which are essentially the functions of repair genes to successfully repair DNA<sup>18</sup>.

Epigenetics is a dynamic situation during the study of cell fate and epigenetic modifications such as DNA methylation, histone modifications and nucleosome position, which play important roles in cancer formation<sup>19,20</sup>. Cancer cells are characterized by a vast reduction in DNA methylation (about 5–6% reduction in the total amount of 5-methyl cytosine)<sup>21</sup>. Overall reduction of mono-acetylated H4K16 forms the majority of histone modifications in cancer cells<sup>22</sup>. All families of chromatin modifying proteins are associated with cancer, although in most cases, the molecular

\* Corresponding author. Department of Toxicology, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

E-mail address: [Dr.hossein1366@yahoo.com](mailto:Dr.hossein1366@yahoo.com) (S.H. Hassanpour).

Peer review under responsibility of Taiwan Oncology Society.

mechanisms underlying their functions remain unknown<sup>23</sup>. In this study, we reviewed cancer from the perspective of the molecular level, in order to get a closer look at this disease.

## 2. Review method

Initially, we searched research papers using keywords such as cancer and molecular process, cancer and treatment and molecular aspects. Subsequently, the papers that matched such word criteria were fully reviewed and their findings duly noted.

## 3. Cancer from the molecular prospective

Genetic changes that lead to oncogene generation and genetic disorders include chromosomal translocation (gene Bcr and oncogene Abl in chronic blood cancer), point mutation (Ras gene in colon cancer), deletion (Erb-B gene in breast cancer), amplification (N-myc in neuroblastoma), and insertion activation (C-myc in acute blood cancer). Chronic blood cancer often occurs in the elderly due to an exchange of genetic material between chromosomes 9 and 22. This condition leads to production of a biomarker called ph1, which is found in 95% of patients and can facilitate a correct diagnosis. The connection of Bcr gene to Abl oncogene results in creation of a gene new combination that translated to protein with kinase activity<sup>24–27</sup>.

Mutation in the p53 gene leads to formation of an unusual protein that has a prominent role in disturbance of molecular process related to p53. Abnormality of these molecular and biological events leads to formation of cancer cells; therefore, the p53 gene has a complex relationship with cancer and it has been reported that p53 abnormality occurs in 60% of cancer cases. Under normal conditions, p53 plays an important role in cell division, cell death, senescence, angiogenesis, differentiation, and DNA metabolism. Additionally, the majority of mutations related to the p53 gene occur in the DNA-binding position, and the disability of genes is controlled by p53 for replication. Cooperation of p53 with CDK1-P2 and CDC2 keeps cancer cells in G1 and G2 phases of cell cycle<sup>28,29</sup>. In fact, p53 is either an inhibitor or a promoter of cancer cells. After DNA damage has been caused by other genes, the p53 protein binds to DNA and leads to stimulation of the WAF1 gene<sup>30,31</sup>. This event leads to connection of p53 with CDK2, and ultimately inhibits the effect of p21 for the next stage of the cell cycle. The anti-cancer action of p53 is active during three routes stimulating, for DNA repairing proteins, induction of apoptosis, and arresting of cell cycle in G1/S phase<sup>16,32–38</sup>.

Overall, hypomethylation primarily occurs in repeated sequences and leads to an increase of movement, and deletion of genes as well as chromosomal instability<sup>21,39</sup>. L1 from the LINE family is a prime example of hypomethylation, which has been observed in a range of cancers including breast, lung, and bladder cancer<sup>40</sup>. Hypomethylation in specific promoters can activate the ectopic expression of oncogenes; for example, this condition occurs for MASPIN as a tumor suppressor gene in breast and prostate cancer. Other examples include S100P in pancreatic cancer, SNCG in breast and ovarian cancer, MAGE and DPP6 in melanoma<sup>41</sup>. Unlike overall hypomethylation, hypermethylation occurs only in a specific CpG region. Transcription inactivation caused by hypermethylation of promoter influences genes involved in repair (Hmlh1, WRN and BBRCA1), response to vitamin (CRBP1, RARB2), cell cycle control (P16INK4b, P16INK4a) and apoptosis (TMS1, DAPK1, and WIF-1). These events have a pivotal role in the induction of cancer<sup>42</sup>. Therefore, hypermethylated promoters can be considered as novel biomarkers for the diagnosis and prognosis of cancer because most studies are focused on the CpG regions of promoters. In addition, it is an important option that frequently

aberrant methylation such as hypermethylation in CpG region occurs during cancer (45–65%)<sup>43,44</sup>. Overall disturbance in DNA methylation patterns can be due to impaired regulation of DNMT, such that it has been determined that DNMT1 and DNMT3b have a high level of expression in many tumors. In addition, miRNA regulates DNMT expression, and it has been found that the MIR-29 family reduces expression of DNMT3a, DNMT3b (directly) and DNMT1 (indirectly)<sup>45</sup>.

Deacetylation occurs by HDAC so that this condition is involved in different tumors. The Sirtuin family is a main class of HDAC enzymes<sup>46</sup>. The increase of SirT1 expression and activity has been shown to have various types. In addition, cooperation between SirT1 and DNMT1 affects DNA methylation<sup>47</sup>. HDAC expression can be regulated by microRNA; for example, miR-449a regulates cell growth and survival during prostate cancer through suppression of HDAC-1 expression. In addition to alteration of HDAC expression in many cancers such as colon cancer, lung and leukemia, disturbance in histone acetylation occurs through ectopic mutations and deletion in HAT and its related genes. These events can be a main reason for cancer formation<sup>48</sup>. Moreover, cancer cells lost H4K16ac, H3K4me3, H4K20me3 and H3K27me3<sup>49</sup>. Change in the distribution of histone methylation is mainly due to expression of histone methyltransferases and histone demethylase. In addition, disabling mutations in SETD2 (a histone methyltransferase) and UTX (a histone demethylase) occurs during renal carcinoma<sup>50</sup>. In leukemia, MLL oncoprotein leads to abnormal patterns of H3K4 and H3K29 methylation, and ultimately changes in the expression of the target genes of MLL<sup>51</sup>.

BRG1 and BRM (as subunits of ATPase related to SWI/SNF complex) are known as tumor suppressors that manifest a pivotal role in 15–20% of lung cancer. Interestingly, it has been reported that BRG1 is a destabilizing of p53 and SWI/SNF complex, and regulates gene expression as local. The SWI/SNF complex is involved in formation of many cancers through its reaction with RB, p53, MYC, MLL, and BRCA1. Therefore, disabling of the SWI/SNF complex disrupts cell growth. In addition, changes in the position of nucleosomes leads to suppression of transcription by promoter hypermethylation. Promoter hypermethylation causes TSS occupation by nucleosomes, and this issue has been reported involving MLH1 in colon cancer. Encoding genes related to subunits of changing complexes of chromatin such as CHD5 are the main target of CpG hypermethylation during cancer. This condition leads to a reduction of its expression and a disturbance in the normal structure of chromatin<sup>52</sup>. In addition to the situation associated with nucleosomes position, histone variants are also associated with cancer such as an increase of MacroH2A expression during the senescence process. Thus, lung tumors along with an increased MacroH2A expression have a better prognosis due to reduction of cell proliferation amount<sup>53</sup>.

## 4. Conclusion

In the past three decades, researchers have reported a substantial volume of information about genes and proteins and their roles in the production of cancer cells. In fact, the role of mutated genes in cancer cells was one of the most important discoveries. Recently, environmental factors related to genetic mutations have been identified. With the help of different molecular methods, we are able to determine the potency of gene expression and defective proteins, as well as detecting novel cancer biomarkers. These findings can be useful to treat cancer and reduce cancer complications. In addition, various studies to explore the epigenetic mechanisms and their relationship together with the development, and progression of various diseases, especially cancer are continuing. Besides, it seems that many aspects of epigenetic

remain unknown. However, by identifying all environmental factors and pivotal genes, this gives us a comprehensive map for further efforts to reduce cancer in the future.

### Conflict of interest

The authors declare that there is no conflict of interest regarding this paper.

### References

1. Siegel R, Naishadham D, Jemal A. *Cancer statistics 2013*, CA Cancer J. Clin. 2013;63:11–30.
2. Meacham CE, Morrison SJ. Tumour heterogeneity and cancer cell plasticity. *Nature*. 2013;501:328–337.
3. Fisher R, Pusztai L, Swanton C. Cancer heterogeneity: implications for targeted therapeutics. *Br J Cancer*. 2013;108:479–485.
4. Siegel RL, Miller KD, Jemal A. *Cancer statistics. 2016*, CA Cancer J Clin. 2016;66:7–30.
5. Schottenfeld D, Fraumeni Jr JF. *Cancer Epidemiology and Prevention*. Oxford University Press; 2006.
6. Yoo KY, Shin HR. Cancer epidemiology and prevention. *Korean J Epidemiol*. 2003;25:1–15.
7. Aizawa K, Liu C, Tang S, et al. Tobacco carcinogen induces both lung cancer and non-alcoholic steatohepatitis and hepatocellular carcinomas in ferrets which can be attenuated by lycopene supplementation. *Int J Cancer*. 2016;139:1171–1181.
8. Poon SL, McPherson JR, Tan P, Teh BT, Rozen SG. Mutation signatures of carcinogen exposure: genome-wide detection and new opportunities for cancer prevention. *Genome Med*. 2014;6:24.
9. Trafalek J, Kolanowski W. Dietary exposure to meat-related carcinogenic substances: is there a way to estimate the risk? *Int J Food Sci Nutr*. 2014;65:774–780.
10. Cumberbatch MG, Cox A, Teare D, Catto JW. Contemporary occupational carcinogen exposure and bladder cancer: a systematic review and meta-analysis. *JAMA Oncol*. 2015;1:1282–1290.
11. Antwi SO, Eckert EC, Sabaque CV, et al. Exposure to environmental chemicals and heavy metals, and risk of pancreatic cancer. *Cancer Causes Control*. 2015;26:1583–1591.
12. Parkin DM. The global health burden of infection-associated cancers in the year 2002. *Int J Cancer*. 2006;118:3030–3044.
13. Seto M, Honma K, Nakagawa M. Diversity of genome profiles in malignant lymphoma. *Cancer Sci*. 2010;101:573–578.
14. Cigudosa JC, Parsa NZ, Louie DC, et al. Cytogenetic analysis of 363 consecutively ascertained diffuse large B-cell lymphomas. *Genes, Chromosomes Cancer*. 1999;25:123–133.
15. Shtivelman E, Lifshitz B, Gale RP, Canaani E. Fused transcript of abl and bcr genes in chronic myelogenous leukaemia. *Nature*. 1985;315:550–554.
16. Matlashewski G, Lamb P, Pim D, Peacock J, Crawford L, Benchimol S. Isolation and characterization of a human p53 cDNA clone: expression of the human p53 gene. *EMBO J*. 1984;3:3257.
17. Wei Q, Li L, Chen D. *DNA Repair, Genetic Instability, and Cancer*: World Scientific. 2007.
18. Wood RD, Mitchell M, Sgouros J, Lindahl T. Human DNA repair genes. *Science*. 2001;291:1284–1289.
19. Alvarez-Buylla ER, Chaos Á, Aldana M, et al. Floral morphogenesis: stochastic explorations of a gene network epigenetic landscape. *PLoS One*. 2008;3:e3626.
20. Portela A, Esteller M. Epigenetic modifications and human disease. *Nat Biotechnol*. 2010;28:1057–1068.
21. Goelz SE, Vogelstein B, Hamilton SR, Feinberg AP. Hypomethylation of DNA from benign and malignant human colon neoplasms. *Science*. 1985;228:187–191.
22. Fraga MF, Ballestar E, Villar-Garea A, et al. Loss of acetylation at Lys16 and trimethylation at Lys20 of histone H4 is a common hallmark of human cancer. *Nat Genet*. 2005;37:391–400.
23. Sharma S, Kelly TK, Jones PA. Epigenetics in cancer. *Carcinogenesis*. 2010;31:27–36.
24. Joensuu H, Dimitrijevic S. Tyrosine kinase inhibitor imatinib (STIS71) as an anticancer agent for solid tumours. *Ann Med*. 2001;33:451–455.
25. King CR, Kraus MH, Aaronson SA. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science*. 1985;229:974–977.
26. Heinrich MC, Blanke CD, Druker BJ, Corless CL. Inhibition of KIT tyrosine kinase activity: a novel molecular approach to the treatment of KIT-positive malignancies. *J Clin Oncol*. 2002;20:1692–1703.
27. Thomas RK, Baker AC, DeBiasi RM, et al. High-throughput oncogene mutation profiling in human cancer. *Nat Genet*. 2007;39:347–351.
28. Chae SW, Sohn JH, Kim D-H, et al. Overexpressions of Cyclin B1, cdc2, p16 and p53 in human breast cancer: the clinicopathologic correlations and prognostic implications. *Yonsei Med J*. 2011;52:445–453.
29. Taylor WR, Stark GR. Regulation of the G2/M transition by p53. *Oncogene*. 2001;20:1803.
30. Bukholm IK, Nesland JM. Protein expression of p53, p21 (WAF1/CIP1), bcl-2, Bax, cyclin D1 and pRb in human colon carcinomas. *Virchows Arch*. 2000;436:224–228.
31. Roninson IB. Oncogenic functions of tumour suppressor p21 Waf1/Cip1/Sdi1: association with cell senescence and tumour-promoting activities of stromal fibroblasts. *Cancer Lett*. 2002;179:1–14.
32. May P, May E. Twenty years of p53 research: structural and functional aspects of the p53 protein. *Oncogene*. 1999;18:7621.
33. McBride O, Merry D, Givol D. The gene for human p53 cellular tumor antigen is located on chromosome 17 short arm (17p13). *Proc Natl Acad Sci*. 1986;83:130–134.
34. Isobe M, Emanuel B, Givol D, Oren M, Croce CM. Localization of gene for human p53 tumour antigen to band 17p13. *Nature*. 1986;320:84–85.
35. Miller C, Koeffler H. P53 mutations in human cancer. *Leukemia*. 1993;7:S18–S21.
36. Koshland Jr DE. Molecule of the year. *Science*. 1991;254:1705–1706.
37. Muller PA, Voudsen KH. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell*. 2014;25:304–317.
38. Selivanova G. p53: fighting cancer. *Curr Cancer Drug Targets*. 2004;4:385–402.
39. Kanai Y, Hirohashi S. Alterations of DNA methylation associated with abnormalities of DNA methyltransferases in human cancers during transition from a precancerous to a malignant state. *Carcinogenesis*. 2007;28:2434–2442.
40. Wilson AS, Power BE, Molloy PL. DNA hypomethylation and human diseases. *Biochim Biophys Acta*. 2007;1775:138–162.
41. Futscher BW, O Meara MM, Kim CJ, et al. Aberrant methylation of the maspin promoter is an early event in human breast cancer. *Neoplasia*. 2004;6:380–389.
42. Esteller M. Epigenetic gene silencing in cancer: the DNA hypermethylome. *Hum Mol Genet*. 2007;16:R50–R59.
43. Li M, Chen W-D, Papadopoulos N, et al. Sensitive digital quantification of DNA methylation in clinical samples. *Nat Biotechnol*. 2009;27:858–863.
44. Doi A, Park I-H, Wen B, et al. Differential methylation of tissue- and cancer-specific CpG island shores distinguishes human induced pluripotent stem cells, embryonic stem cells and fibroblasts. *Nat Genet*. 2009;41:1350–1353.
45. Mirealdi A, Oestergaard MZ, Pharoah PD, Caldas C. Cancer genetics of epigenetic genes. *Hum Mol Genet*. 2007;16:R28–R49.
46. Vaquero A, Sternglanz R, Reinberg D. NAD<sup>+</sup>-dependent deacetylation of H4 lysine 16 by class III HDACs. *Oncogene*. 2007;26:5505–5520.
47. Espada J, Ballestar E, Santoro R, et al. Epigenetic disruption of ribosomal RNA genes and nucleolar architecture in DNA methyltransferase 1 (Dnmt1) deficient cells. *Nucleic Acids Res*. 2007;35:2191–2198.
48. Noonan E, Place R, Pookot D, et al. miR-449a targets HDAC-1 and induces growth arrest in prostate cancer. *Oncogene*. 2009;28:1714–1724.
49. Kondo Y, Shen L, Suzuki S, et al. Alterations of DNA methylation and histone modifications contribute to gene silencing in hepatocellular carcinomas. *Hepatology Res*. 2007;37:974–983.
50. Dalglish GL, Furge K, Greenman C, et al. Systematic sequencing of renal carcinoma reveals inactivation of histone modifying genes. *Nature*. 2010;463:360–363.
51. Wang P, Lin C, Smith ER, et al. Global analysis of H3K4 methylation defines MLL family member targets and points to a role for MLL1-mediated H3K4 methylation in the regulation of transcriptional initiation by RNA polymerase II. *Mol Cell Biol*. 2009;29:6074–6085.
52. Mulero-Navarro S, Esteller M. Chromatin remodeling factor CHD5 is silenced by promoter CpG island hypermethylation in human cancer. *Epigenetics*. 2008;3:210–215.
53. Sporn JC, Kustatscher G, Hothorn T, et al. Histone macroH2A isoforms predict the risk of lung cancer recurrence. *Oncogene*. 2009;28:3423–3428.