

Real analysis

1. Sets, relations, functions, and order

1.1. Logic

Theorem 1.1.1 (De Morgan's Laws for statements)

AND and OR interchange under complementation.

In C notation:

$$!(A \ || \ B) == (!A) \ \&\& \ (!B)$$

$$!(A \ \&\& \ B) == (!A) \ || \ (!B)$$

In mathematical logical notation:

$$\neg(A \vee B) = (\neg A) \wedge (\neg B)$$

$$\neg(A \wedge B) = (\neg A) \vee (\neg B)$$

Theorem 1.1.2 (De Morgan's Laws for set theory)

Union and intersection interchange under complementation.

$$(A \cup B)^c = A^c \cap B^c$$

$$(A \cap B)^c = A^c \cup B^c$$

Theorem 1.1.3

Given the true statement $P \implies Q$,

- $\neg P \implies \neg Q$ is the **inverse**, which might be true or false.
- $Q \implies P$ is the **converse**, which is true iff the inverse is true.
- $\neg(P \implies Q)$ is the **negation**, which is always false.
- $\neg Q \implies \neg P$ is the **contrapositive**, which is always true.

1.2. Sets

See 04 Set Theory, [Topology] 01 Set Theory

A set is defined by selecting elements from another set such that $S(x)$ is true:

$$\{x \in A : S(x)\} \text{ or } \{x \in A \mid S(x)\}$$

$\emptyset$ denotes the empty set.

$P \implies Q$ means P implies Q (if P , then Q).

$P \Leftrightarrow Q$ means P and Q are equivalent (P iff Q).

$A \subset B$ means A is a subset of B (**$A = B$ is allowed**).

$A \subsetneq B$ means A is a strict/proper subset of B .

$A = B$ means $A \subset B$ and $B \supset A$.

$A \cup B$ means the union of A and B .

$A \cap B$ means the intersection of A and B .

$A \parallel B$ means $A \cap B = \emptyset$.

$A \setminus B = \{x \in A : x \notin B\}$, everything in A that's not in B .

$A \times B = \{(a, b) : a \in A, b \in B\}$ is called the **Cartesian product** of A and B .

Definition 1.2.1 (Power set)

The set of all subsets of A , denoted $\mathcal{P}(A)$.

1.3. Ordered pair

See 04 Set Theory

Definition 1.3.1 (Ordered pair)

The **ordered pair** (a, b) is the set $\{\{a\}, \{a, b\}\}$.

Theorem 1.3.2

$$(a, b) = (c, d) \iff a = c \text{ and } b = d.$$

1.4. Relations

See 04 Set Theory, [Topology] 02 Relations

Definition 1.4.1 (Relation)

A **relation** $R : A \rightarrow B$ is any subset of $A \times B$.

aRb means $(a, b) \in R$.

Theorem 1.4.2 (Relations can compose)

If $R : A \rightarrow B$ and $S : B \rightarrow C$, then $S \circ R : A \rightarrow C$ is a subset of $A \times C$.

Warning

$R \circ R^{-1}$ is not necessarily the identity map!

Theorem 1.4.3

$$(S \circ R)^{-1} = R^{-1} \circ S^{-1}$$

$$(T \circ S) \circ R = T \circ (S \circ R)$$

Definition 1.4.4

R is **reflexive** iff aRa for all $a \in A$.

R is **symmetric** iff $aRb \Leftrightarrow bRa$.

R is **transitive** iff $aRb, bRc \Rightarrow aRc$.

1.4.1. Equivalence relations

Definition 1.4.5 (Equivalence relation)

A relation that is reflexive, symmetric, and transitive.

Definition 1.4.6 (Equivalence class)

Let $R : A \rightarrow B$ be an equivalence relation. Then the **equivalence class** of $a \in A$ is

$$[a] = [a]_R := \{b \in A : aRb\}$$

The set of all equivalence relations of R in A is called A/R .

Lemma 1.4.7

Two equivalence classes are either identical or disjoint.

Definition 1.4.8 (Partition)

A collection of disjoint nonempty subsets of A whose union is all of A .

Theorem 1.4.9

Given an equivalence relation on nonempty A , its set of equivalence classes forms a partition of A .

Given a partition P of A , there is a unique equivalence relation whose set of equivalence classes forms P .

1.4.2. Functions (maps)

See 03 Functions

Definition 1.4.10 (Function)

A **function** (also called a **map** or **mapping**) is a relation $f : A \rightarrow B$ where each $a \in A$ appears as the first element (input) of an ordered pair $(a, b) = afb$ exactly once.

If $(a, b) \in f$, we can also say $f(a) = b$. b is the **value** of f at a , or the **image** of a under f .

A is the **domain** and B the **codomain** of f .

Definition 1.4.11 (Injective)

A function f is **injective** or **one-to-one** iff:

- $f(a) = f(b) \Rightarrow a = b$, or equivalently
- $a \neq b \Rightarrow f(a) \neq f(b)$

Definition 1.4.12 (Surjective)

A function $f : A \rightarrow B$ is **surjective** or **onto** iff:

- $f(A) = B$ (its image is its codomain), or equivalently
- for every $b \in B$ there exists $a \in A$ s.t. $f(a) = b$

Definition 1.4.13 (Bijective)

A function is **bijective** and has **one-to-one correspondence** if it is both injective and surjective.

Theorem 1.4.14

A composition of two functions is a function. If both are sur/in/bijective, so is their composition.

Definition 1.4.15

The set of all functions $A \rightarrow B$ is denoted B^A .

Lemma 1.4.16

If A and B are finite, then the size of B^A is $|B|^{|A|}$, the size of B to the size of A .

Theorem 1.4.17

For any set A , there is a bijection between $\mathcal{P}(A)$ and 2^A .

Definition 1.4.18 (Induced map)

The **induced map** of the relation $f : A \rightarrow B$ is a function $\bar{f} : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ defined as

$$\bar{f}(X) := \{b \in B : b = f(a) \text{ for some } a \in X\}$$

$\bar{f}(X)$ is called the **image** of X . This is sometimes denoted $f(X)$ (which is an abuse of notation).

$\bar{f}$ is always a function regardless of if f is a function.

1.5. Sequence

Definition 1.5.1 (Sequence)

A **sequence** is a function whose domain is the natural numbers, or some subset of the natural numbers.

If a is a sequence, $a(n)$ is also denoted a_n , and is called the n th element of a .

1.6. Cardinality

See 06 Counting and Sizes of Sets

Assume that there is some “overset” containing all of the sets we consider.

Definition 1.6.1 (Cardinality)

$\text{card}(A)$ is the equivalence class of A under bijection.

Equivalently, $\text{card}(A) = \text{card}(B)$ iff there is a bijection $A \rightarrow B$.

Two sets are **equipotent**, **equinumerous**, **equipollent**, or **similar** if they have the same cardinality.

Definition 1.6.2

$\text{card } A \leq \text{card } B$ iff there is an injection from A to B .

Theorem 1.6.3 (Schroeder-Bernstein Theorem)

$$\begin{aligned} \text{card}(A) \leq \text{card } B \text{ AND } \text{card}(B) \leq \text{card}(A) \\ \implies \text{card}(A) = \text{card}(B) \end{aligned}$$

Theorem 1.6.4 (Cantor's Theorem)

For any set A ,

$$\text{card}(A) < \text{card}(\mathcal{P}(A))$$

Theorem 1.6.5

Every infinite set is similar to a proper subset of itself.

No finite set is similar to a proper subset of itself.

1.7. Countability

Definition 1.7.1 (Countability)

S is countable iff there is an injection $f : S \rightarrow \mathbb{Z}^+$, or equivalently iff $\text{card } S \leq \text{card } \mathbb{Z}^+$.

Corollary 1.7.1.1

1. Every subset of a countable set is countable.
 - The elements of a countable set can be indexed by the positive integers.
 - All finite sets are countable.
2. A countable union of countable sets is countable.

Theorem 1.7.2

The product of a finite number of countable sets is countable.

However, a product of an infinite number of countable sets is not countable.

Theorem 1.7.3

$$\text{card}(\mathbb{R}) = \text{card}(\mathcal{P}(\mathbb{Z}^+))$$

Thus, the reals are not countable.

1.8. Axiom of choice

Axiom 1.8.1 (Axiom of Choice)

Given a infinite collection of disjoint finite sets, it is possible to select one element from each of the sets.

Equivalently, if A is a set of nonempty sets, there is always a function $f : A \rightarrow \bigcup A$ such that $f(a) \in a$ for all $a \in A$.

In this class, we assume the Axiom of Choice.

Theorem 1.8.2 (Zermelo's Theorem)

The following theorems are equivalent:

- the Axiom of Choice
- the Well-Ordering Property
- the Hausdorff Maximum Principle
- Zorn's Lemma

Theorem 1.8.3 (Well-Ordering Property (Zermelo's Postulate))

Every set can be well-ordered.

9

Theorem 1.8.4 (Hausdorff Maximum Principle)

Every poset has a “maximal” totally ordered subset, i.e. a totally ordered subset that cannot be extended without escaping the poset.

Theorem 1.8.5 (Zorn's Lemma)

If A is a poset in which every totally ordered subset has an upper bound, then A has a maximal element, i.e. there is no larger element in the poset.

2. Fields and order

2.1. Equality

Definition 2.1.1 (Equality)

Equality is an equivalence relation on the reals, so:

- **Reflexive:** $x = x$.
- **Symmetric:** $x = y \implies y = x$.
- **Transitive:** $x = y$ and $y = z \implies y = z$.

2.2. Fields

Definition 2.2.1 (Field)

A *field* $\mathbb{F}$ is any set equipped with two relations, **addition** (+) and **multiplication** (*), which satisfy the following properties:

- **Closure axiom:** If x and y are in $\mathbb{F}$, so are both $x + y$ and xy .
- **Commutativity axioms:** $x + y = y + x$ and $xy = yx$.
- **Associativity axioms:** $(x + y) + z = x + (y + z)$ and $(xy)z = x(yz)$.
- **Distributivity axiom:** $x(y + z) = xy + xz$.
- **Substitution axioms:** $w = x$ and $y + z \implies w + y = x + z$ and $wy = xz$.
- The **additive inverse axiom** and **multiplicative inverse axiom**, defined below.

Axiom 2.2.2 (Additive inverse axiom)

Given $x, y \in \mathbb{F}$, there exists $z \in \mathbb{F}$ s.t. $x + z = y$ (i.e. all addition problems have solutions).

We denote this z by the expression $(y - x)$. We also denote, for every x :

$$0_x := (x - x)$$

$$-x := (0_x - x)$$

Theorem 2.2.3 (Zero is unique)

There is some $0 \in \mathbb{F}$ s.t. $0 = 0_x = 0_y$ for all x and y in $\mathbb{F}$.

Axiom 2.2.4 (Multiplicative inverse axiom)

There exists some number in $\mathbb{F}$ which is not equal to 0. If $x, y \in \mathbb{F}$ s.t. $x \neq 0$, then there exists a real number z s.t. $xz = y$. This z is denoted (y/x) .

The number (x/x) is denoted 1_x , and $(1_x/x)$ is denoted x^{-1} and called the **reciprocal** of x .

Theorem 2.2.5 (One is unique)

There is some $1 \in \mathbb{F}$ s.t. for all nonzero x and y , $1 = 1_x = 1_y$.

Lemma 2.2.6

In any field $\mathbb{F}$, the following hold:

- $0x = 0$ for any x
- $0/x = 0$ for any nonzero x
- $-(-x) = x$
- $(x^{-1})^{-1} = x$ for any nonzero x
- $-(x - y) = y - x$
- $x - y = x + (-y)$
- $-(x + y) = (-x) + (-y)$
- $-x = -1x$

2.3. Order

Definition 2.3.1 (Order)

Let S be a set. An **order** $<$ is a relation that satisfies the order axioms:

- **Trichotomy:** For any $x, y \in S$, exactly one of:
 - $x < y$
 - $x = y$
 - $x > y$
- **Transitivity:** If $x < y$ and $y < z$, then $x < z$.
- **Translation:** If $x < y$, then for any $x, x + z < y + z$.
- **Positive closure:** If $x > 0$ and $y > 0$ then $xy > 0$.

Definition 2.3.2 (Upper bound)

u is an **upper bound** for A iff $a \leq u$ for all $a \in A$.

Definition 2.3.3 (Supremum)

The **supremum** or **least upper bound** of A , denoted $\sup A$, is the unique object s.t. for every upper bound u of A , $\sup A \leq u$.

Definition 2.3.4 (Lower bound)

l is an **lower bound** for A iff $l \leq a$ for all $a \in A$.

Definition 2.3.5 (Infimum)

The **infimum** or **greatest lower bound** of A , denoted $\inf A$, is the unique object s.t. for every upper bound l of A , $\inf A \leq l$.

Theorem 2.3.6

If a set has a maximum, the maximum is its supremum.

If a set has a minimum, the minimum is its infimum.

Definition 2.3.7 (Dedekind completeness)

An ordered set is **complete**, **Dedekind-complete**, or has the **least upper bound property** iff:

- every subset of the set that has an upper bound has a supremum, or equivalently
- every subset of the set that has a lower bound has an infimum.

2.4. Ordered fields

Definition 2.4.1 (Ordered field)

A field $\mathbb{F}$ is an **ordered field** iff it has an order $<$ **compatible with arithmetic**, i.e. satisfying the properties:

- **Translation:** If $x < y$, then $x + z < y + z$ for all z .
- **Positive closure:** If $x > 0$ and $y > 0$, then $xy > 0$.

An element x is **positive** if $x > 0$ and **negative** if $x < 0$. The set containing all positive elements of $\mathbb{F}$ is denoted $\mathbb{F}^+$, and the set containing all negative elements of $\mathbb{F}$ is denoted $\mathbb{F}^-$.

Lemma 2.4.2

- $x > 0$ and $y > 0 \implies x + y > 0$
- $x > 0 \implies -x < 0$
- $-x > 0 \implies x < 0$
- 1 is positive
- $x > y$ and z is positive $\implies xz > yz$
- $x > y > 0 \implies 0 < x^{-1} < y^{-1}$

Theorem 2.4.3

If $a, b \in \mathbb{F}$ and $a \leq b + \varepsilon$ for every positive ε , then $a \leq b$ (if $\mathbb{F}$ is an ordered field).

2.5. Real numbers

Theorem 2.5.1

There is a unique complete ordered field, called $\mathbb{R}$, the **real numbers**.

Theorem 2.5.2 (Approximation Property)

Let S be a nonempty set of reals with a supremum. Then for every real number $a < \sup S$ there exists $x \in S$ with $a < x \leq \sup S$.

Theorem 2.5.3 (Additive Property)

Let A and B be sets of reals with suprema. Consider

$$C := \{a + b : a \in A, b \in B\}.$$

Then,

$$\sup C = \sup A + \sup B.$$

Theorem 2.5.4 (Comparison Property)

Let S and T be two nonempty sets of reals where $s \leq t$ for all $s \in S, t \in T$. Then, if T has a supremum, S also has a supremum and $\sup S \leq \sup T$.

Lemma 2.5.5

The positive integers have no upper bound.

Proof. Assume that $z = \sup \mathbb{Z}^+$ exists. We know that $z - 1 < z$, so by the approximation property there is a positive integer s with $z - 1 < s$. Thus $z < s + 1$. $\square$

Definition 2.5.6 (Comensural)

The set X is **comensural** iff for any $x, y \in X$ with $x > 0$, then there is a positive integer n s.t. $nx > y$.

Theorem 2.5.7 (Archimedean property)

The reals are comensural.

Proof. Consider $\frac{y}{x}$. If $\frac{y}{x} < 0$, then $n = 0$. Otherwise, $\frac{y}{x} > 0$. Since the positive integers are unbounded, choose $n > \frac{y}{x}$. Then $nx > y$. $\square$

Theorem 2.5.8 (Density of the rationals / rationals between reals)

Between any two reals there is a rational number.

Proof. The other cases are trivial, so consider $0 < x < y$. Choose an integer $n > \frac{1}{y-x}$.

We define $q := \frac{k}{n}$, and

We claim $x < q$. Then:

$$\frac{1}{y-x} < n$$

$$y-x > \frac{1}{n}$$

$$y - \frac{1}{n} > x$$

$$y - q \leq \frac{1}{n} < y - x$$

$\square$

2.6. Extended reals

Definition 2.6.1 (Extended reals)

The set $\mathbb{R}^*$ consisting of the reals, plus two elements $-\infty$ and $+\infty$.

The order on it is defined such that

$$-\infty < x < +\infty \text{ for all } x \in \mathbb{R}$$

and when comparing two reals it still has the standard order.

All expressions whose value is defined follow the rules of complete ordered fields, but there are many undefined expressions.

2.7. Absolute values

Theorem 2.7.1 (Corollaries of the Triangle Inequality)

For any $x, y \in \mathbb{R}$:

$$|x + y| \leq |x| + |y|$$

$$|x - y| \leq |x| + |y|$$

$$||x| + |y|| \leq |x - y|$$

2.8. Bounded function

Definition 2.8.1 (Bounded function)

$f : S \rightarrow \mathbb{R}$ is **bounded** iff the image $f(S)$ has both a supremum and infimum.

Equivalently, f is bounded iff there exists M s.t. $|f(x)| < M$ for all x .

Equivalently, f is bounded iff $f(S)$ is bounded.

Theorem 2.8.2

If $f(x) \leq g(x)$ for all $x \in D$, then $\sup f(D) \leq \sup g(D)$.

2.9. Decimal representation

Definition 2.9.1 (Decimal representation)

A string of the form

$$(-)d_n d_{n-1} \dots d_1 d_0 . d_{-1} d_{-2} \dots$$

where each $d_i \in \{0, 1, 2, \dots, 9\}$.

i.e.

- optionally, a minus sign
- a finite string of digits
- a decimal point
- an infinite string of digits

Theorem 2.9.2

There is a bijection between $\mathbb{R}^+$ and the set of decimal representations without a minus sign and such that there is a last 9, i.e. the set of nonterminating positive decimal representations.

In order to prove this, we need some lemmas:

Lemma 2.9.3

There is a bijection between $\mathbb{Z}^+$ and the set of decimal representations ending in an infinite string of 0s.

Proof. Every integer is representable as

$$n = 10m + r.$$

So keep dividing the integer by 10, and pushing the remainder onto the front of the string. This gives us a base-10 representation of the integer, i.e. a finite string of digits.

To make a full decimal representation, put an infinite string of 0s after the decimal point. $\square$

Now let's prove the original theorem.

Proof. Let $x \in \mathbb{R}^+$. Take $n = \lfloor x \rfloor$ and find its decimal representation. Then $0 < x - n < 1$. $\square$

Now we need to prove more lemmas:

Lemma 2.9.4

Every decimal representation represents a real number.

3. Sequences

Definition 3.1 (Sequence)

A function $x : \mathbb{Z}^+ \rightarrow \mathbb{R}$.

$x(n)$ is usually denoted x_n and called the n th element of the sequence. The sequence itself is often denoted $\{x_n\}$, $\{x_n\}_{n=1}^\infty$, (x_n) , or similar.

If the terms of the sequence are specified by a formula, the sequence might be left unnamed and written as, for example, $\{\frac{1}{n}\}_{n=1}^\infty$.

Definition 3.2 (Limit of a sequence)

Let $\{x_n\}$ be a sequence. L is the **limit**, denoted $\lim_{n \rightarrow \infty} x_n$, of the sequence iff for every $\varepsilon > 0$ there exists N s.t. whenever $n > N$, $|x_n - L| < \varepsilon$. If such a limit exists, the sequence **converges** to that limit.

A convergent sequence has exactly one limit.

If a sequence has no limit, it **diverges**.

Theorem 3.3

If a sequence is unbounded, it diverges.

Definition 3.4 (Monotonic)

Let $\{x_n\}$ be a sequence. If $x_n \leq x_{n+1}$ for all n , the sequence is **non-decreasing** or **increasing**. If $x_n < x_{n+1}$ for all N , the sequence is **strictly increasing**. If $x_n \geq x_{n+1}$ for all n , the sequence is **non-increasing** or **decreasing**. If $x_n > x_{n+1}$ for all N , the sequence is **strictly decreasing**.

In all these cases, the sequence is **monotonic** or **monotone**, and if it is strictly increasing or strictly decreasing, then the sequence is **strictly monotonic**.

Theorem 3.5

A bounded monotonically increasing sequence converges to its supremum.

A bounded monotonically decreasing sequence converges to its infimum.

Theorem 3.6

Every nonempty bounded subset of the reals has an increasing sequence converging to its supremum and a decreasing sequence converging to its infimum.

3.1. Subsequences

Definition 3.1.1 (Subsequence)

If $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ is a strictly increasing function and $\{x_n\}$ is a sequence, $\{x_{k(n)}\}$ is a **subsequence** of $\{x_n\}$.

Theorem 3.1.2

The following are equivalent:

1. The sequence $\{x_n\}$ converges.
2. The t -tail $\{x_{n+t}\}$ converges for every t .
3. There is some t for which the t -tail converges.

3.2. Limit properties and theorems

Theorem 3.2.1

Let $\{x_n\} \rightarrow x$ and $\{y_n\} \rightarrow y$, s.t. $x_n \leq y_n$ for all n . Then, $x \leq y$.

Theorem 3.2.2 (Squeeze Theorem)

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences s.t. $a_n \leq b_n \leq c_n$ for all n and $a_n \rightarrow L$ and $c_n \rightarrow L$. Then $b_n \rightarrow L$.

Proof. Let $\varepsilon > 0$. Choose n so large that both $|a_n - L|, |c_n - L| < \varepsilon$. Then,

$$-\varepsilon < c_n - L < \varepsilon$$

$$-\varepsilon < a_n - L < \varepsilon$$

and thus

$$-\varepsilon < a_n - L \leq b_n - L \leq c_n - L < \varepsilon$$

so clearly

$$|b_n - L| < \varepsilon.$$

Since this works for all ε , $b_n \rightarrow L$. □

Theorem 3.2.3

Let $\{x_n\} \rightarrow x$ and $\{y_n\} \rightarrow y$. Then

1. $\{x_n + y_n\} \rightarrow x + y$.
2. $\{x_n - y_n\} \rightarrow x - y$.
3. $\{x_n y_n\} \rightarrow xy$.
4. $\{x_n / y_n\} \rightarrow x / y$ if $y_n \neq 0$ and $y \neq 0$ for all n .

Theorem 3.2.4

Let $\{x_n\}$ be a sequence and x a number. Then,

$$\{x_n\} \rightarrow x \iff \{x_n - x\} \rightarrow 0.$$

Theorem 3.2.5 (Convergence of geometric series)

Let $c > 0$. Then,

1. $c < 1 \implies \lim_{n \rightarrow \infty} c^n = 0$.
2. $c > 1 \implies \{c^n\}$ is unbounded.

Theorem 3.2.6 (Ratio Test)

Let $\{x_n\}$ be an always-nonzero sequence s.t.

$$\lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|} = L.$$

Then,

- $L < 1 \implies \{x_n\} \rightarrow 0$.
- $L > 1 \implies \{x_n\}$ is unbounded.

Theorem 3.2.7

$$\lim_{n \rightarrow \infty} n^{\frac{1}{n}} = 1$$

3.3. Limsup and liminf

Definition 3.3.1 (Limsup and liminf)

Let $\{x_n\}$ be a bounded sequence. For each n , define

$$a_n := \sup\{x_k : k \geq n\}$$

$$b_n := \inf\{x_k : k \geq n\}$$

Thus, a_n is above the n -tail of the sequence, while b_n is below it.

Define

$$\limsup x_n = \lim_{n \rightarrow \infty} a_n$$

$$\liminf x_n = \lim_{n \rightarrow \infty} b_n.$$

We know that $\limsup x_n \geq \liminf x_n$.

Theorem 3.3.2

If $\{x_n\}$ is bounded, then:

- there is a subsequence $\{x_{n_k}\} \rightarrow \limsup x_n$.
- there is a subsequence $\{x_{m_k}\} \rightarrow \liminf x_n$.

Theorem 3.3.3

Let $\{x_n\}$ be a bounded sequence with $\limsup x_n = \liminf x_n = L$. Then, $\{x_n\} \rightarrow L$.

Theorem 3.3.4

$\{x_n\} \rightarrow L$ iff every subsequence converges to L .

3.4. Bolzano-Weierstrass

Theorem 3.4.1 (Bolzano-Weierstrass for Sequences)

Every bounded sequence of real numbers (or in $\mathbb{R}^n$) has a convergent subsequence.

In topology terms, a subset of $\mathbb{R}^n$ is sequentially compact iff it is closed and bounded.

Definition 3.4.2 (Open ball)

Let $x \in \mathbb{R}^n$. The ball $B(x, \varepsilon)$ of radius ε about x is defined as

$$B(x, \varepsilon) := \{y \in \mathbb{R}^n : |x - y| < \varepsilon\}.$$

Definition 3.4.3 (Accumulation point)

Let S be a set. x is a **accumulation point** of S iff every ball $B(x, \varepsilon)$ contains a point of S other than x .

Theorem 3.4.4 (Bolzano-Weierstrass theorem)

Every bounded infinite subset of $\mathbb{R}^n$ has an accumulation point.

Theorem 3.4.5 (Cantor Intersection Theorem)

Let $S_1, S_2, \dots$ be a sequence of bounded closed subsets of $\mathbb{R}^n$ nested as follows: $S_1 \supset S_2 \supset \dots$. Then, the intersection $S_1 \cap S_2 \cap \dots$ is closed and nonempty.

3.5. Infinite limits

See 16 Infinite Limits

Definition 3.5.1 (Diverging to infinity)

A sequence **diverges to** $+\infty$ iff for any $M \in \mathbb{R}$ there is an N s.t. for all $n \geq N$ then $x_n \geq M$.

A sequence **diverges to** $-\infty$ iff for any $M \in \mathbb{R}$ there is an N s.t. for all $n \leq N$ then $x_n \leq M$.

Theorem 3.5.2

An unbounded increasing sequence diverges to $+\infty$.
An unbounded decreasing sequence diverges to $-\infty$.

Definition 3.5.3

Let $\{x_n\}$ be a sequence.

$\limsup x_n = +\infty$ iff $\{x_n\}$ diverges to $+\infty$.

$\liminf x_n = -\infty$ iff $\{x_n\}$ diverges to $-\infty$.

3.6. Cauchy sequences

Definition 3.6.1 (Cauchy sequence)

A sequence $\{x_n\}$ is called **Cauchy sequence** iff for every $\varepsilon > 0$, there exists N s.t. for all $m, n > N$, $|x_n - x_m| < \varepsilon$.

In short, beyond a certain term, all the following terms are arbitrarily close.

Theorem 3.6.2

Every Cauchy sequence is bounded.

Theorem 3.6.3

A sequence of reals converges iff it is a Cauchy sequence.

4. Series

4.1. Series

Definition 4.1.1 (Series)

Given a sequence $\{x_n\}$, the object

$$\sum_{n=1}^{\infty} x_n$$

is called an **infinite series**.

Given a series $\sum_{n=1}^{\infty} x_n$, define its k^{th} **partial sum** by

$$s_k = \sum_{n=1}^k x_n := x_1 + x_2 + \cdots + x_n.$$

Since this sum is finite it is a real number.

Additionally, we can recursively define this sum as

$$\begin{aligned} s_1 &:= x_1 \\ s_{k+1} &:= s_k + x_{k+1}. \end{aligned}$$

A series **converges** iff its sequence of partial sums converges, and it converges to the same value as its sequence of partial sums. It is considered to be equal to said value.

Series with different starting indices are defined in the obvious way:

$$\sum_{n=a}^{\infty} x_n := \sum_{n=1}^{\infty} x_{n+a-1}$$

4.2. Basic convergence tests

Theorem 4.2.1 (n^{th} term test for divergence)

If $\sum x_n$ converges, then $x_n \rightarrow 0$.

Equivalently, if $x_n \not\rightarrow 0$, $\sum x_n$ diverges.

Theorem 4.2.2

Let $\sum x_n = x$ and $\sum y_n = y$. Then, $\sum (x_n + y_n) = x + y$.

For any $c \in \mathbb{R}$, $\sum (cx_n) = cx$.

4.3. Absolute convergence

See 19 Absolute Convergence

Lemma 4.3.1

If $x_n \geq 0$ for all n , then $\sum x_n$ converges iff the sequence of partial sums is bounded above.

Furthermore, each partial sum is less than or equal to the sum.

Definition 4.3.2 (Absolute convergence)

Iff $\sum |x_n|$ converges, then $\sum x_n$ **absolutely converges**. If $\sum |x_n|$ diverges, but $\sum x_n$ converges, then $\sum x_n$ **conditionally converges**.

Theorem 4.3.3

If a series converges absolutely, it converges.

Theorem 4.3.4 (Comparison test for positive series)

Let $\sum x_n$ and $\sum y_n$ be series with $0 \leq x_n \leq y_n$ for all n . Then:

1. If $\sum y_n$ converges, then $\sum x_n \leq \sum y_n$ ($\sum x_n$ converges).
2. If $\sum x_n$ diverges, so does $\sum y_n$.

Corollary 4.3.4.1

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges iff $p > 1$.

4.4. Ratio test and root test

Theorem 4.4.1 (Ratio test)

Let $\sum x_n$ be a series of nonzero terms. Suppose that

$$L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$$

exists. Then,

- If $L < 1$, $\sum x_n$ converges absolutely.
- If $L > 1$, $\sum x_n$ diverges.
- If $L = 1$, the test gives no information.

Theorem 4.4.2 (Root test)

Let $\sum x_n$ be a series and $L := \limsup \sqrt[n]{|x_n|}$ (which may be infinite). Then,

- If $L < 1$, $\sum x_n$ converges absolutely.
- If $L > 1$, $\sum x_n$ diverges.
- If $L = 1$, the test gives no information.

4.5. Alternating series

See 20 Alternating Series

Theorem 4.5.1 (Alternating series test)

Let $x_n \rightarrow 0$ be monotone decreasing. Then, $\sum (-1)^n x_n$ converges.

Theorem 4.5.2

Let $\sum x_n$ be an absolutely convergent series converging to L . Let $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ be a bijection. Then, $\sum x_{\sigma n}$ is also absolutely convergent and converges to L .

In short, if the series is absolutely convergent, we can rearrange the terms and the sum is the same.

Theorem 4.5.3 (Alternating Series Rearrangement Theorem)

Let $\sum x_n$ be a series that satisfies the alternating series test but which is not absolutely convergent. Then, for any $L \in \mathbb{R}$ there is a rearrangement of the series that sums to L .

4.6. Multiplying series

Definition 4.6.1 (Cauchy product)

The **Cauchy product** of $\sum a_n$ and $\sum b_n$ is defined as

$$\sum_{k=1}^n a_k b_{n+1-k}.$$

Essentially, we collect the terms of the product of $\sum a_n$ with $\sum b_n$ in diagonal strips when the terms are arranged in a grid.

Theorem 4.6.2 (Mertens' theorem)

Let $\sum a_n \rightarrow A$ and $\sum b_n \rightarrow B$. Then if at least one of these is absolutely convergent, then $\sum c_n \rightarrow AB$, where $\sum c_n$ is the Cauchy product of $\sum a_n$ and $\sum b_n$.

Additionally, if both converge absolutely, so does the sum of c_n s.

4.7. Power series

Definition 4.7.1 (Power series)

Given $x_0 \in \mathbb{R}$ and a sequence a_n , we can create the **power series**, a function of x :

$$\sum_{n=0}^{\infty} a_n (x - x_0)^n.$$

A power series **converges** iff there exists $x \neq x_0$ that makes the series converge. If $x = x_0$ makes the series converge, then the series converges for any x .

Theorem 4.7.2

A power series either diverges for all $x \neq x_0$, converges absolutely for all x , or there exists a real number ρ , called the **radius of convergence**, s.t. the series converges absolutely on $(x_0 - \rho, x_0 + \rho)$ and diverges if $|x - x_0| > \rho$.

When $x = x_0 + \rho$ or $x = x_0 - \rho$, this theorem is inconclusive.

5. Limits and continuity

5.1. Limits

Definition 5.1.1 (Accumulation point)

Let S be a set. x is a **accumulation point** of S iff every ball $B(x, \varepsilon)$ contains a point of S other than x .

Lemma 5.1.2

x is an accumulation point of S iff there is a sequence of points in $S - \{x\}$ converging to x .

Definition 5.1.3 (Limit)

Let S be a subset of $\mathbb{R}$ and c be an accumulation point of S . Let $f : S \rightarrow \mathbb{R}$ be a function.

The **limit** of $f(x)$ as $x \rightarrow c$ is L iff for each $\varepsilon > 0$, there exists a $\delta > 0$ s.t. whenever $x \in S - \{c\}$ and $|x - c| < \delta$, then $|f(x) - L| < \varepsilon$. If so, we say that $f(x)$ converges to L as $x \rightarrow c$.

If there is no such limit, then the limit does not exist and f **diverges**.

Limits are unique.

Theorem 5.1.4

Let S be a subset of $\mathbb{R}$ and c be an accumulation point of S . Let $f : S \rightarrow \mathbb{R}$ be a function.

Then $f(x) \rightarrow L$ as $x \rightarrow c$ iff for every sequence $\{x_n\}$ in $S - \{c\}$ whose limit is c , the sequence $\{f(x_n)\} \rightarrow L$.

5.2. Continuity

Definition 5.2.1 (Continuity)

Let $S \subset \mathbb{R}$, $c \in S$, and $f : S \rightarrow \mathbb{R}$ be a function. Then, f is **continuous** at c iff for every $\varepsilon > 0$, there is a $\delta > 0$ s.t. whenever $|x - c| < \delta$, then $|f(x) - f(c)| < \varepsilon$.

Theorem 5.2.2

Let $S \in \mathbb{R}$ and $f : S \rightarrow \mathbb{R}$ be a function. Let c be a point in S . Then:

1. If c is not an accumulation point of S , then f is continuous at c .

2. If c is an accumulation point of S , then f is continuous at c iff $\lim_{x \rightarrow c} f(x) = f(c)$.
3. f is continuous at c iff for every sequence $\{x_n\} \in S$ with limit c , $f(x_n) \rightarrow f(c)$.

Theorem 5.2.3 (Composition of continuous functions is continuous)

Let $f : S \rightarrow \mathbb{R}$. If g is continuous at $c \in T$ and f is continuous at $g(c)$, then $f \circ g : T \rightarrow \mathbb{R}$ is continuous at c .

Theorem 5.2.4

There are no functions that are continuous at every rational but discontinuous at every irrational.

5.3. Big Continuity Theorems

Definition 5.3.1 (Bounded function)

A function $f : S \rightarrow \mathbb{R}$ is bounded iff its image is bounded.

Lemma 5.3.2

Let $[a, b]$ be a closed finite interval, and f continuous on $[a, b]$. Then f is bounded.

Theorem 5.3.3 (Extreme Value Theorem)

Let $f(x)$ be a continuous function defined on a closed interval $[a, b]$. Then, f achieves an absolute maximum and minimum on $[a, b]$.

Lemma 5.3.4

Let $f(x)$ be continuous on the closed bounded interval $[a, b]$. If $f(a) < 0$ and $f(b) > 0$, then there exists $c \in [a, b]$ s.t. $f(c) = 0$.

Theorem 5.3.5 (Intermediate Value Theorem)

Let f be continuous on closed, bounded $[a, b]$. Let $y \in \mathbb{R}$ s.t. $f(a) < y < f(b)$ or $f(a) > y > f(b)$. Then, there exists $c \in (a, b)$ with $f(c) = y$.

5.4. Uniform continuity

Definition 5.4.1 (Uniform continuity)

Let $S \subset \mathbb{R}$ and $f : S \rightarrow \mathbb{R}$ be a function. For any $\varepsilon > 0$, f is **uniformly continuous** on S iff we can find δ s.t. for all $x, c \in S$ with $|x - c| < \delta$, then $|f(x) - f(c)| < \varepsilon$.

This differs from continuity iff the same δ works for every point in S .

Uniform continuity obviously implies continuity.

Theorem 5.4.2

If $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function defined on a closed bounded interval, it is uniformly continuous.

Lemma 5.4.3

If $f : S \rightarrow \mathbb{R}$ is uniformly continuous and $\{x_n\}$ is a Cauchy sequence in S , then $\{f(x_n)\}$ is a Cauchy sequence.

Theorem 5.4.4

Let $f : (a, b) \rightarrow \mathbb{R}$. Then f is uniformly continuous iff it has a right-handed limit at a , a left-handed limit at b , and the extension of f to the closed interval $[a, b]$ is continuous.

5.5. Lipschitz continuity

Definition 5.5.1 (Lipschitz continuity)

Let $f : S \rightarrow \mathbb{R}$ be a function. If there exists $K \in \mathbb{R}$ s.t. for every $x, y \in S$, $|f(x) - f(y)| < K|x - y|$ then f is **Lipschitz continuous**.

Theorem 5.5.2

Lipschitz continuity implies uniform continuity.

6. Derivatives

Definition 6.1 (Derivative)

Let $f(x)$ be a real-valued function defined on an open interval (a, b) , and let $x_0 \in (a, b)$. The **derivative** of f at x_0 is

$$\frac{df}{dx}(x_0) := f'(x_0) := \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}.$$

Iff this limit is defined, the function is **differentiable** at x_0 .

Theorem 6.2 (Continuity theorem)

Let f be defined on (a, b) with $x_0 \in (a, b)$. Then, f is differentiable at x_0 iff there is a function f^* (which depends on f and on x_0) s.t.

$$f(x) = f(x_0) + f^*(x)(x - x_0)$$

with f^* defined on (a, b) and continuous at x_0 , with $f^*(x_0) = f'(x_0)$.

If f is continuous, f^* can be defined as

$$f^*(x) := \begin{cases} \frac{f(x) - f(x_0)}{x - x_0} & \text{if } x \neq x_0 \\ f'(x_0) & \text{if } x = x_0. \end{cases}$$

Definition 6.3 (One-sided derivative)

Let f be defined on $[a, b)$. The **right-hand derivative** of f at a is

$$f'_+(a) := \lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{x - a}.$$

Let f be defined on $(a, b]$. The **left-hand derivative** of f at a is

$$f'_-(a) := \lim_{x \rightarrow a^-} \frac{f(x) - f(a)}{x - a}.$$

Definition 6.4 (Infinite derivatives)

Let f be continuous on $[a, b)$. The right-hand derivative of f at a is $+\infty$ iff

$$\lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{x - a} = +\infty.$$

The definition is similar for left-hand and negative infinite derivatives.

Theorem 6.5 (Sum, product, and quotient rules)

Let f and g be functions defined on an interval (a, b) , both differentiable at the point $x_0 \in (a, b)$. (In particular, the derivatives there cannot be infinite.) Then, the following derivatives exist and take on the following values:

- $(f \pm g)'(x_0) = f'(x_0) \pm g'(x_0)$.
- $(f * g)'(x_0) = f(x_0)g'(x_0) + f'(x_0)g(x_0)$.
- $(f/g)'(x_0) = \frac{g(x_0)f'(x_0) - f(x_0)g'(x_0)}{g(x_0)^2}$.

Theorem 6.6 (Chain rule)

Let $f : (a, b) \rightarrow \mathbb{R}$ and $g : f(a, b) \rightarrow \mathbb{R}$. Let $x_0 \in (a, b)$ with f differentiable at x_0 and $f(x_0)$ contained in the interior of $f(a, b)$, and let g be differentiable at $f(x_0)$. Then, $(g \circ f)(x)$ is differentiable at x_0 , and

$$(g \circ f)'(x_0) = g'(f(x_0))f'(x_0),$$

or equivalently,

$$\left. \frac{dg}{dx} \right|_{x_0} = \left. \frac{dg}{df} \right|_{f(x_0)} * \left. \frac{df}{dx} \right|_{x_0}.$$

6.1. Mean value theorem

Theorem 6.1.1

Let $f'(c) > 0$ and f continuous in some interval about c . Then, there is a $\delta > 0$ s.t.:

- $c < x < c + \delta \implies f(c) < f(x)$
- $c - \delta < x < c \implies f(x) < f(c)$

Equivalently, if the derivative is positive, then the function is increasing at that point.

Similarly, if the derivative is negative, then the function is increasing at that point.

Corollary 6.1.1.1

If $f(c)$ is a local min or local max, then $f'(c) = 0$.

Definition 6.1.2 (Critical point)

A point where the derivative of the function is zero.

Theorem 6.1.3 (Rolle's theorem)

Let $f(x)$ be continuous on $[a, b]$ and differentiable on (a, b) , s.t. $f(a) = f(b)$. Then, there exists $c \in (a, b)$ for which $f'(c) = 0$.

Theorem 6.1.4 (Mean value theorem)

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then, there exists some point $c \in (a, b)$ for which

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

Theorem 6.1.5 (Cauchy's mean value theorem)

Let $f(t)$ and $g(t)$ be two functions continuous on $[a, b]$ and differentiable on (a, b) . Then, there exists some point $c \in (a, b)$ for which

$$f'(c)(g(b) - g(a)) = g'(c)(f(b) - f(a)).$$

Theorem 6.1.6

Let $f(x)$ be continuous on $[a, b]$ and differentiable on (a, b) . Then:

- $f'(x) > 0$ for all $x \in (a, b) \iff f$ is strictly increasing
- $f'(x) < 0$ for all $x \in (a, b) \iff f$ is strictly decreasing
- $f'(x) = 0$ for all $x \in (a, b) \iff f$ is constant

Let $g(x)$ also be continuous on $[a, b]$ and differentiable on (a, b) . Then:

- $g'(x) = f'(x)$ for all $x \in (a, b) \iff f(x) - g(x)$ is constant

Theorem 6.1.7

Let $f(x)$ be continuous on $(a, b]$ and differentiable on (a, b) , with $\lim_{x \rightarrow b^-} f'(x) = L$. Then, f is left-hand differentiable at b , and $f'_-(b) = L$.

A similar theorem exists for right-hand derivatives.

Theorem 6.1.8 (Darboux theorem)

Let $f(x)$ be a function that is differentiable on $[a, b]$. Let y be a value s.t. $f'(a) < y < f'(b)$ or $f'(a) > y > f'(b)$. Then there is some $c \in (a, b)$ s.t. $f'(c) = y$.

Corollary 6.1.8.1

- f' cannot change signs without going through zero.
- If $f'(x) \neq 0$ for any x , then f is monotonic.
- If f' itself is monotonic then it must be continuous:
 - the only kinds of discontinuities of monotonic functions are jumps
 - derivatives can't make jumps

6.2. Taylor series

Theorem 6.2.1 (Taylor's formula with the Lagrange form of the remainder)

Let $f : (a, b) \rightarrow \mathbb{R}$ have $n + 1$ derivatives on the interval (a, b) , and let x_0 be a point in (a, b) . Then, for any point y in (a, b) **Taylor's formula** states that

$$f(y) = p_n(y) + R_n(y; x_0).$$

We define the **n th Taylor polynomial centered at x_0** as:

$$p_n(x) := \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k.$$

We define the **Lagrange form of the remainder** as:

$$R_{n(x; x_0)} := \frac{f^{(n+1)}(c)(y - x_0)^{n+1}}{(n + 1)!}$$

for some c between y and x_0 [this theorem does not specify which c this is].

Definition 6.2.2 (Taylor series)

If f is infinitely differentiable, its **Taylor series** around x_0 is the sequence of Taylor polynomials centered at x_0 :

$$p_1, p_2, p_3, \dots$$

Ideally, this would converge to some nice function, and the error $R_n(x; x_0)$ would converge to 0. But there are many functions for which this is not the case; some such functions are not really that nasty.

7. Integrals

7.1. Partitions

Definition 7.1.1 (Partition)

A **partition** P of $[a, b]$ is a finite set of points, $\{x_0, x_1, \dots, x_n\}$ with $a = x_0 < x_1 < \dots < x_n = b$. The k th interval of the partition is $[x_{k-1}, x_k]$, with width $\Delta x_k := x_k - x_{k-1}$.

Definition 7.1.2 (Norm/mesh)

The **norm** or **mesh** of a partition is the width of its largest interval, denoted

$$\|P\| := \max\{\Delta x_k : k \leq n\}.$$

Definition 7.1.3 (Refinement)

$\tilde{P}$ is a **refinement** of P iff $P \subseteq \tilde{P}$.

7.2. Darboux integral

Definition 7.2.1 (Darboux sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded and P a partition of $[a, b]$. Then, define

$$m_k := \inf\{f(x) : x_{k-1} \leq x \leq x_k\}$$

$$M_k := \sup\{f(x) : x_{k-1} \leq x \leq x_k\}.$$

Define the **upper** and **lower Darboux sums**, respectively, as

$$L(P, f) := \sum_{k=1}^n m_k \Delta x_k$$

$$U(P, f) := \sum_{k=1}^n M_k \Delta x_k.$$

Definition 7.2.2 (Upper and lower Darboux integrals)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Then, define the **upper** and **lower Darboux integrals** as follows:

$$\int_a^b f(x) dx := \inf\{U(P, f) : P \text{ is a partition of } [a, b]\}$$

$$\int_a^b f(x) dx := \sup\{L(P, f) : P \text{ is a partition of } [a, b]\}.$$

Definition 7.2.3 (Darboux integrability)

If $f : [a, b] \rightarrow \mathbb{R}$ is bounded, and $\int_a^b f(x) dx = \int_a^b f(x) dx$, then we say f is **Darboux-integrable** ($f \in \mathcal{R}[a, b]$), and we define $\int_a^b f(x) dx$ to be the common value of the lower and upper Darboux integral.

7.3. Riemann integral

Definition 7.3.1 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded and P a partition of $[a, b]$.

For each interval $[x_{i-1}, x_i]$, select any t_i in that interval. Then, the **Riemann sum** of f under this selection of t_i is

$$\sum_{i=1}^n f(t_i) \Delta x_i.$$

Definition 7.3.2 (Riemann integrability)

f is **Riemann-integrable** ($f \in \mathcal{R}[a, b]$) iff there exists some number I s.t. for every $\varepsilon > 0$, there exists a partition P_0 s.t. for every refinement P of P_0 and every choice of t_i on P , the corresponding Riemann sum is within ε of I .

Theorem 7.3.3

Riemann integrability is equivalent to Darboux integrability.

7.4. Integrability

Theorem 7.4.1

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Then, f is Darboux-integrable iff for every $\varepsilon > 0$, there exists a partition P s.t. $U(P, f) - L(P, f) < \varepsilon$.

Theorem 7.4.2

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded and monotonic. Then, $f \in \mathcal{R}[a, b]$.

7.5. Core properties of the integral

See 35 More Integral Basics

The integral operator is additive, linear, and monotonic:

Lemma 7.5.1

Let $a < b < c$ and $f : [a, c] \rightarrow \mathbb{R}$ be a bounded function. Then, both the upper and lower Darboux integrals are additive over the subintervals. That is,

$$\int_a^c f = \int_a^b f + \int_b^c f$$

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

Theorem 7.5.2 (Additivity)

Let $a < b < c$ and $f : [a, c] \rightarrow \mathbb{R}$ be a bounded function. Then,

$$f \in \mathcal{R}[a, c] \iff f \in \mathcal{R}[a, b] \text{ and } f \in \mathcal{R}[b, c],$$

and if f is integrable on the whole interval then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

Lemma 7.5.3

If f and g are bounded functions on $[a, b]$, then

$$\overline{\int_a^b (f + g)} \leq \overline{\int_a^b f} + \overline{\int_a^b g}.$$

Theorem 7.5.4 (Linearity)

Let $f, g \in \mathcal{R}[a, b]$, and α any real constant. Then,

$$\int_a^b \alpha f(x) + g(x) dx = \alpha \int_a^b f(x) dx + \int_a^b g(x) dx.$$

Theorem 7.5.5 (Monotonicity)

Let f and g be bounded functions on $[a, b]$. Assume $f(x) \leq g(x)$ for all $x \in [a, b]$. Then, $\int f < \int g$ and $\overline{\int f} < \overline{\int g}$. If both are in $\mathcal{R}[a, b]$ then $\int f < \int g$.

7.6. Additional properties of the integral

See 34 Working with Integrals

Theorem 7.6.1

Let $f \in [a, b]$. Let $f(x) \in [m, M]$ for all $x \in [a, b]$. Then,

$$m(b - a) \leq \int_a^b f(x) dx \leq M(b - a).$$

Theorem 7.6.2 (Integral of a constant)

$$\int_a^b c dx = c(b - a).$$

Theorem 7.6.3 (Integral of a step function)

Let

$$f(x) = \begin{cases} r & \text{for } x < c \\ s & \text{for } x = c, \\ t & \text{for } x > c \end{cases}$$

with $c \in [a, b]$ and $f : [a, b] \rightarrow \mathbb{R}$.

Then,

$$\int_a^b f(x) dx = r(c - a) + t(b - c).$$

Definition 7.6.4

$$\int_a^b f(x) dx := - \int_b^a f(x) dx.$$

7.7. Integrals of continuous functions

See 36 Integrals of Continuous Functions

Lemma 7.7.1

A bounded function that has only finitely many discontinuities is integrable.

Lemma 7.7.2

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Let $\{a_n\} \rightarrow a$ and $\{b_n \rightarrow b\}$ be sequences in $[a, b]$, with $a_n < b_n$ for all n . Then, if $f \in \mathcal{R}[a_n, b_n]$ for all n , then $f \in \mathcal{R}[a, b]$ and $\int_a^b f = \lim_{n \rightarrow \infty} \int_{a_n}^{b_n} f$.

Essentially, bounded functions need only be integrable inside the interval to be integrable on the entire interval.

Theorem 7.7.3

Continuous functions are integrable.

7.8. Fundamental theorem of calculus

See 37 Fundamental Theorem of Calculus

Theorem 7.8.1 (First Fundamental Theorem of Calculus)

Let $F : [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable* on (a, b) , with $F' \in \mathcal{R}[a, b]$. Then,

$$\int_a^b F'(x) dx = F(b) - F(a).$$

* F may have finitely many points where it is not differentiable. It must be continuous over the whole interval.

Theorem 7.8.2 (Second Fundamental Theorem of Calculus)

Let $f \in \mathcal{R}[a, b]$. Define $F(x) := \int_a^x f$. Then, F is continuous on $[a, b]$, and if f is continuous at c , then F is differentiable at c and $F'(c) = f(c)$.

Theorem 7.8.3 (u-substitution)

Let $f : [c, d] \rightarrow \mathbb{R}$ be continuous and $g : [a, b] \subset [c, d]$ continuously differentiable. Then,

$$\int_a^b f(g(x)) g'(x) dx = \int_{g(a)}^{g(b)} f(u) du.$$

Note that in this case, $u = g(x)$ and $du = g'(x) dx$.

7.9. Logarithm and exponential

See 38 Logs and Exponentials

Definition 7.9.1 (Natural logarithm)

$$\ln(x) := \int_1^x \frac{1}{t} dt.$$

Theorem 7.9.2

$\ln(x)$ is the unique function that satisfies:

- $\ln(1) = 0$
- $\ln$ is differentiable and $L'(x) = \frac{1}{x}$
- $\ln(xy) = \ln(x) + \ln(y)$
- $\ln$ is strictly increasing; it's domain is $(0, \infty)$ and its image is $\mathbb{R}$
- If q is rational, then $\ln(x^q) = q \ln(x)$

Definition 7.9.3 (Logarithm)

$$\log_b(a) := \frac{\ln(a)}{\ln(b)}.$$

Thus,

$$\ln(x) = \log_e(x).$$

Definition 7.9.4 (Natural exponent)

$$\exp(x) := \ln^{-1}(x).$$

Equivalently, $\exp$ is the unique solution to the initial-value differential equation problem

$$y' = y, y(0) = 1.$$

Equivalently,

$$\exp(x) := \sum_{n=0}^{\infty} \frac{x^n}{n!}.$$

Theorem 7.9.5

$\exp(x)$ is the unique function that satisfies:

- $\exp(0) = 1$
- $\exp(x + y) = \exp(x) \exp(y)$
- For any rational q , $\exp(qx) = \exp(x)^q$
- $\exp(x)$ is strictly increasing and bijective from $\mathbb{R}$ to $(0, \infty)$
- $\lim_{x \rightarrow \infty} \exp(x) = \infty$
- $\lim_{x \rightarrow -\infty} \exp(x) = 0$
- $\frac{d}{dx} \exp(x) = \exp(x)$

Definition 7.9.6 (Exponent)

Let $x > 0$ and $y \in \mathbb{R}$. Then,

$$x^y := \exp(y \ln(x)).$$

Thus,

$$\exp(x) = e^x.$$

8. Sequences of functions

8.1. Pointwise convergence

Definition 8.1.1 (Pointwise convergence)

Let $\{f_1, f_2, \dots\}$ be a sequence of functions. Let S be the set of all points x for which the sequence $\{f_1(x), f_2(x), \dots\}$ converges. Then, S is the domain of the limit function

$$f(x) := \lim_{n \rightarrow \infty} f_n(x).$$

$\{f_n\}$ **converges pointwise** to f on S .

8.2. Uniform convergence

Definition 8.2.1 (Uniform convergence)

Let $\{f_n\}$ be a sequence of functions. $\{f_n\}$ **converges uniformly** to f on S iff for every $\varepsilon > 0$ there exists N s.t. $n > N$ implies $|f(x) - f_n(x)| < \varepsilon$ for all $x \in S$.

Definition 8.2.2

Let $\{f_n\}$ be a sequence of functions on S . Let $M > 0$ be a constant so that $|f_n(x)| \leq M$ for all x and n . Then, $\{f_n\}$ is **uniformly bounded** by M on S .

Theorem 8.2.3

Let $f_n \rightarrow f$ uniformly on S and let each f_n be bounded. Then, f is bounded and the sequence $\{f_n\}$ is uniformly bounded on S .

Definition 8.2.4 (Uniform norm)

Let f be a bounded function on domain S . Then, define the **uniform norm** or **infinity norm** of f on S is

$$\|f\|_{\infty} = \|f\|_u := \sup\{|f(x)| : x \in S\}.$$

Theorem 8.2.5

A sequence of bounded functions f_n converges uniformly to f iff

$$\lim_{n \rightarrow \infty} \|f_n - f\|_{\infty} = 0.$$

Theorem 8.2.6

Let $\{f_n\}$ be a sequence of functions. Then there exists a function f s.t. $f_n \rightarrow f$ uniformly iff the sequence satisfies the **Cauchy condition**: for every $\varepsilon > 0$ there exists N s.t. if $m, n > N$, then $|f_n(x) - f_m(x)| < \varepsilon$ for all x .

Definition 8.2.7

A series converges uniformly iff its sequence of partial sums converges uniformly.

Theorem 8.2.8 (Weierstraß M-test)

Let M_n be a sequence of nonnegative numbers s.t. $|f_n(x)| \leq M_n$ for all x and n . Then, $\sum f_n$ converges uniformly if $\sum M_n$ converges.

8.3. Continuity, integrability, and differentiability of limit functions

See 41 Interchanging limits

Theorem 8.3.1 (Interchanging limits)

When $f_n \rightarrow f$ uniformly, then

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \lim_{x \rightarrow a} f_n(x) = \lim_{n \rightarrow \infty} f_n(a) = f(a)$$

Theorem 8.3.2

Let $\{f_n\}$ be a sequence of continuous functions $S \rightarrow \mathbb{R}$, where S is some domain. If the sequence converges uniformly to a function f , then f is also continuous.

Theorem 8.3.3

Let $\{f_n\}$ be a sequence of integrable functions on $[a, b]$ that converges uniformly to f . Then f is integrable, and

$$\int_a^b f = \lim_{n \rightarrow \infty} \int_a^b f_n.$$

Theorem 8.3.4

Let each term of $\{f_n\}$ be differentiable at each point of (a, b) and assume that these derivatives converge $f'_n \rightarrow g$ uniformly on (a, b) . Further, assume that there is at least one point $x_0 \in (a, b)$ s.t. $\{f_n(x_0)\}$ converges. Then,

- There is a function f s.t. $f_n \rightarrow f$ uniformly on (a, b) , and
- For each point in x in (a, b) , the derivative $f'(x)$ exists and equals $g(x)$.

Theorem 8.3.5

- The power series of a function converges uniformly upon its interval of convergence.
- Inside the interval of convergence, the power series represents a continuous function.
- The function is integrable, and the integral of the sum of the series is the sum of the integrals of the terms
- The derivatives of the power sums converge uniformly, and the partial sums of the series add to $f(x_0)$ at x_0 , so the derivative theorem's conditions are satisfied, and you can differentiate a power series term-by-term to obtain the derivative of the function to which it converges.

9. Picard's theorem

Definition 9.1 (Interior)

If I is an interval with endpoints a and b , I° is its **interior**, defined as

$$(a, b)$$

The interior of a finite union of intervals is the union of the interiors of those intervals.

For more complicated sets, we can use the definition from topology: if A is a set, then A° is the union of all open sets contained in A .

Definition 9.2 (First-order ordinary differential equation)

Let I, J in $\mathbb{R}$ be closed, bounded intervals. Let $F : I \times J \rightarrow \mathbb{R}$ be a function. Then,

$$f'(x) = F(x, f(x))$$

is a **first-order differential equation**. This is also written

$$y' = F(x, y).$$

If we know that

$$f(x_0) = y_0,$$

then we have an **initial-value problem** whose solution is the function $f : I \rightarrow J$ which satisfies both conditions:

$$f'(x) = F(x, f(x)) \quad f(x_0) = y_0.$$

Definition 9.3 (Lipschitz in the second variable)

$F : \mathbb{R}^2 \rightarrow \mathbb{R}$ is **Lipschitz in the second variable** iff there exists $L \in \mathbb{R}$ s.t.

$$|F(x, y) - F(x, z)| \leq L|y - z| \text{ for all } y, z \in J \text{ and } x \in I.$$

Theorem 9.4 (Picard's theorem)

Let $I, J \in \mathbb{R}$ be closed bounded intervals with $(x_0, y_0) \in I^\circ \times J^\circ$.

If $F : I \times J \rightarrow \mathbb{R}$ is continuous and Lipschitz in the second variable, then there exists $h > 0$ s.t. $[x_0 - h, x_0 + h] \subset I$ and a unique differentiable function $f : [x_0 - h, x_0 + h] \rightarrow J$ s.t.

$$f'(x) = F(x, f(x)) \quad f(x_0) = y_0.$$

In short, if F is continuous and Lipschitz in the second variable, the initial value problem has a unique solution on some interval around x_0 .

The unique solution can be constructed via the following procedure:

1. Define $f_0(x) := y_0$.
2. For each $k \in \mathbb{N}^+$, define the ***k*th Picard iterate**:

$$f_k(x) := y_0 + \int_{x_0}^x F(t, f_{k-1}(t)) dt$$

3. The sequence of Picard iterates converges uniformly to the solution f .

10. Construction of the reals

Let's assume that we have constructed the rationals. We must now construct the reals by taking Cauchy sequences of rationals, but we don't yet know that their (possibly irrational) limits exist, and there are multiple Cauchy sequences which converge to the same real which must be considered the same.

Definition 10.1 (Null sequence)

A sequence N of rational numbers is a **null sequence** iff its limit (in ε - N) is zero.

Definition 10.2 (Equivalence of Cauchy sequences)

Two Cauchy sequences are **equivalent** if the sequence of their differences is a null sequence:

$$\{a_n\} \equiv \{b_n\} \text{ iff } \{a_n - b_n\} \rightarrow 0.$$

Equivalence of Cauchy sequences is an equivalence relation.

Definition 10.3 (Real numbers)

$\mathbb{R}$ is the set of equivalence classes of Cauchy sequences of rational numbers under the above definition. Define $(\{a_n\})$ to be the equivalence class of $\{a_n\}$. Define

$$\begin{aligned} (\{a_n\}) + (\{b_n\}) &:= (\{a_n + b_n\}) \\ (\{a_n\})(\{b_n\}) &:= (\{a_n b_n\}). \end{aligned}$$

This satisfies the definition of the reals as the unique complete ordered field.

11. Lebesgue integrability condition

11.1. Topology of the reals

Definition 11.1.1 (Open set)

The following are **open**:

- The empty set
- Open intervals
- Unions of open sets
- Finite intersections of open sets
- The whole space

A set is **closed** if it is the complement of an open set.

The collection of closed sets is closed under finite union and arbitrary intersection.

Definition 11.1.2 (Continuity)

Let $f : X \rightarrow Y$ be a function from topological space X to topological space Y . Then, f is continuous iff $f^{-1}(A) \subset X \Leftrightarrow A$ is open in Y .

Theorem 11.1.3

The following are equivalent:

- f is continuous (under ε - δ)
- f is continuous under the above definition
- The inverse image of any closed set in $\mathbb{R}$ is closed in $\mathbb{R}$

Definition 11.1.4 (Oscillation)

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $S \subset \mathbb{R}$. The **oscillation** of f on S is defined as

$$\omega_f(S) := \sup\{f(S)\} - \inf\{f(S)\}.$$

If the supremum or infimum does not exist, it is defined as $+\infty$.

The **oscillation** of f at the point $x \in S$ is defined as

$$\lim_{\varepsilon \rightarrow 0} \omega_f(S \cap (x - \varepsilon, x + \varepsilon)).$$

Theorem 11.1.5

f is continuous at x_0 iff the oscillation of f at x_0 is zero.

Definition 11.1.6 (Open cover)

An **open cover** of X is a collection of open sets whose union contains all of X .

Definition 11.1.7 (Compactness)

A space where every open cover has a finite subcover.

Theorem 11.1.8

Subsets of $\mathbb{R}^n$ are compact iff they are closed and bounded.

11.2. Measure

Definition 11.2.1 (Measure)

A **measure** is a function that takes a subset S of the reals, and returns its “size” denoted $|S|$, following the following properties:

- $0 \leq |S| \leq \infty$.
- $|\emptyset| = 0$.
- $S' := \{x + a : x \in S\} \Rightarrow |S'| = |S|$.
- $|(0, 1)| = |[0, 1]| = 1$.
- **Countably additive**: If $\{S_i\}$ is a countable collection of disjoint sets, $|\cup S_i| = \sum_i |S_i|$.

Axiom of choice implies that no such function can exist. Axiom of determinacy implies such a function exists.

To avoid this, we define some nice sets as **measurable**, and leave the measure undefined for other sets.

11.2.1. Lebesgue outer measure

Definition 11.2.2 (Lebesgue outer measure)

Cover the set with a countable number of disjoint open intervals, and sum their totals. Take the infimum of all such coverings.

$$\lambda^*(S) := \inf \left\{ \sum (b_i - a_i) \mid \{(a_i, b_i)\}_{i \in \mathbb{Z}^+} \text{ is such a cover} \right\}.$$

Every set has such an outer measure, but it does not necessarily satisfy countable additivity. We define sets to be measurable if their outer measures satisfy said property.

A set has **measure 0** iff its outer measure is zero.

Theorem 11.2.3

- Every countable subset of $\mathbb{R}$ has measure 0.
- A subset of a measure 0 set also has measure 0.
- Any countable union of measure 0 sets has measure 0.

11.3. Lebesgue integrability condition

Theorem 11.3.1 (Lebesgue Integrability Condition)

$f : [a, b] \rightarrow \mathbb{R}$ is Riemann-integrable iff it is bounded and its set of discontinuities has measure 0.

Corollary 11.3.1.1

Let $f : [a, b] \rightarrow [c, d]$ be integrable and $g : [c, d] \rightarrow \mathbb{R}$ continuous. Then, $g \circ f$ is integrable.

12. Index

12.1. Headings

1. Sets, relations, functions, and order	1	5. Limits and continuity	9
1.1. Logic	1	5.1. Limits	9
1.2. Sets	1	5.2. Continuity	9
1.3. Ordered pair	1	5.3. Big Continuity Theorems	9
1.4. Relations	1	5.4. Uniform continuity	10
1.4.1. Equivalence relations	1	5.5. Lipschitz continuity	10
1.4.2. Functions (maps)	2	6. Derivatives	10
1.5. Sequence	2	6.1. Mean value theorem	11
1.6. Cardinality	2	6.2. Taylor series	11
1.7. Countability	3	7. Integrals	12
1.8. Axiom of choice	3	7.1. Partitions	12
2. Fields and order	3	7.2. Darboux integral	12
2.1. Equality	3	7.3. Riemann integral	12
2.2. Fields	3	7.4. Integrability	12
2.3. Order	4	7.5. Core properties of the integral	12
2.4. Ordered fields	4	7.6. Additional properties of the integral	13
2.5. Real numbers	4	7.7. Integrals of continuous functions	13
2.6. Extended reals	5	7.8. Fundamental theorem of calculus	13
2.7. Absolute values	5	7.9. Logarithm and exponential	13
2.8. Bounded function	5	8. Sequences of functions	14
2.9. Decimal representation	5	8.1. Pointwise convergence	14
3. Sequences	6	8.2. Uniform convergence	14
3.1. Subsequences	6	8.3. Continuity, integrability, and differentiability of limit functions	14
3.2. Limit properties and theorems	6	9. Picard's theorem	15
3.3. Limsup and liminf	7	10. Construction of the reals	15
3.4. Bolzano-Weierstras	7	11. Lebesgue integrability condition	16
3.5. Infinite limits	7	11.1. Topology of the reals	16
3.6. Cauchy sequences	7	11.2. Measure	16
4. Series	7	11.2.1. Lebesgue outer measure	16
4.1. Series	7	11.3. Lebesgue integrability condition	17
4.2. Basic convergence tests	8	12. Index	17
4.3. Absolute convergence	8	12.1. Headings	17
4.4. Ratio test and root test	8	12.2. Definitions	18
4.5. Alternating series	8	12.3. Theorems	18
4.6. Multiplying series	8	12.4. Axioms	19
4.7. Power series	9	12.5. Lemmata	19
		12.6. Corollary	19

12.2. Definitions

Definition 1.2.1	Power set	1
Definition 1.3.1	Ordered pair	1
Definition 1.4.1	Relation	1
Definition 1.4.4		1
Definition 1.4.5	Equivalence relation	1
Definition 1.4.6	Equivalence class	1
Definition 1.4.8	Partition	2
Definition 1.4.10	Function	2
Definition 1.4.11	Injective	2
Definition 1.4.12	Surjective	2
Definition 1.4.13	Bijjective	2
Definition 1.4.15		2
Definition 1.4.18	Induced map	2
Definition 1.5.1	Sequence	2
Definition 1.6.1	Cardinality	2
Definition 1.6.2		2
Definition 1.7.1	Countability	3
Definition 2.1.1	Equality	3
Definition 2.2.1	Field	3
Definition 2.3.1	Order	4
Definition 2.3.2	Upper bound	4
Definition 2.3.3	Supremum	4
Definition 2.3.4	Lower bound	4
Definition 2.3.5	Infimum	4
Definition 2.3.7	Dedekind completeness	4
Definition 2.4.1	Ordered field	4
Definition 2.5.6	Comensural	5
Definition 2.6.1	Extended reals	5
Definition 2.8.1	Bounded function	5
Definition 2.9.1	Decimal representation	5
Definition 3.1	Sequence	6
Definition 3.2	Limit of a sequence	6
Definition 3.4	Monotonic	6
Definition 3.1.1	Subsequence	6
Definition 3.3.1	Limsup and liminf	7
Definition 3.4.2	Open ball	7
Definition 3.4.3	Accumulation point	7
Definition 3.5.1	Diverging to infinity	7

Definition 3.5.3		7
Definition 3.6.1	Cauchy sequence	7
Definition 4.1.1	Series	7
Definition 4.3.2	Absolute convergence	8
Definition 4.6.1	Cauchy product	8
Definition 4.7.1	Power series	9
Definition 5.1.1	Accumulation point	9
Definition 5.1.3	Limit	9
Definition 5.2.1	Continuity	9
Definition 5.3.1	Bounded function	9
Definition 5.4.1	Uniform continuity	10
Definition 5.5.1	Lipschitz continuity	10
Definition 6.1	Derivative	10
Definition 6.3	One-sided derivative	10
Definition 6.4	Infinite derivatives	10
Definition 6.1.2	Critical point	11
Definition 6.2.2	Taylor series	11
Definition 7.1.1	Partition	12
Definition 7.1.2	Norm/mesh	12
Definition 7.1.3	Refinement	12
Definition 7.2.1	Darboux sum	12
Definition 7.2.2	Upper and lower Darboux integrals	12
Definition 7.2.3	Darboux integrability	12
Definition 7.3.1	Riemann sum	12
Definition 7.3.2	Riemann integrability	12
Definition 7.6.4		13
Definition 7.9.1	Natural logarithm	13
Definition 7.9.3	Logarithm	13
Definition 7.9.4	Natural exponent	14
Definition 7.9.6	Exponent	14
Definition 8.1.1	Pointwise convergence	14
Definition 8.2.1	Uniform convergence	14
Definition 8.2.2		14
Definition 8.2.4	Uniform norm	14
Definition 8.2.7		14
Definition 9.1	Interior	15
Definition 9.2	First-order ordinary differential equation	15

Definition 9.3	Lipschitz in the second variable	15
Definition 10.1	Null sequence	15
Definition 10.2	Equivalence of Cauchy sequences	15
Definition 10.3	Real numbers	16
Definition 11.1.1	Open set	16
Definition 11.1.2	Continuity	16
Definition 11.1.4	Oscillation	16
Definition 11.1.6	Open cover	16
Definition 11.1.7	Compactness	16
Definition 11.2.1	Measure	16
Definition 11.2.2	Lebesgue outer measure	16

12.3. Theorems

Theorem 1.1.1	De Morgan's Laws for statements	1
Theorem 1.1.2	De Morgan's Laws for set theory	1
Theorem 1.1.3		1
Theorem 1.3.2		1
Theorem 1.4.2	Relations can compose	1
Theorem 1.4.3		1
Theorem 1.4.9		2
Theorem 1.4.14		2
Theorem 1.4.17		2
Theorem 1.6.3	Schroeder-Bernstein Theorem	2
Theorem 1.6.4	Cantor's Theorem	2
Theorem 1.6.5		2
Theorem 1.7.2		3
Theorem 1.7.3		3
Theorem 1.8.2	Zermelo's Theorem	3
Theorem 1.8.3	Well-Ordering Property (Zermelo's Postulate)	3
Theorem 1.8.4	Hausdorff Maximum Principle	3
Theorem 1.8.5	Zorn's Lemma	3
Theorem 2.2.3	Zero is unique	3
Theorem 2.2.5	One is unique	3
Theorem 2.3.6		4
Theorem 2.4.3		4
Theorem 2.5.1		4
Theorem 2.5.2	Approximation Property	4
Theorem 2.5.3	Additive Property	4
Theorem 2.5.4	Comparison Property	4

Theorem 2.5.7	Archimedean property	5	Theorem 5.1.4		9	Theorem 8.2.3		14
Theorem 2.5.8	Density of the rationals / rationals between reals	5	Theorem 5.2.2		9	Theorem 8.2.5		14
Theorem 2.7.1	Corollaries of the Triangle Inequality	5	Theorem 5.2.3	Composition of continuous functions is continuous	9	Theorem 8.2.6		14
Theorem 2.8.2		5	Theorem 5.2.4		9	Theorem 8.2.8	Weierstraß M -test	14
Theorem 2.9.2		5	Theorem 5.3.3	Extreme Value Theorem	9	Theorem 8.3.1	Interchanging limits	14
Theorem 3.3		6	Theorem 5.3.5	Intermediate Value Theorem	9	Theorem 8.3.2		14
Theorem 3.5		6	Theorem 5.4.2		10	Theorem 8.3.3		14
Theorem 3.6		6	Theorem 5.4.4		10	Theorem 8.3.4		15
Theorem 3.1.2		6	Theorem 5.5.2		10	Theorem 8.3.5		15
Theorem 3.2.1		6	Theorem 6.2	Continuity theorem	10	Theorem 9.4	Picard's theorem	15
Theorem 3.2.2	Squeeze Theorem	6	Theorem 6.5	Sum, product, and quotient rules	10	Theorem 11.1.3		16
Theorem 3.2.3		6	Theorem 6.6	Chain rule	10	Theorem 11.1.5		16
Theorem 3.2.4		6	Theorem 6.1.1		11	Theorem 11.1.8		16
Theorem 3.2.5	Convergence of geometric series	6	Theorem 6.1.3	Rolle's theorem	11	Theorem 11.2.3		16
Theorem 3.2.6	Ratio Test	6	Theorem 6.1.4	Mean value theorem	11	Theorem 11.3.1	Lebesgue Integrability Condition	17
Theorem 3.2.7		7	Theorem 6.1.5	Cauchy's mean value theorem	11			
Theorem 3.3.2		7	Theorem 6.1.6		11	12.4. Axioms		
Theorem 3.3.3		7	Theorem 6.1.7		11	Axiom 1.8.1	Axiom of Choice	3
Theorem 3.3.4		7	Theorem 6.1.8	Darboux theorem	11	Axiom 2.2.2	Additive inverse axiom	3
Theorem 3.4.1	Bolzano-Weierstrass for Sequences	7	Theorem 6.2.1	Taylor's formula with the Lagrange form of the remainder	11	Axiom 2.2.4	Multiplicative inverse axiom	3
Theorem 3.4.4	Bolzano-Weierstrass theorem	7	Theorem 7.3.3		12	12.5. Lemmata		
Theorem 3.4.5	Cantor Intersection Theorem	7	Theorem 7.4.1		12	Lemma 1.4.7		2
Theorem 3.5.2		7	Theorem 7.4.2		12	Lemma 1.4.16		2
Theorem 3.6.2		7	Theorem 7.5.2	Additivity	12	Lemma 2.2.6		4
Theorem 3.6.3		7	Theorem 7.5.4	Linearity	13	Lemma 2.4.2		4
Theorem 4.2.1	n^{th} term test for divergence	8	Theorem 7.5.5	Monotonicity	13	Lemma 2.5.5		4
Theorem 4.2.2		8	Theorem 7.6.1		13	Lemma 2.9.3		5
Theorem 4.3.3		8	Theorem 7.6.2	Integral of a constant	13	Lemma 2.9.4		6
Theorem 4.3.4	Comparison test for positive series	8	Theorem 7.6.3	Integral of a step function	13	Lemma 4.3.1		8
Theorem 4.4.1	Ratio test	8	Theorem 7.7.3		13	Lemma 5.1.2		9
Theorem 4.4.2	Root test	8	Theorem 7.8.1	First Fundamental Theorem of Calculus	13	Lemma 5.3.2		9
Theorem 4.5.1	Alternating series test	8	Theorem 7.8.2	Second Fundamental Theorem of Calculus	13	Lemma 5.3.4		9
Theorem 4.5.2		8	Theorem 7.8.3	u-substitution	13	Lemma 5.4.3		10
Theorem 4.5.3	Alternating Series Rearrangement Theorem	8	Theorem 7.9.2		13	Lemma 7.5.1		12
Theorem 4.6.2	Mertens' theorem	8	Theorem 7.9.5		14	Lemma 7.5.3		12
Theorem 4.7.2		9				Lemma 7.7.1		13
						Lemma 7.7.2		13
						12.6. Corollary		

Corollary 1.7.1.1 3

Corollary 4.3.4.1 8

Corollary 6.1.1.1 11

Corollary 6.1.8.1 11

Corollary 11.3.1.1 17