

Math 122A Homework 7

Ethan Martirosyan

February 22, 2024

Bak Newmann Chapter 5

Problem 17

First, let us consider the case $n = 2$. Then, we note that

$$a_1 z_1 + a_2 z_2 = a_1 z_1 + (1 - a_1) z_2$$

which is in the convex set S by the definition of convex set. Next, we may suppose that $n \geq 2$ and that

$$a_1 z_1 + \cdots + a_n z_n$$

is in S whenever $a_i \geq 0$ for all i and $\sum a_i = 1$. We must show that

$$a_1 z_1 + \cdots + a_{n+1} z_{n+1}$$

is in S whenever $a_i \geq 0$ for all i and $\sum a_i = 1$. Notice that

$$a_1 z_1 + \cdots + a_{n+1} z_{n+1} = (1 - a_{n+1}) \left(\frac{a_1}{1 - a_{n+1}} z_1 + \cdots + \frac{a_n}{1 - a_{n+1}} z_n \right) + a_{n+1} z_{n+1}$$

By our induction hypothesis, we know that

$$\frac{a_1}{1 - a_{n+1}} z_1 + \cdots + \frac{a_n}{1 - a_{n+1}} z_n$$

is in S . By the definition of convex set, we may deduce that

$$(1 - a_{n+1}) \left(\frac{a_1}{1 - a_{n+1}} z_1 + \cdots + \frac{a_n}{1 - a_{n+1}} z_n \right) + a_{n+1} z_{n+1}$$

is in S , thus completing the proof.

Problem 19

Notice that an antiderivative of

$$P(z) = 1 + 2z + \cdots + nz^{n-1}$$

is

$$Q(z) = 1 + z + \cdots + z^n$$

Using the formula for geometric sums, we obtain

$$Q(z) = 1 + z + \cdots + z^n = \frac{z^{n+1} - 1}{z - 1}$$

Notice that the zeroes of the latter expression are

$$e^{\frac{2\pi ia}{n+1}}$$

where $1 \leq a \leq n$. That is, all the zeroes of $Q(z)$ lie on the unit circle. Thus the convex hull of these solutions must lie inside the unit circle. By the Gauss-Lucas Theorem, we may deduce that all the zeroes of $P(z)$ lie inside the unit circle.

Bak Newmann Chapter 6

Problem 1

Let $\beta = 1 + i$. Then, we have

$$\frac{1}{z} = \frac{1}{\beta + (z - \beta)}$$

Factoring out β yields

$$\frac{1}{\beta + (z - \beta)} = \frac{1}{\beta(1 + \frac{z - \beta}{\beta})} = \frac{1}{\beta} \cdot \frac{1}{1 - (-\frac{z - \beta}{\beta})}$$

Notice that

$$\frac{1}{\beta} \cdot \frac{1}{1 - (-\frac{z - \beta}{\beta})} = \frac{1}{\beta} \left(1 - \frac{z - \beta}{\beta} + \frac{(z - \beta)^2}{\beta^2} + \dots \right)$$

This is equal to

$$\frac{1}{\beta} \sum_{n=0}^{\infty} \frac{(-1)^n (z - \beta)^n}{\beta^n} = \sum_{n=0}^{\infty} \frac{(-1)^n (z - \beta)^n}{\beta^{n+1}}$$

Substituting $\beta = 1 + i$, we obtain

$$\frac{1}{z} = \sum_{n=0}^{\infty} \frac{(-1)^n (z - 1 - i)^n}{(1 + i)^{n+1}}$$

Problem 2

Notice that $1 - z - 2z^2 = (1 - 2z)(1 + z)$. Now, we may write

$$\frac{1}{1 - z - 2z^2} = \frac{A}{1 - 2z} + \frac{B}{1 + z}$$

where A and B are constants to be determined. Multiplying both sides by $1 - z - 2z^2$ yields

$$1 = A(1 + z) + B(1 - 2z)$$

so that

$$1 = A + Az + B - 2Bz = A + B + (A - 2B)z$$

Thus, we know that $A + B = 1$ and $A - 2B = 0$. Since $A = 2B$, we obtain $2B + B = 3B = 1$ so that $B = 1/3$ and $A = 1 - 1/3 = 2/3$. We have

$$\begin{aligned} \frac{1}{1 - z - 2z^2} &= \frac{2/3}{1 - 2z} + \frac{1/3}{1 + z} = \frac{2}{3} \sum_{n=0}^{\infty} 2^n z^n + \frac{1}{3} \sum_{n=0}^{\infty} (-1)^n z^n = \frac{1}{3} \left(\sum_{n=0}^{\infty} 2^{n+1} z^n + \sum_{n=0}^{\infty} (-1)^n z^n \right) \\ &= \frac{1}{3} \left(\sum_{n=0}^{\infty} (2^{n+1} + (-1)^n) z^n \right) = \sum_{n=0}^{\infty} \frac{2^{n+1} + (-1)^n}{3} z^n \end{aligned}$$

Problem 3

Differentiating the identity

$$1 + z + z^2 + \cdots = \frac{1}{1 - z}$$

yields

$$1 + 2z + \cdots = \frac{1}{(1 - z)^2}$$

Multiplying both sides by z , we obtain

$$z + 2z^2 + \cdots = \frac{z}{(1 - z)^2}$$

That is, we have established the following identity:

$$\sum n z^n = \frac{z}{(1 - z)^2}$$

Differentiating this identity yields

$$\sum n^2 z^{n-1} = \frac{1}{(1 - z)^2} + \frac{2z}{(1 - z)^3} = \frac{1 - z}{(1 - z)^3} + \frac{2z}{(1 - z)^3} = \frac{1 + z}{(1 - z)^3}$$

Multiplying by z , we obtain

$$\sum n^2 z^n = \frac{z(1 + z)}{(1 - z)^3}$$

Problem 4

For the sake of contradiction, suppose that for every positive integer n , we have

$$f(n) = \frac{1}{n+1} = \frac{1/n}{1/n+1}$$

Notice that the functions f and $g(z) = z/(z+1)$ agree at all points in the set $S = \{1/n : n \in \mathbb{N}\}$. This set has an accumulation point at 0. By Corollary 6.10 in Bak and Newman, we may deduce that $f = g$ throughout the region $|z| < 1$. By continuity, we know that $f = g$ on $|z| = 1$ except when $z = -1$. No matter how $f(-1)$ is defined, we know that f is not continuous at -1 since

$$\lim_{z \rightarrow -1} f(z) = \lim_{z \rightarrow -1} \frac{z}{z+1} = \infty$$

Thus f is not analytic at -1 , contrary to our assumptions about f .

Needham Chapter 9

Problem 1

Let us rewrite the integral

$$\int_0^{2\pi} \frac{dt}{1 + a^2 - 2a \cos t}$$

as a contour integral around the unit circle C by introducing the substitutions $\cos t = \frac{1}{2}(z + 1/z)$ and $dz = iz dt$, as the author does on page 437 of Visual Complex Analysis. Then, we find that

$$\int_0^{2\pi} \frac{dt}{1 + a^2 - 2a \cos t} = \oint_C \frac{-i(dz/z)}{1 + a^2 - a(z + 1/z)}$$

Manipulating the integrand, we find that

$$\begin{aligned} \frac{-i(dz/z)}{1 + a^2 - a(z + 1/z)} &= \frac{i dz}{-z - za^2 + az^2 + a} = \frac{i dz}{az^2 - za^2 - z + a} = \frac{i dz}{az(z - a) - (z - a)} \\ &= \frac{i dz}{(z - a)(az - 1)} \end{aligned}$$

Thus, we may conclude that

$$\int_0^{2\pi} \frac{dt}{1 + a^2 - 2a \cos t} = \oint_C \frac{i dz}{(z - a)(az - 1)}$$

Next, we claim that

$$\oint_C \frac{i dz}{(z - a)(az - 1)} = \frac{2\pi}{1 - a^2}$$

if $0 < a < 1$. Let $f(z) = i/(az - 1)$. Notice that f is analytic on the unit disk since its only singularity $z = \frac{1}{a}$ has modulus strictly greater than 1. Appealing to the Cauchy Integral Formula, we find that

$$\oint_C \frac{i dz}{(z - a)(az - 1)} = \oint_C \frac{f(z)}{z - a} dz = 2\pi i \cdot f(a) = 2\pi i \cdot \frac{i}{a^2 - 1} = \frac{-2\pi}{a^2 - 1} = \frac{2\pi}{1 - a^2}$$

We may deduce that

$$\int_0^{2\pi} \frac{dt}{1 + a^2 - 2a \cos t} = \frac{2\pi}{1 - a^2}$$

for $0 < a < 1$.

Problem 3

In Visual Complex Analysis, it is shown that

$$f^{(r)}(0) = \frac{r!}{2\pi i} \oint_C \frac{f(z)}{z^{r+1}} dz$$

for any entire function f , any nonnegative integer r , and any closed loop C . Let $f(z) = (1+z)^n$. Then, we have

$$f^{(r)}(0) = \frac{r!}{2\pi i} \oint_C \frac{(1+z)^n}{z^{r+1}} dz$$

By the definition of f , we know that

$$f'(z) = n(1+z)^{n-1}, \quad f''(z) = n(n-1)(1+z)^{n-2}, \quad f^{(3)}(z) = n(n-1)(n-2)(1+z)^{n-3}, \dots$$

By induction, we may deduce that

$$f^{(r)}(z) = n(n-1) \cdots (n-r+1)(1+z)^{n-r}$$

Setting $z = 0$, we obtain

$$f^{(r)}(0) = n(n-1) \cdots (n-r+1) = \frac{n!}{(n-r)!}$$

We have

$$\frac{n!}{(n-r)!} = f^{(r)}(0) = \frac{r!}{2\pi i} \oint_C \frac{(1+z)^n}{z^{r+1}} dz$$

Dividing both sides by $r!$ yields

$$\frac{n!}{r!(n-r)!} = \frac{1}{2\pi i} \oint_C \frac{(1+z)^n}{z^{r+1}} dz$$

or

$$\binom{n}{r} = \frac{1}{2\pi i} \oint_C \frac{(1+z)^n}{z^{r+1}} dz$$

Now, we may suppose that C is the unit circle, and we will use the above formula to prove that

$$\binom{2n}{n} \leq 4^n$$

We have

$$\binom{2n}{n} = \frac{1}{2\pi i} \oint_C \frac{(1+z)^{2n}}{z^{n+1}} dz \leq \frac{1}{2\pi} \oint_C \left| \frac{(1+z)^{2n}}{z^{n+1}} \right| dz$$

Notice that

$$|(1+z)^{2n}| = |1+z|^{2n} \leq 2^{2n} = 4^n$$

and

$$|z^{n+1}| = |z|^{n+1} = 1$$

for all z on the unit circle C . Furthermore, the circumference of C is 2π . Thus, using the M - L formula from Bak and Newman, we find that

$$\frac{1}{2\pi} \oint_C \left| \frac{(1+z)^{2n}}{z^{n+1}} \right| dz \leq \frac{1}{2\pi} \cdot 2\pi \cdot 4^n = 4^n$$

so that

$$\binom{2n}{n} \leq 4^n$$