

Certified analytical bounds for hypersonic reachable sets

Ethan Knox

Independent Researcher, Champaign, IL

PhD Research Summary Draft

September 29, 2026

Abstract

A method for computing *certified* reachable sets of a full six-degree-of-freedom hypersonic entry vehicle, reaching state dimensions where grid-based dynamic programming cannot, with a guarantee that is derived analytically and verified in exact arithmetic rather than produced by a numerical solver.

Open items

- Exact-Arithmetic Computational Pipeline:** The proposal demands a guarantee verified in exact arithmetic rather than produced by a numerical solver. Floating-point inaccuracies inherent to standard simulation tools compromise the analytical certificate. Must implement a custom, exact-arithmetic interval integrator (likely leveraging C/C++ or specialized rational arithmetic libraries). This pipeline must process the hybrid dynamics, manage the boundary-layer bubble transitions, and compute the intersection geometries deterministically to verify the final bounds. 3
- Liouville Equation for Differential Inclusions:** The current framework embeds the flight mechanics $\dot{x} = f_q(x, c, t)$ as a linear Liouville constraint on an occupation measure μ over $[0, T] \times \mathcal{X} \times U$. By bounding true non-polynomial aerodynamics within strict intervals, the vector field becomes a differential inclusion $\dot{x} \in \mathcal{F}(x, c, t)$. We must formulate a robust Liouville equation capable of acting on set-valued maps, and rigorously prove that the support of the resulting occupation measure, $\text{supp } \mu_T$, securely outer-approximates all possible trajectories generated by any vector field selected from the interval inclusion. 3
- Wrapping Effect Suppression in Boundary-Layer Bubbles:** Boundary-layer bubbles represent discrete atmospheric skip events as compact impulse sets $\mathcal{I}_j$ carried by internal scaling groups. Because our framework fixes time as the independent variable, integrating interval bounds through highly dynamic skip events will induce severe "wrapping effects" (artificial inflation of bounding boxes). Must develop advanced intersection geometries (zonotopes, Taylor models, etc.) integrated into the profile decomposition to prove geometries maintain tight corridor bounds through multiple skip events without the reachable set exploding prematurely. 4
- Bahouri-Gérard Under Interval Disturbances:** The framework relies on the Bahouri-Gérard almost-orthogonality argument to separate the multi-parameter scaling groups (atmospheric, attitude, and elastic). Under interval aerodynamics, these groups couple with an adversarial, bounded uncertainty. We must prove the non-interaction theorem for the defect measures while accounting for interval bounds, guaranteeing that the gap between the upper and lower aerodynamic bounds does not cause mass to leak between scaling groups during atmospheric transition, preserving the compact residual bound $O(\sqrt{\epsilon})$ 4
- Analytical Derivation of Corridor Constants:** The ultimate guarantee relies on deriving the residual corridor constant $C_{\mathcal{R}}$ and the maximum boundary-layer bubble count $K_{\max}(Q_{\max})$; The number of bubbles is bounded a priori by the integrated heat-load budget. We must then derive explicit, closed-form algebraic expressions for $C_{\mathcal{R}}$ and $K_{\max}$ strictly from the physical corridor bounds, initial conditions, and the interval widths of the aerodynamics, entirely without relying on numerical interior-point solvers. 5
- State Space Minimization and Kinematic Trade-offs:** The shift to exact interval arithmetic renders traditional Savageau-style polynomial adjunctions (e.g., $y = \sqrt{\rho/\rho_0}$, $\rho = 1/||\vec{r}||$) completely obsolete, as transcendentals are evaluated natively. We must formally strip away these algebraic embeddings and collapse the theoretical state space back to its minimal physical dimensions, which inherently tightens bounds by reducing susceptibility to the wrapping effect. The remaining open implementation task is determining if retaining purely kinematic adjunctions (such as quaternions to avoid Euler angle singularities) is computationally necessary to prevent interval bounding boxes from artificially inflating, or if the raw 13-state formulation is optimal for the pipeline's intersection geometries. 5

The problem

The reachable set of an entry vehicle (every state it can be driven to under admissible control) is what safety and threat analyses need bounded. Computing it exactly means solving a Hamilton–Jacobi–Isaacs equation on a grid, which is exponential in state dimension and capped in practice near four or five states [1, 2]. A rigid-body entry vehicle carries thirteen states, $x = (r, \theta, \phi, u, v, w, q_0, q_3, \omega_{1:3}) \in \mathbb{R}^{13}$, and twenty-five once ablation recession, integrated heat load and aerothermoelastic deformation are coupled [3].

So the exact grid-HJI method quits before the real problem starts.

The standard polynomial barrier with coefficients obtained from an interior-point solver has analytical *form* but inherits the solver’s accuracy [4], so it is not a certificate; a numerical solver’s failure modes are documented, and they include the loss of strict feasibility on which every rounding-based scheme depends [5, 6].

There are then two problems to solve, not one: dimension, and the provenance of the guarantee. The second is not a matter of care. Interval arithmetic is the setting in which a guarantee has meant, for forty years, an enclosure rather than an approximation, and the property that makes an enclosure valid at all—inclusion isotonicity—is exactly what a reimplementaion loses first; not every law of real arithmetic survives the move to intervals, which is the original form of the wrapping effect [7, 8].

Applied interval analysis supplies the machinery the pipeline would be built from, and it is deliberately conservative: certified set inversion, and a documented account of where the looseness enters [9].

Exact-Arithmetic Computational Pipeline: The proposal demands a guarantee verified in exact arithmetic rather than produced by a numerical solver. Floating-point inaccuracies inherent to standard simulation tools compromise the analytical certificate. Must implement a custom, exact-arithmetic interval integrator (likely leveraging C/C++ or specialized rational arithmetic libraries). This pipeline must process the hybrid dynamics, manage the boundary-layer bubble transitions, and compute the intersection geometries deterministically to verify the final bounds.

The approach

Embed the hybrid flight dynamics into a Banach space of occupation measures. The nonlinear flight mechanics $\dot{x} = f_q(x, c, t)$ become *linear*: the Liouville equation is a linear constraint on a measure μ over $[0, T] \times \mathcal{X} \times U$, and $\text{supp } \mu_T$ outer-approximates the reachable set. So far this is the occupation-measure program of Lasserre, Henrion and collaborators [10, 11]. The embedding buys linearity at the cost of uniqueness, and the way it is recovered is standard: a trajectory family satisfying the Liouville constraint lies in the convex hull of the generalized curves, so measures correspond to convex mixtures of trajectories rather than to one of them [12, 13]. The same relaxation is what admits the interval envelope, and that is the first thing that has to be built rather than borrowed. Because the true aerodynamics are transcendental, they are enclosed in strict intervals, and a vector field selected from an interval is a set-valued map; the resulting differential inclusion is the object the measure must describe, not the nominal field [14, 15]. Occupation-measure theory for inclusions of exactly this kind already exists and must be cited as prior art: it defines an occupational invariant measure without assuming solution uniqueness, which is the same move made here [16]. What it assumes is worth stating plainly, because it is where the entry problem is likely to fail: the vector fields must satisfy a linear growth condition, and the singular-perturbation results additionally require a uniform bound on the fast state along the whole ε -family [16, 17].

Liouville Equation for Differential Inclusions: The current framework embeds the flight mechanics $\dot{x} = f_q(x, c, t)$ as a linear Liouville constraint on an occupation measure μ over $[0, T] \times \mathcal{X} \times U$. By bounding true non-polynomial aerodynamics within strict intervals, the vector field becomes a differential inclusion $\dot{x} \in \mathcal{F}(x, c, t)$. We must formulate a robust Liouville equation capable of acting on set-valued maps, and rigorously prove that the support of the resulting occupation measure, $\text{supp } \mu_T$, securely outer-approximates all possible trajectories generated by any vector field selected from the interval inclusion.

The new content is what happens in the thin-atmosphere limit $\varepsilon_{\text{atm}} = H_s/R_e \rightarrow 0$. Rescaled by the phugoid-to-secular time ratio (which, as in the classical Chapman–Loh variables, enters as $\sqrt{\varepsilon_{\text{atm}}}$, not ε_{atm} [18, 19]), the occupation-measure family *is not tight*. Mass concentrates onto the low-dimensional equilibrium-glide slow manifold $\Lambda = L \cos \sigma / [m(g - V^2/r)] = 1$.

This is precisely Lions’ concentration alternative, read on the defect measure of the rescaled family [20]. The bubbles are concentration at a shrinking timescale, so the operative half of the principle is the dilation case rather than the translation case: it is the limit-case modification that applies [21]. Rather than restore compactness, we extract the profile decomposition the dichotomy hands us:

- a finite-dimensional *skeleton* $\mathcal{S}$ —the concentration component, the glide manifold as a graph over its retained coordinates, for which a rigorous characterization with quantitative accuracy estimates already exists [22];
- one *boundary-layer bubble* per skip event, the analogue of a concentrating profile: a compact impulse set $\mathcal{I}_j$ of dimension $\leq m$, carried by an internal scaling group at each atmospheric pass, with orthogonality read off the scale-and-core separation condition that governs concentrating profiles generally [23];

Wrapping Effect Suppression in Boundary-Layer Bubbles: Boundary-layer bubbles represent discrete atmospheric skip events as compact impulse sets $\mathcal{I}_j$ carried by internal scaling groups. Because our framework fixes time as the independent variable, integrating interval bounds through highly dynamic skip events will induce severe “wrapping effects” (artificial inflation of bounding boxes). Must develop advanced intersection geometries (zonotopes, Taylor models, etc.) integrated into the profile decomposition to prove geometries maintain tight corridor bounds through multiple skip events without the reachable set exploding prematurely.

- a *compact residual* with a quantitative bound, $O(\sqrt{\varepsilon})$ in the corridor metric, in a norm chosen so that the loss of compactness actually disappears rather than merely being hidden [24].

The number of bubbles is bounded *a priori*, $K \leq K_{\max}(Q_{\max})$, by the integrated heat-load budget (a trajectory can only afford so many skips [25, 26]). The wrapping effect to be suppressed here is not new: for linear systems it is already quantified, and the term is first order in the time step and exponential in the horizon times the matrix norm [27]. The multi-parameter separation of the three scaling groups (atmospheric, attitude, elastic) is the Bahouri–Gérard almost-orthogonality argument, transported to defect measures; it is the one genuinely new obligation and it sits on the critical path.

The argument worth transporting is the constructive one: splitting a bounded sequence into an oscillatory and a singular part at each scale forces those scales to be orthogonal, and iterating extracts a pairwise orthogonal family, so orthogonality is produced by the construction rather than assumed by it [28].

Bahouri-Gérard Under Interval Disturbances: The framework relies on the Bahouri-Gérard almost-orthogonality argument to separate the multi-parameter scaling groups (atmospheric, attitude, and elastic). Under interval aerodynamics, these groups couple with an adversarial, bounded uncertainty. We must prove the non-interaction theorem for the defect measures while accounting for interval bounds, guaranteeing that the gap between the upper and lower aerodynamic bounds does not cause mass to leak between scaling groups during atmospheric transition, preserving the compact residual bound $O(\sqrt{\epsilon})$.

The target statement is

$$\mathcal{R}_{\text{cert}}(w) \subseteq \underbrace{\left(\Phi_{\mathcal{S}}(\Theta) \right)}_{\text{skeleton}} \oplus \underbrace{\left(\bigoplus_{j=1}^K \mathcal{I}_j \right)}_{\text{bubbles}} \oplus \underbrace{\mathbb{B}(C_{\mathcal{R}}\sqrt{\epsilon})}_{\text{residual}},$$

uniformly over the admissible family generated by the uncertainty set, not along a nominal trajectory. The exact residual corridor constant $C_{\mathcal{R}}$ and the maximum bubble count K_{max} are derived as explicit, closed-form algebraic expressions directly from the physical corridor bounds and aerodynamic interval widths, completely independent of numerical optimization logic. Requiring the constants in closed form, rather than as the output of a solver, is what separates a certificate from a computation: exact rational certificate representations are known to exist, with a documented bit-complexity that grows with the number of variables [29, 30]. Because interval arithmetic evaluates transcendentals natively, the traditional algebraic adjunctions required to force the system into a polynomial format [31] are obsolete. The augmented state space is collapsed back to its minimal physical dimensions, retaining only the adjunctions genuinely necessary to avoid coordinate singularities.

Analytical Derivation of Corridor Constants: The ultimate guarantee relies on deriving the residual corridor constant $C_{\mathcal{R}}$ and the maximum boundary-layer bubble count $K_{\text{max}}(Q_{\text{max}})$; The number of bubbles is bounded a priori by the integrated heat-load budget. We must then derive explicit, closed-form algebraic expressions for $C_{\mathcal{R}}$ and K_{max} strictly from the physical corridor bounds, initial conditions, and the interval widths of the aerodynamics, entirely without relying on numerical interior-point solvers.

State Space Minimization and Kinematic Trade-offs: The shift to exact interval arithmetic renders traditional Savageau-style polynomial adjunctions (e.g., $y = \sqrt{\rho/\rho_0}$, $\rho = 1/||\vec{r}||$) completely obsolete, as transcendentals are evaluated natively. We must formally strip away these algebraic embeddings and collapse the theoretical state space back to its minimal physical dimensions, which inherently tightens bounds by reducing susceptibility to the wrapping effect. The remaining open implementation task is determining if retaining purely kinematic adjunctions (such as quaternions to avoid Euler angle singularities) is computationally necessary to prevent interval bounding boxes from artificially inflating, or if the raw 13-state formulation is optimal for the pipeline’s intersection geometries.

A Coupled Entry Dynamics and Polynomial Embedding

A.1 Coupled 6-DOF Entry Dynamics

The entry vehicle state comprises the coupled translational, rotational, thermal, ablative, and modal subsystems. Position is carried in Cartesian Earth-centred Earth-fixed (ECEF) components and attitude as a quaternion from the ECEF frame to the body frame. No angle is a state, so the dynamics have no coordinate singularities anywhere on the globe; geocentric latitude and longitude are read out from the position after integration.

Frames.

- $\mathcal{E}$: Earth-centred Earth-fixed (ECEF), rotating with the constant planetary rate $\vec{\omega}_E = \omega_E \hat{z}_{\mathcal{E}}$.
- $\mathcal{B}$: body frame, fixed at the instantaneous center of mass.

The attitude quaternion $\vec{q} = (q_0, q_1, q_2, q_3)^\top$ carries $\mathcal{E} \rightarrow \mathcal{B}$, with direction-cosine matrix

$$\mathbf{R} \equiv \mathbf{R}_{B/E}(\vec{q}) = \begin{pmatrix} q_0^2 + q_1^2 - q_2^2 - q_3^2 & 2(q_1q_2 + q_0q_3) & 2(q_1q_3 - q_0q_2) \\ 2(q_1q_2 - q_0q_3) & q_0^2 - q_1^2 + q_2^2 - q_3^2 & 2(q_2q_3 + q_0q_1) \\ 2(q_1q_3 + q_0q_2) & 2(q_2q_3 - q_0q_1) & q_0^2 - q_1^2 - q_2^2 + q_3^2 \end{pmatrix}, \quad (\text{A.1})$$

quadratic in $\vec{q}$.

Position Kinematics. With $\vec{r} = (r_1, r_2, r_3)^\top$ the ECEF position and $\vec{V} = (u, v, w)^\top$ the planet-relative velocity resolved in $\mathcal{B}$, the kinematic equation is:

$$\dot{\vec{r}} = \mathbf{R}^\top \vec{V}. \quad (\text{A.2})$$

Translational Dynamics. Newton's law holds in the inertial frame; transporting the planet-relative velocity through $\mathcal{E}$ into $\mathcal{B}$, with inertial body rates $\vec{\omega} = (p, q, r_b)^\top$, mass m , aerodynamic force $\vec{F}_a$, thrust $\vec{F}_t$, and the planetary rate resolved in the body frame $\vec{\omega}_E^{\mathcal{B}} = \mathbf{R} \omega_E \hat{z}_{\mathcal{E}}$, the translational dynamical equations are:

$$\dot{\vec{V}}|_{\mathcal{B}} = \frac{1}{m}(\vec{F}_a + \vec{F}_t) + \mathbf{R} \vec{g}(\vec{r}) - (\vec{\omega} + \vec{\omega}_E^{\mathcal{B}}) \times \vec{V} - \vec{\omega}_E^{\mathcal{B}} \times (\vec{\omega}_E^{\mathcal{B}} \times \mathbf{R} \vec{r}). \quad (\text{A.3})$$

Gravity, including the J_2 oblateness term, in ECEF components:

$$\vec{g}(\vec{r}) = -\frac{\mu}{\|\vec{r}\|^3} \left[\left(1 + \frac{3}{2} J_2 \frac{R_e^2}{\|\vec{r}\|^2} \left(1 - 5 \frac{r_3^2}{\|\vec{r}\|^2} \right) \right) \vec{r} + 3 J_2 \frac{R_e^2}{\|\vec{r}\|^2} r_3 \hat{z}_{\mathcal{E}} \right]. \quad (\text{A.4})$$

Remark A.1. This is the gradient of the J_2 potential $\frac{\mu}{\|\vec{r}\|} [1 - \frac{1}{2} J_2 (R_e / \|\vec{r}\|)^2 (3r_3^2 / \|\vec{r}\|^2 - 1)]$; its radial and northward components reproduce the spherical form, with $r_3 / \|\vec{r}\| = \sin \phi$.

Rotational Dynamics / Euler's Equations. With angular momentum $\vec{h} = \mathbf{I} \vec{\omega}$, combined aerodynamic / reaction control system (RCS) moments $\vec{M} = \vec{M}_a + \vec{M}_{\text{RCS}}$, and the inertia tensor $\mathbf{I} = \mathbf{I}(s, \eta)$ depending on recession depth s and modal amplitudes η (so that $\mathbf{I} \vec{\omega}$ is the variable-inertia term), the rotational dynamical equations are:

$$\mathbf{I} \dot{\vec{\omega}} = \vec{M} - \vec{\omega} \times (\mathbf{I} \vec{\omega}) - \dot{\mathbf{I}} \vec{\omega}. \quad (\text{A.5})$$

Quaternion Kinematics. The quaternion is driven by the body rate *relative to* $\mathcal{E}$, $\vec{\omega}_{B/E} = \vec{\omega} - \vec{\omega}_E^{\mathcal{B}}$. For $\vec{a} = (a_1, a_2, a_3)^\top$,

$$\dot{\vec{q}} = \frac{1}{2} \Omega(\vec{\omega} - \mathbf{R} \omega_E \hat{z}_{\mathcal{E}}) \vec{q}, \quad \Omega(\vec{a}) = \begin{pmatrix} 0 & -a_1 & -a_2 & -a_3 \\ a_1 & 0 & a_3 & -a_2 \\ a_2 & -a_3 & 0 & a_1 \\ a_3 & a_2 & -a_1 & 0 \end{pmatrix}. \quad (\text{A.6})$$

Remark A.2. The rate is **polynomial** — bilinear in $(\vec{\omega}, \vec{q})$ and cubic in $\vec{q}$ through the Earth-rate term. With this Ω and $\mathbf{R}$ of Eq. (A.1), $\dot{\mathbf{R}} = -[\vec{\omega}_{B/E} \times] \mathbf{R}$ holds identically, and $\Omega(\vec{a})$ is skew for every $\vec{a}$, so $|\vec{q}|$ is preserved.

Thermal. With the density-square-root embedding $y = \sqrt{\rho/\rho_0}$ and $V = \sqrt{u^2 + v^2 + w^2}$, the modulus lift $\dot{\chi} = c \chi \dot{T}_w$ replaces T_w with a polynomial via Varma–Savageau–Voit recasting [31]. Stagnation-point heating (Sutton–Graves [32], with nose radius R_n , sea-level density ρ_0) equations are then given by:

$$\dot{Q}_{\text{tot}} = \dot{Q} = k_Q \sqrt{\rho_0/R_n} y V^3, \quad \dot{T}_w = \frac{\dot{Q} - (T_w - T_\infty)/\tau_{\text{cond}}}{C_w}. \quad (\text{A.7})$$

The correlation is used inside its stated validity range and not extrapolated past it: axisymmetric blunt bodies, gases in chemical equilibrium, enthalpy 2.3 to 116.2 MJ/kg, pressure 0.001 to 100 atm, and wall temperature at one of two tabulated values rather than a range[32, 33]. The chemical-equilibrium clause is the one most likely to fail, at the high-altitude end, which is where the nonequilibrium and rarefied treatments take over[34]. The density profile itself should come from a whole-atmosphere model rather than a spliced pair, since a splice would put a discontinuity in the vector field and defeat the polynomial recast[35].

Ablation. With recession rate coupled to heat flux and mass loss from char-layer blow-off, the ablative equations are given by:

$$\dot{s} = \alpha_s \dot{Q}, \quad \dot{m} = -\dot{m}_{\text{abl}}. \quad (\text{A.8})$$

Recession is a *state* with its own differential equation rather than a post-processed output because the interface condition requires the heat input to equal the heat conducted into the solid plus the heat absorbed in melting; the transformation that maps a receding boundary onto a fixed one is the classical one, and it costs only extra convective terms in products of states already carried[36]. A decoupled correlation for the recession rate is not obviously conservative: the surface energy balance couples to a multicomponent boundary layer with unequal diffusion coefficients, and the adequacy of a decoupled treatment depends on the material–environment pair[37].

Modal/Structural Elasticity. With piston-theory forcing and temperature-dependent natural frequency, per mode i our modal equations are:

$$\ddot{\eta}_i + 2\zeta_i \omega_i(T) \dot{\eta}_i + \omega_i(T)^2 \eta_i = Q_i(x, \eta, \dot{\eta}). \quad (\text{A.9})$$

Piston theory is a double expansion in thickness ratio and inverse Mach number valid when the contributions of order δ/M^3 and δ^2/M^2 are negligible, which is where its applicability conditions are stated and where they must be checked rather than assumed[38]. That choice is load-bearing

for the cost in SOS degree: the first-order point-function relation is affine in the normal component of fluid velocity, hence affine in the modal rates and free in the Gram table, whereas each further order costs one degree in the downwash and multiplies the modal block through. Carrying first order exactly and enveloping the higher-order residuals against the paper’s own smallness criterion is therefore the trade to state explicitly. The pressure band over which the aerodynamic law is the one being certified is likewise a checkable, state-dependent condition, and belongs in the constraint set rather than in an assumption[39].

A.2 Polynomial Embedding via Adjunction

A priori, the dynamics contain transcendental and rational terms. We repair each of these by adjoining one variable and one polynomial defining constraint, so the entire augmented system is polynomial in the augmented state:

1. **Transcendental Incidence:** $\alpha = \text{atan2}(w, u)$, $\beta = \arcsin(v/V)$ are inverse-trigonometric functions of body rates, we replace them with polynomial direction-cosine ratios $(u/V, v/V, w/V)$.
2. **Square-Root Speed:** Since $V = \sqrt{u^2 + v^2 + w^2}$ carries a square root, we adjoin V with the velocity magnitude constraint $V^2 = u^2 + v^2 + w^2$.
3. **Exponential Atmosphere:** Since $\rho = \rho_0 e^{-h/H_s}$ with $h = \|\vec{r}\| - R_e$ appears in $\bar{q} = \frac{1}{2}\rho V^2$, we adjoin $y = \sqrt{\rho/\rho_0}$ with rate $\dot{y} = -\frac{y}{2H_s} \frac{d}{dt}\|\vec{r}\|$, where the radial rate $\frac{d}{dt}\|\vec{r}\| = \varrho(\mathbf{R}\vec{r}) \cdot \vec{V}$ is polynomial once ϱ (item 4) is adjoined.
4. **Reciprocal Radius:** Gravity carries $\|\vec{r}\|^{-3}$, $\|\vec{r}\|^{-5}$, $\|\vec{r}\|^{-7}$, so we adjoin $\varrho = 1/\|\vec{r}\|$ with $\varrho^2 \|\vec{r}\|^2 = 1$, $\varrho > 0$, and rate $\dot{\varrho} = -\varrho^3(\mathbf{R}\vec{r}) \cdot \vec{V}$. Gravity is then polynomial:

$$\vec{g} = -\mu\varrho^3 \left[\left(1 + \frac{3}{2}J_2R_e^2\varrho^2(1 - 5r_3^2\varrho^2)\right)\vec{r} + 3J_2R_e^2\varrho^2 r_3 \hat{z}_{\mathcal{E}} \right]. \quad (\text{A.10})$$

No trigonometric adjunctions are needed: no angle is a state, and latitude dependence enters only through $r_3\varrho = \sin\phi$.

5. **Quaternion Norm:** $|\vec{q}|^2 = 1$ is already a polynomial invariant (skew-symmetry of Ω preserves it), no adjunction necessary.
6. **Heat / Load-Factor Square Roots:** $\dot{Q} \propto yV^3$ is already polynomial; however, $n = \|\vec{F}_a\|/(mg_0)$ requires squaring to $(\bar{q}S)^2(C_X^2 + C_Y^2 + C_Z^2) \leq (n_{\max}mg_0)^2$.

The Augmented State. Collecting all coordinates:

$$\tilde{x} = \left(\underbrace{r_1, r_2, r_3, u, v, w, q_0, q_1, q_2, q_3, p, q, r_b}_{13, \text{ core}}, \underbrace{y, V, \varrho, Q_{\text{tot}}, m, s, \chi}_{7, \text{ adjoined}}, \underbrace{\eta, \dot{\eta}}_{2n_{\eta}, \text{ modal}} \right). \quad (\text{A.11})$$

References

- [1] S. Bansal, M. Chen, S. Herbert, and C. J. Tomlin. “Hamilton-Jacobi Reachability: A Brief Overview and Recent Advances”. In: *56th IEEE Conference on Decision and Control (CDC)*. 2017, pp. 2242–2253. DOI: [10.1109/CDC.2017.8263977](https://doi.org/10.1109/CDC.2017.8263977).
- [2] I. M. Mitchell, A. M. Bayen, and C. J. Tomlin. “A Time-Dependent Hamilton-Jacobi Formulation of Reachable Sets for Continuous Dynamic Games”. In: *IEEE Transactions on Automatic Control* 50.7 (2005), pp. 947–957. DOI: [10.1109/TAC.2005.851439](https://doi.org/10.1109/TAC.2005.851439).

- [3] J. J. McNamara and P. P. Friedmann. “Aeroelastic and Aerothermoelastic Analysis in Hypersonic Flow: Past, Present, and Future”. In: *AIAA Journal* 49.6 (2011), pp. 1089–1122. DOI: [10.2514/1.J050882](https://doi.org/10.2514/1.J050882).
- [4] S. Prajna, A. Jadbabaie, and G. J. Pappas. “A Framework for Worst-Case and Stochastic Safety Verification Using Barrier Certificates”. In: *IEEE Transactions on Automatic Control* 52.8 (2007), pp. 1415–1428. DOI: [10.1109/TAC.2007.902736](https://doi.org/10.1109/TAC.2007.902736).
- [5] P. Roux, Y.-L. Voronin, and S. Sankaranarayanan. “Validating Numerical Semidefinite Programming Solvers for Polynomial Invariants”. In: *Formal Methods in System Design* 53.2 (2018). Published online 14 October 2017, pp. 286–312. DOI: [10.1007/s10703-017-0302-y](https://doi.org/10.1007/s10703-017-0302-y).
- [6] H. Peyrl and P. A. Parrilo. “Computing Sum of Squares Decompositions with Rational Coefficients”. In: *Theoretical Computer Science* 409.2 (2008), pp. 269–281. DOI: [10.1016/j.tcs.2008.09.025](https://doi.org/10.1016/j.tcs.2008.09.025).
- [7] R. E. Moore, R. B. Kearfott, and M. J. Cloud. *Introduction to Interval Analysis*. Vol. 110. Other Titles in Applied Mathematics. Read off the copyright page of the front matter in references/, which is split into fm.pdf and ch1-ch11.pdf plus appendices. ix+213 pages. Cross-checked against the SIAM record. Philadelphia: Society for Industrial and Applied Mathematics, 2009. ISBN: 978-0-89871-669-6. DOI: [10.1137/1.9780898717716](https://doi.org/10.1137/1.9780898717716).
- [8] N. S. Nedialkov, K. R. Jackson, and G. F. Corliss. “Validated Solutions of Initial Value Problems for Ordinary Differential Equations”. In: *Applied Mathematics and Computation* 105.1 (1999), pp. 21–68. DOI: [10.1016/S0096-3003\(98\)10083-8](https://doi.org/10.1016/S0096-3003(98)10083-8).
- [9] L. Jaulin, M. Kieffer, O. Didrit, and E. Walter. *Applied Interval Analysis: With Examples in Parameter and State Estimation, Robust Control and Robotics*. Read off the copyright page of the copy in references/ (LoC 2001020164, QA297.75 .A664, 382 pages in this printing). The copy is an OCR’d paper capture, so the text layer is thin but the front matter is legible. London: Springer-Verlag, 2001. ISBN: 978-1-4471-1067-5. DOI: [10.1007/978-1-4471-0249-6](https://doi.org/10.1007/978-1-4471-0249-6).
- [10] J.-B. Lasserre, D. Henrion, C. Prieur, and E. Trélat. “Nonlinear Optimal Control via Occupation Measures and LMI Relaxations”. In: *SIAM Journal on Control and Optimization* 47.4 (2008), pp. 1643–1666.
- [11] R. B. Vinter. “Convex Duality and Nonlinear Optimal Control”. In: *SIAM Journal on Control and Optimization* 31.2 (1993), pp. 518–538.
- [12] D. Henrion and M. Korda. “Convex Computation of the Region of Attraction of Polynomial Control Systems”. In: *IEEE Transactions on Automatic Control* 59.2 (2014), pp. 297–312. DOI: [10.1109/TAC.2013.2283095](https://doi.org/10.1109/TAC.2013.2283095).
- [13] J. M. Ball. “A Version of the Fundamental Theorem for Young Measures”. In: *PDEs and Continuum Models of Phase Transitions*. Vol. 344. Lecture Notes in Physics. VERIFY the booktitle and editors against the published record; the page range 207–215 is confirmed from the file. Berlin: Springer, 1989, pp. 207–215.
- [14] A. F. Filippov. *Differential Equations with Discontinuous Righthand Sides*. Ed. by F. M. Arscott. Mathematics and Its Applications (Soviet Series). Translated from the Russian. Read off the copyright page of the copy in references/ (LoC 88-4531, QA374.F4613 1988, 325 pages) and cross-checked against the Springer record. The 2010 softcover reprint carries the same DOI. Dordrecht: Kluwer Academic Publishers, 1988. ISBN: 902772699X. DOI: [10.1007/978-94-015-7793-9](https://doi.org/10.1007/978-94-015-7793-9).

- [15] J.-P. Aubin and A. Cellina. *Differential Inclusions: Set-Valued Maps and Viability Theory*. Vol. 264. Grundlehren der mathematischen Wissenschaften. Read off the title page and the Internet Archive metadata of the copy in references/ (366 pages); cross-checked against the Springer and WorldCat records. Berlin: Springer-Verlag, 1984. DOI: [10.1007/978-3-642-69512-4](https://doi.org/10.1007/978-3-642-69512-4).
- [16] Z. Artstein. “Invariant Measures of Differential Inclusions Applied to Singular Perturbations”. In: *Journal of Differential Equations* 152.2 (1999). Received 10 September 1997; revised 9 July 1998. RESOLVES the 1999/2000 dating question: this is the 1999 paper, and artstein2000value is the separate 2000 one, pp. 289–307. DOI: [10.1006/jdeq.1998.3527](https://doi.org/10.1006/jdeq.1998.3527).
- [17] Z. Artstein and V. Gaitsgory. “The Value Function of Singularly Perturbed Control Systems”. In: *Applied Mathematics and Optimization* 41.3 (2000), pp. 425–445. DOI: [10.1007/s002459910022](https://doi.org/10.1007/s002459910022).
- [18] D. R. Chapman. *An Approximate Analytical Method for Studying Entry into Planetary Atmospheres*. Tech. rep. Technical Note 4276. Later reissued as NASA TR R-11. 103 pages. Ames Aeronautical Laboratory, Moffett Field, California: National Advisory Committee for Aeronautics, May 1958.
- [19] W. H. T. Loh. “Re-entry and Planetary Entry Physics and Technology. Dynamics, Physics, Radiation, Heat Transfer and Ablation”. In: *Re-entry and Planetary Entry Physics and Technology*. Ed. by W. H. T. Loh. Vol. 2. Applied Physics and Engineering: An International Series. Read off the title and copyright pages of the copy in references/ (LoC 68-18103, Springer title no. 3892, 491 pages in this scan). The volume is edited AND authored by Loh. VERIFY the series/volume assignment against the published record before submission: the scan’s cover reads “Volume 2” while its title page reads “I”, which is the usual two-volume set ambiguity, and the chapter field above is a placeholder for whatever the volume’s own pagination settles. New York: Springer-Verlag, 1968. Chap. Volume 1. ISBN: 978-3-642-48929-7. DOI: [10.1007/978-3-642-48927-3](https://doi.org/10.1007/978-3-642-48927-3).
- [20] P.-L. Lions. “The Concentration-Compactness Principle in the Calculus of Variations. The Locally Compact Case, Part 1”. In: *Annales de l’Institut Henri Poincaré, Analyse Non Linéaire* 1.2 (1984), pp. 109–145.
- [21] P.-L. Lions. “The Concentration-Compactness Principle in the Calculus of Variations. The Limit Case, Part 1”. In: *Revista Matemática Iberoamericana* 1.1 (1985). The copy in references/ is a page scan with no text layer, pp. 145–201.
- [22] P. Lu. “Asymptotic Analysis of Quasi-Equilibrium Glide in Lifting Entry Flight”. In: *Journal of Guidance, Control, and Dynamics* 29.3 (2006), pp. 662–670. DOI: [10.2514/1.15384](https://doi.org/10.2514/1.15384).
- [23] P. Gérard. “Description du Défaut de Compacité de l’Injection de Sobolev”. In: *ESAIM: Control, Optimisation and Calculus of Variations* 3 (1998). In French. The text layer of the copy in references/ is badly mangled by the OCR; the page images are clean, pp. 213–233. DOI: [10.1051/cocv:1998107](https://doi.org/10.1051/cocv:1998107).
- [24] S. Solimini. “A Note on Compactness-Type Properties with Respect to Lorentz Norms of Bounded Subsets of a Sobolev Space”. In: *Annales de l’Institut Henri Poincaré, Analyse Non Linéaire* 12.3 (1995), pp. 319–337.

- [25] Y.-Y. Shi, L. Pottsepp, and M. C. Eckstein. “A Matched Asymptotic Solution for Skipping Entry into Planetary Atmosphere”. In: *AIAA Journal* 9.4 (Apr. 1971). A Technical Note, not a full paper – under two and a half pages. RESOLVES the earlier "VERIFY everything". Authors are with McDonnell Douglas Astronautics Company, Western Division, Huntington Beach, California, pp. 736–738.
- [26] J. Eggers Alfred J., H. J. Allen, and S. E. Neice. *A Comparative Analysis of the Performance of Long-Range Hypervelocity Vehicles*. Tech. rep. Report 1382. Ames Aeronautical Laboratory, Moffett Field, California: National Advisory Committee for Aeronautics, 1958.
- [27] C. Le Guernic and A. Girard. “Reachability Analysis of Linear Systems Using Support Functions”. In: *Nonlinear Analysis: Hybrid Systems* 4.2 (2010). RESOLVES the placeholder leguernic2009reachability: the venue is Nonlinear Analysis: Hybrid Systems as suspected, but the year is 2010, not 2009 (received 24 October 2008, accepted 11 March 2009), pp. 250–262. DOI: [10.1016/j.nahs.2009.03.002](https://doi.org/10.1016/j.nahs.2009.03.002).
- [28] H. Bahouri and P. Gérard. “High Frequency Approximation of Solutions to Critical Nonlinear Wave Equations”. In: *American Journal of Mathematics* 121.1 (1999), pp. 131–175. DOI: [10.1353/ajm.1999.0001](https://doi.org/10.1353/ajm.1999.0001).
- [29] E. L. Kaltofen, B. Li, Z. Yang, and L. Zhi. “Exact Certification in Global Polynomial Optimization via Sums-of-Squares of Rational Functions with Rational Coefficients”. In: *Journal of Symbolic Computation* 47.1 (2012). Received 6 January 2009; accepted 27 July 2011, pp. 1–15. DOI: [10.1016/j.jsc.2011.08.002](https://doi.org/10.1016/j.jsc.2011.08.002).
- [30] V. Magron and M. Safey El Din. “On Exact Reznick, Hilbert-Artin and Putinar’s Representations”. In: *Journal of Symbolic Computation* 107 (2021). SUPERSEDES the placeholder key magron2019exact and its guessed title "On Exact Polya and Putinar’s Representations", which is a DIFFERENT paper by the same authors. Received 17 July 2019, revised 25 November 2020, accepted 29 March 2021, pp. 221–250. DOI: [10.1016/j.jsc.2021.03.003](https://doi.org/10.1016/j.jsc.2021.03.003).
- [31] M. A. Savageau and E. O. Voit. “Recasting Nonlinear Differential Equations as S-Systems: A Canonical Nonlinear Form”. In: *Mathematical Biosciences* 87.1 (1987). Received 6 May 1987; revised 28 July 1987, pp. 83–115. DOI: [10.1016/0025-5564\(87\)90035-6](https://doi.org/10.1016/0025-5564(87)90035-6).
- [32] K. Sutton and J. Graves Randolph A. *A General Stagnation-Point Convective-Heating Equation for Arbitrary Gas Mixtures*. Tech. rep. NASA TR R-376. Langley report number L-7885; work unit 117-07-01-01. Langley Research Center, Hampton, Virginia: National Aeronautics and Space Administration, Nov. 1971.
- [33] J. A. Fay and F. R. Riddell. “Theory of Stagnation Point Heat Transfer in Dissociated Air”. In: *Journal of the Aeronautical Sciences* 25.2 (Feb. 1958). Received 14 May 1957. Avco Research Laboratory; sponsored by the Ballistic Missile Division, ARDC. VERIFY whether the published range runs to p85 or continues on a later page, as was common in this journal, pp. 73–85. DOI: [10.2514/8.7517](https://doi.org/10.2514/8.7517).
- [34] P. A. Gnoffo. “Planetary-Entry Gas Dynamics”. In: *Annual Review of Fluid Mechanics* 31 (1999), pp. 459–494. DOI: [10.1146/annurev.fluid.31.1.459](https://doi.org/10.1146/annurev.fluid.31.1.459).
- [35] J. T. Emmert, D. P. Drob, J. M. Picone, D. E. Siskind, J. Jones M., M. G. Mlynczak, P. F. Bernath, X. Chu, E. Doornbos, B. Funke, L. P. Goncharenko, M. E. Hervig, M. J. Schwartz, P. E. Sheese, F. Vargas, B. P. Williams, and T. Yuan. “NRLMSIS 2.0: A Whole-Atmosphere Empirical Model of Temperature and Neutral Species Densities”. In: *Earth and Space Science* 8.3 (2021). Seventeen authors; the full list is given above rather than et al. because the earlier entry flagged "VERIFY author list", e2020EA001321. DOI: [10.1029/2020EA001321](https://doi.org/10.1029/2020EA001321).

- [36] H. G. Landau. “Heat Conduction in a Melting Solid”. In: *Quarterly of Applied Mathematics* 8.1 (1950). Ballistic Research Laboratories, Aberdeen Proving Ground. SUPERSEDES the bare placeholder key landau1950, pp. 81–94.
- [37] R. M. Kendall, R. A. Rindal, and E. P. Bartlett. “A Multicomponent Boundary Layer Chemically Coupled to an Ablating Surface”. In: *AIAA Journal* 5.6 (June 1967). SUPERSEDES the placeholder kendall1968analytical, which had this as a 1968 Aerotherm technical report. It is a 1967 AIAA Journal article; the authors were with Aerotherm Corporation, Palo Alto, pp. 1063–1071. DOI: [10.2514/3.4138](https://doi.org/10.2514/3.4138).
- [38] H. Ashley and G. Zartarian. “Piston Theory—A New Aerodynamic Tool for the Aeroelastician”. In: *Journal of the Aeronautical Sciences* 23.12 (1956), pp. 1109–1118. DOI: [10.2514/8.3766](https://doi.org/10.2514/8.3766).
- [39] M. J. Lighthill. “Oscillating Airfoils at High Mach Number”. In: *Journal of the Aeronautical Sciences* 20.6 (1953), pp. 402–406. DOI: [10.2514/8.2657](https://doi.org/10.2514/8.2657).