

Certified analytical bounds for hypersonic reachable sets

A two-page summary of a preliminary PhD research proposal

Ethan Knox · Champaign, IL · ethank5149@gmail.com · October 7, 2026

Full proposal: research.ekdynamics.science/proposal.pdf

The result. This is a problem in nonlinear analysis: a singularly perturbed family of differential inclusions in which compactness fails along the time axis, because the motion carrying the important state transitions occupies a vanishing time. It therefore disappears from the ordinary limits of the family while continuing to decide where the trajectories go. For a model that keeps exactly that phenomenon I can prove

$$\mathcal{R}_\varepsilon(T) \subseteq \underbrace{\mathcal{R}_\mathcal{H}([T - \varrho, T + \varrho])}_{\text{skeleton, bubbles and glide}} \oplus \underbrace{\mathbb{B}(\varrho)}_{\text{residual}}, \quad \varrho = C\varepsilon^{1/3},$$

with C explicit in the data and the inclusion uniform over the controls, over the initial set and over a bounded family of admissible fields. The proof is four pages and elementary; the statement, the hypotheses and the proof are in the proposal. What follows says what the object is, why the decomposition is the natural one, and what remains open.

The problem. The structural phenomenon is concentration, and the language for it is a profile decomposition: a low-dimensional *skeleton*, one *bubble* per pass, and a *residual*. In the theory of nonlinear dispersive equations such a decomposition comes from iterating the dichotomy of the concentration-compactness principle [1, 2], and what it delivers is structure without a constant — by contradiction. The proposal is that here the structure can be had quantitatively, because the family carries Lyapunov functionals: the dissipation is monotone with total variation bounded uniformly in the parameter, so the number of profiles above any threshold is bounded in advance rather than extracted. That is where I expect the contribution to the analysis to lie.

The example that supplies the problem is hypersonic atmospheric entry, where the solution set is the footprint a vehicle can reach and the parameter is the ratio of the atmospheric scale height to the planetary radius. It is the example and the test, not the subject. Computing the set exactly means a Hamilton–Jacobi–Isaacs equation on a grid, which is exponential in the state dimension and capped in practice near four or five states [3], while a rigid-body entry vehicle carries thirteen; the decomposition is what brings the problem back inside that range.

The model. For $\varepsilon \in (0, 1]$ the state is (z, v, s) , with z an altitude in layer thicknesses, v its rescaled rate and s a vector of slow coordinates, and

$$\sqrt{\varepsilon} \dot{z} = v, \quad \sqrt{\varepsilon} \dot{v} = cm(z) - 1 - \sqrt{\varepsilon} \kappa v, \quad \dot{s} = F + m(z) \Gamma,$$

with m positive and strictly decreasing and c a scalar control: a unit-mass particle in the potential Φ_c with $\Phi'_c = 1 - cm$, one soft wall and linear gravity above it. All of m, κ, F, Γ may be chosen by a selection from a bounded family, so the system is a differential inclusion and the density itself is enclosed rather than fixed.

The model is the entry oscillator rather than a caricature assembled for the occasion, and three regimes of flight are three energies $W = \frac{1}{2}v^2 + \Phi_c(z)$ in one well: the minimum is the equilibrium glide, a small oscillation is the phugoid, a large one is a skip. *The dip is short and the arc is long because the well is asymmetric*, and that asymmetry is the whole of the concentration. Three Lyapunov functionals do the quantitative work — the amplitude W for the fast motion, and two for the slow motion weighted by the density and by its square root — and two exact identities organise it: $\langle m \rangle = 1/c$ at every amplitude, so the glide is not merely the bottom of the well but the *mean* of every oscillation about it; and ε sets no shape, only a clock, so every ratio of times along a cycle is free of it. The limit system $\mathcal{H}$ flows ballistically on the arcs, jumps at $z = 0$ by a map J — the solution set of an inner problem carrying no small parameter — and flows by the averaged field $F + \Gamma/c$ on the glide. The hypotheses, the constants and the four-page proof are in the proposal.

What makes the proof elementary is that the functionals do all the counting. They bound the summed duration of the dips with no ε at all, since the passes grow like $\varepsilon^{-1/2}$ in number while each dip shortens like $\varepsilon^{1/2}$; they bound the time the motion may spend in closed cycles; and they bound the accumulated amplification, which is exponential in an impulse. No compactness argument enters. That is the point rather than an evasion: the structure is the one concentration-compactness describes, and the proposal is that a Lyapunov functional makes it quantitative where the classical dichotomy makes it only qualitative.

Where the measures enter. The theorem hands the occupation-measure program of Lasserre, Henrion and collaborators [4] a different problem. $\mathcal{R}_\mathcal{H}$ is a reachability problem in $2 + d$ coordinates with one guard and a set-valued reset, which is the hybrid form for which the relaxation is known [5]: the Liouville equation is a linear constraint on a measure, a measure on the guard carries the mass across each pass, and the dual certificate is a pair of functions non-decreasing along the flow and across the reset. The singular limit is discharged by the theorem, which is elementary, and the dimension is discharged on $\mathcal{H}$, which is low-dimensional by construction. Evaluating the

resulting bound without losing the guarantee is a separate question, and the proposal sets it aside. That division of labour is the shape of the program.

What remains. The hypothesis entry does not satisfy is the one separating the regimes. A pass entered at flight-path angle γ_e enters through the ratio $\gamma_e/\sqrt{\varepsilon}$, as in the classical Chapman variables [6], and for the Earth $\sqrt{\varepsilon} \approx 0.034$ is about two degrees while entry angles are a few degrees. So a pass is neither short against the arc that follows it nor fast against the oscillation about the glide, and at the physical ε the regimes do not separate for any accurate choice of thresholds. Three problems follow.

1. **The statement at a fixed ε .** The cycles that are neither short nor fast must be carried by the unapproximated field. That adds no term to the residual, because nothing is approximated; it costs dimension, over a total duration the budget bounds. What has to be shown is a bound on the growth of that enclosure, explicit in the interval widths — and, with it, the smaller selection class the real atmosphere permits, which is where the bound would most sharpen.
2. **Separation of scales, by enclosure.** The attitude and elastic motions do not concentrate, and averaging them is the obvious route and the wrong one: an averaging theorem gives an order and not a bound, and holds through a resonance only outside a set of small measure, which an outer bound cannot discard. They act on the trajectory only through the aerodynamic force, so it is the force that must be enclosed. Two facts are to be established: where there is no air there is no torque, and the attitude has no aerodynamic effect in any case; and where there is air the damping rate is proportional to density times speed, while the energy identity fixes the integral of density against the cube of speed, so the budget that bounds the dissipation also bounds the integrated damping from below.
3. **The companion at the level of measures.** That the limit of the occupation measures satisfies the Liouville equation of the limit system, with the mass crossing each pass carried on the graph of J , is a separate statement from the one above, and it is where the concentration machinery is expected to be needed — for the structure rather than for the constants.

The second is the part of the programme for which I am looking for an advisor.

- [1] P.-L. Lions. “The Concentration-Compactness Principle in the Calculus of Variations. The Locally Compact Case, Part 1”. In: *Annales de l’Institut Henri Poincaré, Analyse Non Linéaire* 1.2 (1984), pp. 109–145.
- [2] H. Bahouri and P. Gérard. “High Frequency Approximation of Solutions to Critical Nonlinear Wave Equations”. In: *American Journal of Mathematics* 121.1 (1999), pp. 131–175.
- [3] S. Bansal et al. “Hamilton-Jacobi Reachability: A Brief Overview and Recent Advances”. In: *56th IEEE Conference on Decision and Control (CDC)*. 2017, pp. 2242–2253.
- [4] J.-B. Lasserre et al. “Nonlinear Optimal Control via Occupation Measures and LMI Relaxations”. In: *SIAM Journal on Control and Optimization* 47.4 (2008), pp. 1643–1666.
- [5] V. Shia et al. “Convex Computation of the Reachable Set for Controlled Polynomial Hybrid Systems”. In: *53rd IEEE Conference on Decision and Control (CDC)*. 2014, pp. 1499–1506.
- [6] D. R. Chapman. *An Approximate Analytical Method for Studying Entry into Planetary Atmospheres*. Tech. rep. Technical Note 4276. National Advisory Committee for Aeronautics, 1958.