

# 2-local Hamiltonian construction

June 3, 2019

- Let  $\lambda(H)$  denote the smallest eigenvalue of  $H$ .
- Let  $H|_S = \prod_S H \prod_S$ .
- Let  $\hat{t}$  denote  $t$  represented in unary.

# Change of definition

An operator  $H : \mathcal{B}^{\otimes n} \rightarrow \mathcal{B}^{\otimes n}$  is a  $k$ -local Hamiltonian if  $H = \sum_{j=1}^r H_j$  where  $H_i$  are hermitian acting on at most  $k$  qubits.

# Change of definition

An operator  $H : \mathcal{B}^{\otimes n} \rightarrow \mathcal{B}^{\otimes n}$  is a  $k$ -local Hamiltonian if  $H = \sum_{j=1}^r H_j$  where  $H_j$  are hermitian acting on at most  $k$  qubits.

note

The positive semidefinite condition is no longer there.

# Assumptions

- The circuit contains only controlled phase gate  $C_\phi$  and one-qubit gates

# Assumptions

- The circuit contains only controlled phase gate  $C_\phi$  and one-qubit gates
- $C_\phi$  gates are preceded and followed by two  $Z$  gates

# Assumptions

- The circuit contains only controlled phase gate  $C_\phi$  and one-qubit gates
- $C_\phi$  gates are preceded and followed by two  $Z$  gates
- $C_\phi$  gates are applied at  $t \in T_2 = \{L, 2L, 3L, \dots\}$ .

# Assumptions

- The circuit contains only controlled phase gate  $C_\phi$  and one-qubit gates
- $C_\phi$  gates are preceded and followed by two  $Z$  gates
- $C_\phi$  gates are applied at  $t \in T_2 = \{L, 2L, 3L, \dots\}$ .
- There are  $T = (|T_2| + 1)L - 1$  gates in total.



# Assumptions

- The circuit contains only controlled phase gate  $C_\phi$  and one-qubit gates
- $C_\phi$  gates are preceded and followed by two  $Z$  gates
- $C_\phi$  gates are applied at  $t \in T_2 = \{L, 2L, 3L, \dots\}$ .
- There are  $T = (|T_2| + 1)L - 1$  gates in total.
- Let  $T_1 = [T] \setminus T_2$  be time corresponding to 1-qubit gates

# Projection Lemma

Let  $H_1, H_2$  be Hamiltonians where  $H_2 \geq 0$ , and  $S = \ker H_2$ . Then  $\exists J \in \mathbb{R}$  s.t.  $\lambda(H_1|_S) - \frac{1}{8} \leq \lambda(H_1 + JH_2) \leq \lambda(H_1|_S)$ .

# Projection Lemma

Let  $H_1, H_2$  be Hamiltonians where  $H_2 \geq 0$ , and  $S = \ker H_2$ . Then  $\exists J \in \mathbb{R}$  s.t.  $\lambda(H_1|_S) - \frac{1}{8} \leq \lambda(H_1 + JH_2) \leq \lambda(H_1|_S)$ .

idea

Construct  $H_2$  to “cut out” vectors outside  $\ker H_2$

## 2-local construction overview

$$H = H_{out} + J_{in}H_{in} + J_2H_{prop2} + J_1H_{prop1} + J_{clock}H_{clock}$$

## 2-local construction overview

$$H = H_{out} + J_{in}H_{in} + J_2H_{prop2} + J_1H_{prop1} + J_{clock}H_{clock}$$

- $H_{clock}$  corresponds to valid (unary) clock states

## 2-local construction overview

$$H = H_{out} + J_{in}H_{in} + J_2H_{prop2} + J_1H_{prop1} + J_{clock}H_{clock}$$

- $H_{clock}$  corresponds to valid (unary) clock states
- $H_{prop1}$  corresponds to correct propagations of 1-qubit gates

## 2-local construction overview

$$H = H_{out} + J_{in}H_{in} + J_2H_{prop2} + J_1H_{prop1} + J_{clock}H_{clock}$$

- $H_{clock}$  corresponds to valid (unary) clock states
- $H_{prop1}$  corresponds to correct propagations of 1-qubit gates
- $H_{prop}$  corresponds to correct propagation

## 2-local construction overview

$$H = H_{out} + J_{in}H_{in} + J_2H_{prop2} + J_1H_{prop1} + J_{clock}H_{clock}$$

- $H_{clock}$  corresponds to valid (unary) clock states
- $H_{prop1}$  corresponds to correct propagations of 1-qubit gates
- $H_{prop}$  corresponds to correct propagation
- $H_{in}$  corresponds to the input qubits being initialized properly



## 2-local construction overview

$$H = H_{out} + J_{in}H_{in} + J_2H_{prop2} + J_1H_{prop1} + J_{clock}H_{clock}$$

- $H_{clock}$  corresponds to valid (unary) clock states
- $H_{prop1}$  corresponds to correct propagations of 1-qubit gates
- $H_{prop}$  corresponds to correct propagation
- $H_{in}$  corresponds to the input qubits being initialized properly
- $S_{legal} \supset S_{prop1} \supset S_{prop} \supset S_{in}$

# Restriction to legal clock states

$$H_{clock} = \sum_{1 \leq i < j \leq T} I \otimes |01\rangle\langle 01|_{ij}$$

# Restriction to legal clock states

$$H_{clock} = \sum_{1 \leq i < j \leq T} I \otimes |01\rangle\langle 01|_{ij}$$

When restricted to  $S_{legal} = \ker H_{clock}$ :

- $|10\rangle\langle 10|_{t,t+1} = |\widehat{t}\rangle\langle \widehat{t}|$

# Restriction to legal clock states

$$H_{clock} = \sum_{1 \leq i < j \leq T} I \otimes |01\rangle\langle 01|_{ij}$$

When restricted to  $S_{legal} = \ker H_{clock}$ :

- $|10\rangle\langle 10|_{t,t+1} = |\widehat{t}\rangle\langle \widehat{t}|$
- $|1\rangle\langle 0|_{t+1} = |\widehat{t+1}\rangle\langle \widehat{t}|$

# Restriction to legal clock states

$$H_{clock} = \sum_{1 \leq i < j \leq T} I \otimes |01\rangle\langle 01|_{ij}$$

When restricted to  $S_{legal} = \ker H_{clock}$ :

- $|10\rangle\langle 10|_{t,t+1} = |\widehat{t}\rangle\langle \widehat{t}|$
- $|1\rangle\langle 0|_{t+1} = |\widehat{t+1}\rangle\langle \widehat{t}|$
- $|11\rangle\langle 00|_{t,t+1} = |\widehat{t+2}\rangle\langle \widehat{t}|$

# Restriction to $S_{prop1}$

$$H_{prop1} = \sum_{t \in T_1} H_{prop,t}$$

$$H_{prop,t} = I \otimes |\widehat{t}\rangle\langle\widehat{t}| + I \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| - U_t \otimes |\widehat{t}\rangle\langle\widehat{t-1}| - U_t^\dagger \otimes |\widehat{t-1}\rangle\langle\widehat{t}|$$

## Restriction to $S_{prop1}$ cont.

Let  $t \in T_2$ . Then the following are true:

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+1}\rangle)(\langle \widehat{t}| + \langle \widehat{t+1}|)$

## Restriction to $S_{prop1}$ cont.

Let  $t \in T_2$ . Then the following are true:

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+1}\rangle)(\langle \widehat{t}| + \langle \widehat{t+1}|)$
- $|1\rangle\langle 1|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle - |\widehat{t+1}\rangle)(\langle \widehat{t}| - \langle \widehat{t+1}|)$



# Restriction to $S_{prop1}$ cont.

Let  $t \in T_2$ . Then the following are true:

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+1}\rangle)(\langle \widehat{t}| + \langle \widehat{t+1}|)$
- $|1\rangle\langle 1|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle - |\widehat{t+1}\rangle)(\langle \widehat{t}| - \langle \widehat{t+1}|)$
- $|0\rangle\langle 0|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle + |\widehat{t+2}\rangle)(\langle \widehat{t+1}| + \langle \widehat{t+2}|)$

# Restriction to $S_{prop1}$ cont.

Let  $t \in T_2$ . Then the following are true:

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+1}\rangle)(\langle \widehat{t}| + \langle \widehat{t+1}|)$
- $|1\rangle\langle 1|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle - |\widehat{t+1}\rangle)(\langle \widehat{t}| - \langle \widehat{t+1}|)$
- $|0\rangle\langle 0|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle + |\widehat{t+2}\rangle)(\langle \widehat{t+1}| + \langle \widehat{t+2}|)$
- $|1\rangle\langle 1|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle - |\widehat{t+2}\rangle)(\langle \widehat{t+1}| - \langle \widehat{t+2}|)$

# Restriction to $S_{prop1}$ cont.

Let  $t \in T_2$ . Then the following are true:

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+1}\rangle)(\langle \widehat{t}| + \langle \widehat{t+1}|)$
- $|1\rangle\langle 1|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle - |\widehat{t+1}\rangle)(\langle \widehat{t}| - \langle \widehat{t+1}|)$
- $|0\rangle\langle 0|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle + |\widehat{t+2}\rangle)(\langle \widehat{t+1}| + \langle \widehat{t+2}|)$
- $|1\rangle\langle 1|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle - |\widehat{t+2}\rangle)(\langle \widehat{t+1}| - \langle \widehat{t+2}|)$
- $(|00\rangle\langle 00|_{f_t, s_t} + |11\rangle\langle 11|_{f_t, s_t}) \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+2}\rangle)(\langle \widehat{t}| + \langle \widehat{t+2}|)$

# Restriction to $S_{prop1}$ cont.

Let  $t \in T_2$ . Then the following are true:

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+1}\rangle)(\langle \widehat{t}| + \langle \widehat{t+1}|)$
- $|1\rangle\langle 1|_{f_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle - |\widehat{t+1}\rangle)(\langle \widehat{t}| - \langle \widehat{t+1}|)$
- $|0\rangle\langle 0|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle + |\widehat{t+2}\rangle)(\langle \widehat{t+1}| + \langle \widehat{t+2}|)$
- $|1\rangle\langle 1|_{s_t} \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t+1}\rangle - |\widehat{t+2}\rangle)(\langle \widehat{t+1}| - \langle \widehat{t+2}|)$
- $(|00\rangle\langle 00|_{f_t, s_t} + |11\rangle\langle 11|_{f_t, s_t}) \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle + |\widehat{t+2}\rangle)(\langle \widehat{t}| + \langle \widehat{t+2}|)$
- $(|01\rangle\langle 01|_{f_t, s_t} + |10\rangle\langle 10|_{f_t, s_t}) \otimes |\widehat{t}\rangle\langle \widehat{t}| = (|\widehat{t}\rangle - |\widehat{t+2}\rangle)(\langle \widehat{t}| - \langle \widehat{t+2}|)$

# Restriction to $S_{prop1}$ cont.

Similarly,

- $|0\rangle\langle 0|_{f_t} \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| = (|\widehat{t-1}\rangle + |\widehat{t-2}\rangle)(\langle\widehat{t-1}| + \langle\widehat{t-2}|)$
- $|1\rangle\langle 1|_{f_t} \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| = (|\widehat{t-1}\rangle - |\widehat{t-2}\rangle)(\langle\widehat{t-1}| - \langle\widehat{t-2}|)$
- $|0\rangle\langle 0|_{s_t} \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| = (|\widehat{t-2}\rangle + |\widehat{t-3}\rangle)(\langle\widehat{t-2}| + \langle\widehat{t-3}|)$
- $|1\rangle\langle 1|_{s_t} \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| = (|\widehat{t-2}\rangle - |\widehat{t-3}\rangle)(\langle\widehat{t-2}| - \langle\widehat{t-3}|)$
- $(|00\rangle\langle 00|_{f_t, s_t} + |11\rangle\langle 11|_{f_t, s_t}) \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| = (|\widehat{t-1}\rangle + |\widehat{t-3}\rangle)(\langle\widehat{t-1}| + \langle\widehat{t-3}|)$
- $(|01\rangle\langle 01|_{f_t, s_t} + |10\rangle\langle 10|_{f_t, s_t}) \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| = (|\widehat{t-1}\rangle - |\widehat{t-3}\rangle)(\langle\widehat{t-1}| - \langle\widehat{t-3}|)$

# Restriction to $S_{prop}$

Naive construction:

$$H_t = I \otimes |\widehat{t}\rangle\langle\widehat{t}| + I \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| - C_\phi \otimes |\widehat{t}\rangle\langle\widehat{t-1}| - C_\phi^\dagger \otimes |\widehat{t-1}\rangle\langle\widehat{t}|$$

$$H' = \sum_{t \in T_2} H_t$$

# Restriction to $S_{prop}$

Naive construction:

$$H_t = I \otimes |\widehat{t}\rangle\langle\widehat{t}| + I \otimes |\widehat{t-1}\rangle\langle\widehat{t-1}| - C_\phi \otimes |\widehat{t}\rangle\langle\widehat{t-1}| - C_\phi^\dagger \otimes |\widehat{t-1}\rangle\langle\widehat{t}|$$

$$H' = \sum_{t \in T_2} H_t$$

Equivalently,  $H_t =$

$$\begin{aligned} &|00\rangle\langle 00|_{f_t, s_t} \otimes (|\widehat{t}\rangle - |\widehat{t-1}\rangle)(\langle\widehat{t}| - \langle\widehat{t-1}|) + \\ &|01\rangle\langle 01|_{f_t, s_t} \otimes (|\widehat{t}\rangle - |\widehat{t-1}\rangle)(\langle\widehat{t}| - \langle\widehat{t-1}|) + \\ &|10\rangle\langle 10|_{f_t, s_t} \otimes (|\widehat{t}\rangle - |\widehat{t-1}\rangle)(\langle\widehat{t}| - \langle\widehat{t-1}|) + \\ &|11\rangle\langle 11|_{f_t, s_t} \otimes (|\widehat{t}\rangle + |\widehat{t-1}\rangle)(\langle\widehat{t}| + \langle\widehat{t-1}|) \end{aligned}$$

# Restriction to $S_{prop}$ cont.

$$H_{prop2} = \sum_{t \in T_2} H_{prop2,t}$$

$$H_{prop2,t} =$$

$$\begin{aligned} &|00\rangle\langle 00|_{f_t, s_t} \otimes 4(|\widehat{t}\rangle - |\widehat{t-1}\rangle)(\langle\widehat{t}| - \langle\widehat{t-1}|) + \\ &|01\rangle\langle 01|_{f_t, s_t} \otimes 2(|\widehat{t}\rangle - |\widehat{t-1}\rangle)(\langle\widehat{t}| - \langle\widehat{t-1}|) + \\ &|10\rangle\langle 10|_{f_t, s_t} \otimes 2(|\widehat{t}\rangle - |\widehat{t-1}\rangle)(\langle\widehat{t}| - \langle\widehat{t-1}|) + \\ &|11\rangle\langle 11|_{f_t, s_t} \otimes (|\widehat{t}\rangle + |\widehat{t-1}\rangle)(\langle\widehat{t}| + \langle\widehat{t-1}|) \end{aligned}$$



## Restriction to $S_{prop}$ cont.

$$H_{prop2,t} = H_{qubit,t} + H_{time,t} =$$

## Restriction to $S_{prop}$ cont.

$$\begin{aligned} H_{prop2,t} = H_{qubit,t} + H_{time,t} = \\ (-2|0\rangle\langle 0|_{f_t} - 2|0\rangle\langle 0|_{s_t} + |1\rangle\langle 1|_{f_t} + |1\rangle\langle 1|_{s_t}) \otimes \\ (|\widehat{t}\rangle\langle \widehat{t-1}| + |\widehat{t-1}\rangle\langle \widehat{t}|) + \end{aligned}$$

# Restriction to $S_{prop}$ cont.

$$H_{prop2,t} = H_{qubit,t} + H_{time,t} =$$

$$(-2|0\rangle\langle 0|_{f_t} - 2|0\rangle\langle 0|_{s_t} + |1\rangle\langle 1|_{f_t} + |1\rangle\langle 1|_{s_t}) \otimes$$

$$(|\widehat{t}\rangle\langle \widehat{t-1}| + |\widehat{t-1}\rangle\langle \widehat{t}|) +$$

$$(2|0\rangle\langle 0|_{f_t} + 2|0\rangle\langle 0|_{s_t} + |1\rangle\langle 1|_{f_t} + |1\rangle\langle 1|_{s_t} - 2|01\rangle\langle 01|_{f_t, s_t} - 2|10\rangle\langle 10|_{f_t, s_t}) \otimes$$

$$(|\widehat{t-1}\rangle\langle \widehat{t-1}| + |\widehat{t}\rangle\langle \widehat{t}|)$$

# Restriction to $S_{in}$

$$H_{in} = \sum_{i=m+1}^N |1\rangle\langle 1|_i \otimes |0\rangle\langle 0|$$

$$H_{out} = (T+1)|0\rangle\langle 0|_1 \otimes |T\rangle\langle T|$$

Then  $\langle \eta | H_{out} | \eta \rangle$  is the eigenvalue, as well the probability of rejection.

# Conclusion

By the projection lemma,

$$\lambda(H_{out}|_{S_{in}} - \frac{4}{8}) \leq \lambda(H) \leq \lambda(H_{out}|_{S_{in}})$$

# Conclusion

By the projection lemma,

$$\lambda(H_{out}|_{S_{in}} - \frac{4}{8}) \leq \lambda(H) \leq \lambda(H_{out}|_{S_{in}})$$

- If the circuit rejects with probability less than  $\varepsilon$ , then  $\lambda(H) < \varepsilon$ .

# Conclusion

By the projection lemma,

$$\lambda(H_{out}|_{S_{in}} - \frac{4}{8}) \leq \lambda(H) \leq \lambda(H_{out}|_{S_{in}})$$

- If the circuit rejects with probability less than  $\varepsilon$ , then  $\lambda(H) < \varepsilon$ .
- Otherwise the circuit should reject with probability greater than  $1 - \varepsilon$ . Which means  $\lambda(H) > 1 - \varepsilon - \frac{4}{8} = \frac{1}{2} - \varepsilon$ .

# Conclusion

By the projection lemma,

$$\lambda(H_{out}|_{S_{in}} - \frac{4}{8}) \leq \lambda(H) \leq \lambda(H_{out}|_{S_{in}})$$

- If the circuit rejects with probability less than  $\varepsilon$ , then  $\lambda(H) < \varepsilon$ .
- Otherwise the circuit should reject with probability greater than  $1 - \varepsilon$ . Which means  $\lambda(H) > 1 - \varepsilon - \frac{4}{8} = \frac{1}{2} - \varepsilon$ .
- So 2-local Hamiltonian is QMA-hard.