



SORBONNE UNIVERSITÉ

MASTER THESIS

Teichmüller theory and Thurston Earthquake flow

Bonnaifoux Etienne

supervised by
Carlos MATHEUS

May 27, 2020

Contents

1	Introduction	2
2	Notations	4
3	Teichmuller Theory	5
3.1	First definitions	5
3.2	Decomposition of hyperbolic surface	7
3.3	Flow on Teichmüller space	9
3.4	The collaring theorem	12
4	Isomorphisme of Mirzhakani	15
4.1	Tightening map	15
4.2	Correspondance between foliations and quadratic differentials	17
4.3	Shear Cordinate	18
4.4	Conjugaison between the horocyclic flow and the earthquake flow	20
5	Mixing rate	22
5.1	Mixing proprieties of the geodesic and horocyclic flows	22
5.2	Rate of mixing of this flows	23

1 Introduction

Since Bernhard Riemann, mathematicians knew that a geometric hyperbolic surface can be described with only a finite number of parameters. Knowing this, for a given surface, we can be interested in the set of all geometries we can give it, modulo composition by map isotopic to the identity map, this set is called the Teichmuller space. Other problems rise shortly after this definition. How can we deform in a natural way a surface's geometry into another? What does it mean that a geometry is close to another? What are the natural boundaries of the Teichmuller space?

Oswald Teichmuller, a german mathematician, studied and answered this question in the year preceding the second World War. He created the first metric on this space by finding a solution to an extremal problem: between two hyperbolic geometry on the same surface is there a function which minimize the deformation ? The resulting theorem [?] proves not only the existence, but also the unicity of this function. It naturally induced a distance in the now called Teichmuller space by considering the logarithm of the deformation of the extremal function.

Thurston then added other important steps to this theory. He underlined the role of laminations. A lamination is a generalisation of simple closed curve. And he introduced the earthquake flow, in a course in 1976-1977 at Priceton, which plays an important role in Teichmuller theory . Kerckhoff used this tool to show the Nielsen realisation conjecture in 1983 [10] which states that every finite subgroup of the mapping class group have a fixed point in the Teichmuller space. This theory have been actively studied and therefore generated a lot of literature and contributions. One can refer to the following books [5], [8], [9] or course notes [15] [16] for a more detailed study of the theory.

Still, some questions remain open, raising further interest of the community in the topic of hyperbolic geometry. Among the main questions encountered, figures the asymptotic number of closed geodesics. To begin with, one can wonder what is the number $\pi(X, L)$ of closed geodesic on a hyperbolic surface X of length less than L . The answer was found in the 1940s and 1950s by Delsarte, Huber and Selberg and -due to its resemblance to the prime number theorem- is called the prime number theorem for hyperbolic surfaces. It states that:

$$\pi(X, L) \simeq e^L/L$$

as $L \rightarrow \infty$. The reader can refer to [3] for more details.

A much harder problem was to find the number, $\sigma(X, L)$, of simple (which don't intersect themselves) closed geodesic of length less than L on a hyperbolic surface X . It was found years later, in Mirzakhani's PhD [Ref?citation?], and we have

$$\sigma(X, L) \simeq C_X L^{6g-6}$$

As $L \rightarrow \infty$ where g is the genus of the surface X and C_X is a constant which depend of the geometry X .

To obtain such a result, Myriam Mirzhakhani conjugated the earthquake flow to the horocycle



[!ht]

Figure 1: The mathematician Oswald Teichmuller

flow. This step provides that the Earthquake flow is ergodic and allows us to use Birkoff theorem [[Ref?citation?]] to understand asymptotic quantities.

The question is now to give error terms to this quantity. In order to do that, we need to understand better the mixing rate of the earthquake flow. This flow is conjugate to the horocyclic flow which has a polynomial mixing rate. But as the conjugacy is only a measurable map it does not transport the rate of mixing. We should analyse the rate of mixing of the earthquake flow by other means. One research direction would be to consider natural functions on Teichmuller space such as the systole. The systole is the length of the shortest simple closed curve on the surface. This function behaves nicely along earthquake path, it is continuous, convex and we know its first derivative at the origin. Moreover in the case of the once punctured torus we can give a frame determined by the continued fraction of the slope.

In this master thesis, I will first give an introduction to Teichmuller theory and some useful and classical tools in this theory such as the collar lemma or a surface's decomposition in pair of pants. Then, we will review the proof of the Mirzakhani's conjugacy between the horocyclic flow and the earthquake flow. After doing so, we will focus on the mixing properties of the ergodic and horocyclic flow, and discuss their mixing rate. Finally we will discuss a special case: the once punctured torus, as it is one of the simplest example of hyperbolic surface.

2 Notations

\mathcal{ML}	measured lamination
$\mathcal{MF}(S)$	space of all equivalence classes of measured foliations.
\mathcal{QD}	bundle of nonzero quadratic differential.
\mathcal{T}_g	Teichmuller space of surface of genus g
$Mod(S)$	The modular group of a surface S
$\mathcal{P}^1\mathcal{M}_g$	Product of \mathcal{T}_g and $\mathcal{MF}(S)$ quotiented by the modular group.
$\mathcal{GC}(S)$	the space of geodesic currents
\mathbb{D}	the Poincaré disk
\mathbb{S}_∞	the unit circle seen as the boundary of \mathbb{D}
$\bar{\mathbb{D}}$	$\mathbb{D} \cup \mathbb{S}$
\mathbf{S}	set of all simple closed curve of a surface
$D_\gamma(\alpha)$	Dehn's twist of α around γ

3 Teichmuller Theory

3.1 First definitions

To begin we will in this section give the very definition to introduce the theory of Teichmuller space

Definition 3.1. Let S be a surface of genus g , a marking of S is a couple (X, f) made of a closed Riemann surface X and of one homeomorphism $f : S \rightarrow X$ which preserve the orientation. On the set of the marking S , we have a equivalence relation, $(X_1, f_1) \sim (X_2, f_2)$ if there exist $\alpha : X_1 \rightarrow X_2$ such that $f_2 \circ \alpha \circ f_1^{-1}$ be an homeomorphism of S preserving the orientation and isotope to the identity map. The set of the marking quotient by the previous relation is the *Teichmuller space* and is written \mathcal{T}_g .

Remark. If $g \geq 2$, for every closed curve α of S , there is only one closed geodesic of X freely isotope to $f(\alpha)$. Moreover if α is simple, so is the geodesic. We will note $l_\alpha(X)$ its hyperbolic length and we will take the weakest topologie on \mathcal{T}_g which make this map continuous.

This give a map $L : \mathcal{T}_g \rightarrow \mathbb{R}^{\mathbf{S}}$, where \mathbf{S} is the set of all simple closed curve of a given surface. We can ask if this map is injective, i.e. that a geometry is given by the length of the set of simple closed curve. The answer is yes and more precisely one can choose only $9g - 9 + 3n$ curve so that this map is injective [5] Theorem 10.7. This give the intuition that Teichmuller space can be describe only by using a finite set of parameter. We will see after that other coordinates which have nice properties for other utilisations.

An other curious but important fact about simple closed geodesic is that the sum of a function of theirs length is equal to a constant which does not depend on the genus or the geometry of the surface.

Theorem 3.1. *Let X be a hyperbolic surface then*

$$\sum_{\gamma} \frac{1}{1 + e^{l_\gamma(X)}} = \frac{1}{2}$$

Where the sum is taken over all simple closed geodesic.

This theorem was prove by McShane and bear his name [17]. A good introduction with a proof using probabilistic method on tree can be found in this paper [11].

Definition 3.2. The *modular group* is the group of homeomorphisms of S which respect orientation quotiented by the subgroup of homeomorphisms isotopic to the identity map. We will call this group Mod_g . It act discretely on the Teichmuller space \mathcal{T}_g and the quotient is called the *moduli space* and is written \mathcal{M}_g .

The Teichmuller space is the universal covering of the modular space. In fact the Teichmuller space is homeomorph to a sphere and so is contractible and in particular, connected and simply connected. It is the universal covering of the moduli space.

We can ask ourselves how this spaces look and if we can give an easy representation of them. The fact is that the modular space is not manifold but an orbifold, that is a space which locally look like a ball of a vector space quotiented by a finite group.

We will now describe objects that are natural to understand the Teichmuller space.

Definition 3.3. A *foliation* on a surface S is the collection of the following data: a finite set of points $P = (p_1, p_2, \dots)$, given a open covering U_i on $S - P$, a collection of C^1 real function ν_i

such than $\|d\nu_j\| = \|d\nu_i\|$ on $U_i \cap U_j$, and near each singular point p_s a coordinate neighborhood V with complex coordinate z such that $\|d\nu\| = \|Im(z^{\frac{k}{2}}dz)\|$ for some positive integer k called the degree of the singular point.

Leaves of the foliations are the graphs immersed in S along which $d\nu$ is constant. In addition if the surface S have a boundary we is required that it is contained in a singular leaf.

The height $h_\gamma(\|d\nu\|)$ of a (free homotopy class) of a loop γ on S is the infimum in the homotopy class of the integral by $\|d\nu\|$

$$h_\gamma(\|d\nu\|) = \inf_{\gamma \sim \gamma'} \int_\gamma \|d\nu\|$$

The topology on the measured lamination that we will use is the weakest that make the height functions continous.

Remark. We won't actually study the set of measured foliation but the equivalence class of

$$h_\gamma(\|d\nu\|) = h_\gamma(\|d\mu\|), \text{ for each loop } \gamma \in S$$

. We can equivalently use Whitehead equivalence relation on singular foliations by collapsing critical intervals to points and taking isotopy of foliation.

Let $\mathcal{MF}(S)$ be the space of all equivalence classes of measured foliations.

Definition 3.4. A *lamination* is a closed set made of an union (non necessarily finite) of geodesics. For each point x in λ , a lamination, passed only one geodesic of the lamination. We will write this space $\mathcal{ML}(x)$.

To understand the space of geodeisc lamination it is sometimes easier to understand the "trace" of this space in the boundary circle of the universal cover of tge surface. This observation gives the following definition.

Definition 3.5. Let \mathcal{M}_∞ be the space of unordered pairs of distinct points in \mathbb{S}^1

$$\mathcal{M}_\infty := (z, w) \in \mathbb{S}^1 \times \mathbb{S}^1, z \neq w / (z, w) \equiv (w, z)$$

Let G be a discret torsion-free group in $PSL(2, \mathbb{R})$ such that $\mathbb{H}/G = S$ is a hyperbolic surface. A *geodesic current* μ on S is a G -invariant Radon measure on \mathcal{M}_∞ . We will note $\mathcal{GC}(S)$ the space of geodesic currents, see [9]

Remark. $\mathcal{GC}(S)$ have a natural topology which is the weak $*$ convergence on continous functions.

Exemple 3.1. Let $g \in G$ be an element that represent a simple closed geodesic on the surface S . The G -orbit of the fix point of g is a discret set on which we can put diracs measure. This link the geodesic currents to the measured lamination.

Remark. A multicurve is a formel sum of geodesics $\gamma = \sum a_i \gamma_i$. The space of lamination is in some aspect the closure of the set of all multicurve

Definition 3.6. Consider the square $\mathcal{M}_\infty^2 := \mathcal{M}_\infty \times \mathcal{M}_\infty$. In this space we can consider the open subset \mathcal{IM}_∞^2 corresponding to pair pairs of geodesics which have transversal intersections in \mathbb{H} . G act on \mathcal{IM}_∞^2 . If μ and ν are geodesic currents in $\mathcal{GC}(S)$, the product $\mu \times \nu$ define a G -invariant measure on \mathcal{IM}_∞^2 . Finally if we take the mass of the total space \mathcal{IM}_∞^2/G , the reasult is called the *intersection number*, $i(\mu, \nu)$

Proposition 3.2.

$$i : \mathcal{GC}(S) \times \mathcal{GC}(S) \rightarrow \mathbb{R}_+$$

is continuous and bilinear [2].

Remark. If α and β are simple closed geodesics (dirac measure in $\mathcal{GC}(S)$), then the intersection number is the number of intersection between α and β . Actually, one can define intersection in this way, first on simple closed geodesic, then by bilinearity on multi-curves and finally by continuity on geodesic current.

Remark. The topology on \mathcal{ML} is the weakest that make $i(.,.)$ a continuous function.

Definition 3.7. A *quadratic differential* is a section of the square of the tangent space to X . It is locally as $\phi = \phi(z)dz^2$.

Remark. If $\phi(p) \neq 0$ we can find a map including p in which $\phi = dz^2$. Hence ϕ determine a flat metric on X and a foliation \mathcal{F} corresponding to horizontal lignes.

A quadratic differential is said to be integral if


$$\|\phi\| = \int_X |\phi| < \infty$$

We will write $\mathcal{Q}(X)$ the Banach space of integral quadratic differentials.

3.2 Decomposition of hyperbolic surface

One way to construct all hyperbolic surface is to decompose them in elementary piece, that we will call pair of pant.

A hyperbolic geometric exercise show that a hexagone which side are geodesics and with right angles is determined by the length of three sides which are not consecutifs.



On the image above γ_i , γ_j and γ_k determined the hexagone. Then we can glued them to have a pair of pant.

Definition 3.8. A *pair of pant* is a hyperbolic surface with three geodesic boundaries and no ponctured.

Remark. The pair of pant is uniquely determined by the length of the three boundary geodesics.

Proof. As a hyperbolic polygon is uniquely determined by three non consecutif sides, the surface with boundaries that we obtain by gluing two together is determined by the length of its three boundary geodesics. \square

Remark. The length of one or more geodesic can go to zero and the boundaries become a punctured.

We can now decompose, with the following theorem, all hyperbolic surfaces in a collection of pair of pants.

Theorem 3.3. *Let S be a surface of genus g and with n punctured. There is a set of $3g - 3 + n$ simple closed curves $(\gamma_1, \dots, \gamma_{3g-3+n})$ such that $S \setminus \gamma_i$ is a disjoint collection of pair of pants.*

Definition 3.9. Given a surface S and a pant decomposition $\gamma_1, \dots, \gamma_{3g-3+n}$, we have a map

$$(S) \rightarrow (\mathbb{R}^{+3g-3+n}, \mathbb{R}^{3g-3+n}) X \mapsto (l_{\gamma_1}(X), \dots, l_{\gamma_{3g-3+n}}(X), \tau_{\gamma_1}(X), \dots, \tau_{\gamma_{3g-3+n}}(X))$$

This map is injective and is call the *Fenchel-Nielsen* coordinates.

Definition 3.10. Dehn twist Let γ be a simple closed curve. There is a tubular neighborhood of γ called A homeomorph to $[0; 1] \times S^1$. A *Dehn's twist* around γ is the homeomorphism which is the identity out of A and is $(t, s) \mapsto (t, e^{2i\pi t} s)$ on A .




Figure 2: A Dehn twist

We can now give a set of generator of the modular group.

Remark. The Lickorisk theorem states that the modular group is generated by the Dehn's twist and more precisely that one can choose only $2g + 1$ generators [13]. We will give here a easier version of this theorem.

Theorem 3.4. *There is a collection $\delta_1, \dots, \delta_{9g-9}$ of simple closed curves such that $\mathcal{T}_g \rightarrow \mathbb{R}^{9g-9}$ is injective.*

Proof. Let take $(\gamma_1, \dots, \gamma_{9g-9})$ a decomposition in pair of pants, $(\alpha_1, \dots, \alpha_{9g-9})$ be a collection of simple closed curves such that $i(\gamma_i, \alpha_i) > 0$ and $i(\gamma_i, \alpha_j) = 0$ for $i \neq j$, finnely we take $\beta_i = D_{\gamma_i}(\alpha_i)$. We want to show that the length of this collection of $9g - 9$ curves determined the hyperbolic structure $X \in \mathcal{T}_g$. X already has the Fenchel-Nielsen coordinate, of the pant decomposition $(\gamma_1, \dots, \gamma_{3g-3})$, $(l_{\gamma_1}(X), \tau_{\gamma_1}(X), \dots, l_{\gamma_{3g-3}}(X), \tau_{\gamma_{3g-3}}(X))$ so we need only to show that the parameters $\tau_{\gamma_i}(X)$ are determined by the lenght of the collection. Up to a renormalisation we can take $\tau_{\gamma_i}(X) = 0$ for every i . Now let's take $t = (t_1, \dots, t_{3g-3}) \in \mathbb{R}^{3g-3} \setminus 0$. We will note X_t the hyperbolic geometry which has the same length as X and twist parameters t in the Fenchel-Nielsen coordinate of the pair of pants. So $X_0 = X$. We will consider the function $A(t) = l_{\alpha_1}(X_t)$ and $B(t) = l_{\beta_1}(X_t)$. This function depend only of t_1 as $i(\gamma_i, \alpha_j) = 0$ for

$i \neq j$. Moreover they are strictly convex and by definition we have $A(t_1 + l_{(\gamma_1)}(X)) = B(t_1)$. We will show that there is no $t_1 \neq 0$ such that $A(t_1) = A(0)$ and $B(t_1) = B(0)$ that is $A(t_1 + l_{(\gamma_1)}(X)) = A(l_{(\gamma_1)}(X))$. We will note $s = t_1$ and $L = l_{(\gamma_1)}(X)$. Suppose there is $s \neq 0$ such that $A(s) = A(0)$, we can take $s > 0$, the other case is symmetric. If $s < L$, then by convexity for every $t \in]0; s[$, $A(t) < A(0) = A(s)$ and A is strictly increasing after s so as $s < L < L + s$ we have $A(L) < A(L + s)$. If $s > L$, then $L < s < L + s$ and $A(L) < A(L + s)$. The final case $s = L$ is also impossible since we would have $A(0) = A(L) = A(2L)$. Finally we can make the same argument for the other twist parameters which conclude the proof. \square

3.3 Flow on Teichmüller space

We will define the main object of this paper, earthquake flow.

Definition 3.11. The *earthquake flow* is family of maps defined for $t \in \mathbb{R}$

$$\begin{aligned} E_t : \mathcal{ML} \times \mathcal{T}_g &\rightarrow \mathcal{ML} \times \mathcal{T}_g \\ (\lambda, X) &\mapsto (\lambda, E_{t\lambda}X) \end{aligned}$$

where $E_{t\lambda}$ is first defined if λ is a simple closed curve. In this case, we open the surface along λ , then twist the left part of t unit and put it together again. Then if γ_1 and γ_2 are two curves that don't intersect, E_{γ_1} and E_{γ_2} commute. So we can define the earthquake map on weighted multicurves by twisting one curve after the other by amounts proportional to the weight. Finally as multicurves are dense in the set of lamination we can extend this map for any lamination [10].




Figure 3: Effect of a twist on a transverse curve, image from [20]

Remark. If the lamination is just a simple closed curve γ then $E_{l_\gamma(X)}(X, \gamma)$ is just a Dehn twist around γ . Moreover if we take a decomposition in a pair of pant that contain γ , it is just a translation in the coordinate of the twist of γ .

One could give a precise proof that the extension of the earthquake map from the multicurves to the lamination is rigorous, that is if we take two sequence α_n and α'_n of multicurves which converge to the same lamination, then the sequences of earthquake map along this multicurves converge to the same map.

We will work on universal covering of the surface, the half-plane \mathbb{H} . If $v \in T^1\mathbb{H}$ we will call $\gamma(v)$ the geodesic passing by the basepoint of v and with v as tangent vector at this point.

We have to use a useful lemma to control the distortion of the earthquake map. We will not prove them, but their proofs are in [10].

Lemma 3.5. *Let l and l' be two geodesics and $x \in l$, $y \in l'$ two points at most ϵ apart. Then if v and v' are the two geodesic tangents to l and l' respectively at point x and y , we have $d(v, v') < C\epsilon$ for a universal constant C .*

Lemma 3.6. *Let v and v' be two vectors such that $d(v, v') < \epsilon$, let denote $\gamma = \gamma(v)$ and $\gamma' = \gamma(v')$. Let $w \in T^1\mathbb{H}$, then*

1. $d(E_{t\gamma}w, E_{t\gamma'}w) \leq Kt\epsilon$
2. $d(E_{t\gamma}w, w) \leq Kt$

for all $t \leq T$ and for a constant K depending on T and on the distance between the basepoint of v and w .

With this two lemma, we can describe what happend if we change a discret lamination by a simple closed curve that average it.

Lemma 3.7. *Let $x, y \in \mathbb{H}$, $v \in T_1\mathbb{H}$ based at y , \bar{A} the geodesic from x to y . Suppose γ is a discrete lamination with equal measure on each leaf whose intersection with \bar{A} is include in a subarc A . Let l be a single geodesic intersecting A with angle equal to the average angle of the intersections of γ and A and with mass equal to $\mu = i(A, \gamma)$.*

If A has length less than δ then for every $T \in \mathbb{R}^+$ the distance between $E_{t\gamma}v$ and $E_{tl}v$ is less than $Kt\mu\delta$, for all $t \leq T$ and K is a constant depending only of $T\mu$ and $d(x, y)$.

Proof. Let denote l_1, l_2, \dots, l_n the leaf of γ . The image of \bar{A} is a disconnected arc. The point x is connected to $E_{t\gamma}y$ by a staircase path going along component of $\bar{A} \setminus \gamma$ and subarc of l_i . Let denote the successive component A_0, A_1, \dots, A_n and δ_i the length of A_i . So the staircase path is moving by δ_0 along A_0 then by μ/n along l_1 , and so on.




Figure 4: Image of the construction described, image from [10]

We now alter the path by replacing the shearing along l_n distance μ/n by a shearing along l_{n-1} distance $2\mu/n$. The change is less than $Kt\frac{\mu}{n}C\delta_{n-1}$ by lemma 3.5 and lemma 3.6. Then we change the shearing by a shearing along l_{n-2} of distance $3\mu/n$ and we continue until we shear a distance μ along l_1 . The total change is less than $KtC \sum_{i=1}^{n-1} \frac{i\mu}{n} \delta_{n-i}$ which is less than $KCt\mu\delta$. We now pass from l_1 to l with lemma 3.5 and 3.6, and we obtain the new lemma. \square

We need a final lemma to conclude of the well-founded definition of the earthquake flow. In this case we control the difference after earthquaking by two simple geodesic going by the same point of the arc.

Lemma 3.8. *Let x, y, v , and \bar{A} be as above, if l and l' be geodesic of \mathbb{H} with measure μ and μ' such that $l \cup \bar{A} = l' \cup \bar{A} = p \notin \{x, y\}$ and the difference between the vectors tangent of l and l' at p of length μ and μ' is less than ϵ . Then for any T , $d(E_{tl}v, E_{tl'}v) < Kt\epsilon$, for $t \leq T$ and K a constant which depend only on $d(x, y)$ and $T\mu$*

Finnally this give the theorem which control the distance between two earthquake path.

Proposition 3.9. *Let $\nu \in \mathcal{ML}$ be a lamination and let x, y be in \mathbb{H} , A be the geodesic from x to y and $v \in T^1\mathbb{H}$ be based at y , and x and y do not lie on the atomic part of ν . Then for any ϵ, T , there is a neighborhood U of ν in \mathcal{ML} such that for all $\gamma, \bar{\gamma}$ weighted multicurve in U , $d(E_{t\gamma}v, E_{t\bar{\gamma}}v) < Kt\epsilon$, for all $t \leq T$, K a constant depending only on $d(x, y)$ and $Ti(\nu, A)$*

Corollaire. *The earthquake flow is well defined along any lamination $\nu \in \mathcal{ML}$ and for all time t .*

Remark. The earthquake flow is an isometry outside the support of the lamination and is continuous outside the atomic part, i.e. the simple closed geodesics of the lamination.

Thurston show that given two point in the Teichmüller space, there is a lamination λ such that the earthquake flow from one point with respect to λ reach the other point (also in [10]).

We can ask ourselves what is an invariant measure of this flow.

Definition 3.12. The *Weil-Peterson form* is the the form

$$\omega_{WP} = \sum dl_i \wedge d\tau_i$$

Where (l_1, \dots, τ_1) are the Fenchel-Nielsen according to a pant decomposition.

This give a measure μ_{WP} .

There is a finite measure ν_g in the Lebesgue measure class on $\mathcal{P}^1\mathcal{M}_g$ which is invariant under the earthquake flow. This measure projects to the volume form given by $B(X) \times \mu_{WP}$ on \mathcal{M}_g , where

$$B(X) = \mu_{Th}(\lambda \in \mathcal{ML}, l_\lambda(X) \leq 1)$$

There are two other important flows, the geodesic flow and the horocyclic flow. First there is a natural homeomorphism between $T^1\mathbb{H} \simeq SL_2(\mathbb{R})/SO_2(\mathbb{R})$, since $SL_2(\mathbb{R})/SO_2(\mathbb{R})$ act simply transitively on it. This morphism can be choosen up to a conjugaison via an other element of $SL_2(\mathbb{R})/SO_2(\mathbb{R})$. We we will be interested in a special kind of subgroup.

Definition 3.13. A *fuchsian group* Γ is a finitely generated and discrete subgroup of $SL_2(\mathbb{R})$. Then Γ act discontinuously on \mathbb{H} .

Then a Hyperbolic surface can be represented as $\mathbb{H}/\Gamma = SL_2(\mathbb{R})/SO_2(\mathbb{R})/\Gamma$. If U is a one parameter subgroup of $SL_2\mathbb{R}$ it act on the quotient.

There are two important exemple:

Definition 3.14. The *geodesic flow* is a flow on the Teichmuller space given by the action of the diagonal matrices

$$u_t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}$$

Definition 3.15. The *horocycle flow* is a flow on the bundle of nonzero quadratic differential, QD , of the Teichmuller space given by the unipotent action of

$$u_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$$

The geodesic flow is also the flow that we obtain by following geodesic line on \mathbb{H} and the horocycle is the flow we obtain by following curves which are everywhere orthogonal to the geodesic, which is the horizontal line and the circle tangent to the real line.




Figure 5:




Figure 6: The conjugaison of the horocycle flow by the geodesic one.

An important relation is how this two flows interact between each other

$$\begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix} u_t = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^t \end{pmatrix} = \begin{pmatrix} 1 & se^{2t} \\ 0 & 1 \end{pmatrix}$$

So the the conjugaison of the horocyclic flow by the geodesic one is still the horocyclic flow.

We give the definition of a third flow that we will used in demonstration of the ergodicity of the two previous one.

Definition 3.16. The *rotational flow* is a flow on the bundle of nonzero quadratic differential, \mathcal{QD} , of the Teichmuller space given by the action of

$$e_\theta = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix}$$

We have the relation

$$h_s = e_\theta g_t e_{\pi+\theta}$$

3.4 The collaring theorem

We will now give a useful tool to give necessary condition on length of two intersecting geodesics.

The collar function $\eta :]0; \infty[\rightarrow]0; \infty[$ is defined as follow. We draw a segment of length $l > 0$ on a geodesic γ , then we project perpendiculary to the geodesic the end of this segment to infinity and draw the geodesic δ which have this endpoint. So we have $\eta(l) = d(\gamma, \delta)$.




Figure 7: In red the segment of length l and in green the geodesic whose endpoints are orthogonal projection of the end of the segment.

It is an exercise to show that:

$$\eta(l) = \frac{1}{2} \ln \left(\frac{\cosh(l/2) + 1}{\cosh(l/2) - 1} \right)$$

This quantity will code a long in the "tube" generated by a simple closed geodesic. We give a definition to make this a little more precise.

Definition 3.17. Let γ be a simple closed geodesic of length l on a hyperbolic surface X . If the δ -neighborhood

$$A_\delta(\gamma) := \{x \in X \mid d(x, \gamma) < \delta\}$$

is isometric to the δ -neighborhood of the unique simple closed geodesic on the cylinder of modulus $\frac{\pi}{l}$, we say that γ admits a δ -collar, or that $A_\delta(\gamma)$ is the δ -collar of γ .

We can now state a useful theorem.

Theorem 3.10. Let X be a complete hyperbolic surface, and let $\Gamma := \gamma_1, \dots$ be a collection of disjoint simple closed geodesics, each γ_i of length l_i . Then $A_{\eta(l_i)}(\gamma_i)$ are collars around the γ_i , and they are disjoint.

Proof. Choose γ_1 and γ_2 and add other simple closed curves to have a maximal multicurve that includes both. Now cutting along this curve we have a set of pairs of pants so we only have to show that the $\eta(l_i)$ neighborhood of γ_i the boundaries of the pair of pants do not intersect each other. We cut the pair of pants along geodesics coming from a boundary C and meeting the two other boundaries A and B . We unfold this figure in the hyperbolic plane and name the side of the octogone following the figure below.

Since a' and b' have the common perpendicular C' , they do not intersect and similarly for a'' and b'' . The theorem follows easily by the definition of the function η .

□

There are some corollaries which follow from this theorem and are ready to use in many occasions.




Figure 8: image from [8]

Corollaire. Let X be a hyperbolic surface, and γ_1, γ_2 two simple closed geodesics on X of lengths l_1 and l_2 . If $l_2 < 2\eta(l_1)$, then either $\gamma_1 = \gamma_2$ or $\gamma_1 \cap \gamma_2 = \emptyset$

Proof. If $\gamma_1 \neq \gamma_2$ and $\gamma_1 \cap \gamma_2 \neq \emptyset$ then γ_2 must cross the collar neighborhood of γ_1 from one boundary to the other and so have length strictly superior than $2\eta(l_1)$. \square

Corollaire. Let X be a hyperbolic surface, and let γ_1, γ_2 be two simple closed geodesics with lengths $< \ln(3 + 2\sqrt{2})$. Then either $\gamma_1 = \gamma_2$ or $\gamma_1 \cap \gamma_2 = \emptyset$.

Corollaire. Let X be a complete hyperbolic surface, γ a simple closed geodesic on X of length l , and A_γ the collar around γ . Then any simple geodesic δ on X that enter A_γ either intersect γ or spirals towards γ .

Proof. Suppose the geodesic δ enters A_γ . We can lift the situation in the universal cover of the hyperbolic disc, where $\tilde{\gamma}$ is a lift of γ can be a diameter of the circle. Then if $\tilde{\delta}$ do not intersect $\tilde{\gamma}$ and do not have the same point at infinity, then its two endpoint are on the same side of $\tilde{\gamma}$ in the disc. Now the translation along $\tilde{\gamma}$ of length $l_\gamma(X)$ is in the representation of $\pi_1(X)$. If $\tilde{\delta}$ intersect A_γ then by the definition of δ it will intersect with its image by the translation cited before and hence is not simple in X . \square

4 Isomorphisme of Mirzhakani

The aim of this part is to demonstrate the following statement

Theorem 4.1. *There is a measurable conjugacy F between the earthquake flow $(\lambda, X) \mapsto (\lambda, E_{t\lambda}(X))$ on $\mathcal{ML} \times \mathcal{T}_g$ and the Teichmüller unipotent flow action of*

$$u_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$$

on the bundle \mathcal{QD} of nonzero quadratic differentials over Teichmüller space \mathcal{T}_g .

$$\begin{array}{ccc} \mathcal{ML} \times \mathcal{T}_g & \xrightarrow{E_t} & \mathcal{ML} \times \mathcal{T}_g \\ F \downarrow & & \downarrow F \\ \mathcal{QD} & \xrightarrow{u_t} & \mathcal{QD} \end{array}$$


To do this we will decompose F between intermediates maps.

4.1 Tightening map

A first coresspondance, found by Thurston, exist between measured foliations and measured laminations. We will mostly follow the paper of Levitt [12].

Definition 4.1. We say that two foliations are equivalent if we can pass from one to the other by Whithead (see definition below) moves or isotopy (homeomorphism isotopic to the identity).

Definition 4.2. Given a measure foliation, an critical segment γ is an arc between two singularities along a leaf which is not a simple closed curve. There is a map f homotopic to the identity that collapse γ to a point x and is identity outside a neighborhood of γ which contain no other singularity. Doing so we reduce the number of singularities of the foliation and if the extremities of γ are singularities of order k_1 and k_2 , x is now a singularity of the new foliation of order $k_1 + k_2 - 2$. This action is called a *Whithead move*.



Theorem 4.2. *Let X be a closed orientable hyperbolic surface and \mathcal{F} a foliation. There is a canonical geodesic lamination $\gamma(\mathcal{F})$ associated to \mathcal{F} . If \mathcal{F} and \mathcal{F}' are associated foliation then $\gamma(\mathcal{F}) = \gamma(\mathcal{F}')$. In the opposite direction given a geodesic lamination γ , one can find a foliation \mathcal{F} such that $\gamma(\mathcal{F}) = \gamma$ and it's unique up to equivalence.*

Definition 4.3. A *tranverse curve* is a simple closed curve C which is never tangent to \mathcal{F} and contain no singularity of \mathcal{F} .

Remark. Since \mathcal{F} contain only saddle singularities, C cannot be contractible, therefor C is isotopic to a simple closed geodesic.

We will work on the universal cover of X , which is the Poincaré disc \mathbb{D} with circle "at infinity" \mathbb{S}_∞ . We call $p : X \rightarrow \mathbb{D}$ the universal projection. $\tilde{\mathcal{F}}$ is $p^{-1}(\mathcal{F})$.

We will say that a foliation follow the (*) condition if the following is true:

If f_1 and f_2 are two compact homotopic leaves then all leaf in the open annulus between them is also compact.

Lemma 4.3. *Let h be a leaf of $\tilde{\mathcal{F}}$. Each end of h converge to a point of S ; the two point at infinity cannot be the same.*

Proof. First, we should notice that the behavior of leaf at infinity do not change if we take an equivalent foliation. Indeed a homeomorphism ϕ on a compact fundamental domain isotopic to the identity can be extend to an homeomorphism $\tilde{\phi}$ on \mathbb{D} such that $dist(x, \tilde{\phi}(x)) \leq K$. This implice that $\tilde{\phi}$ extend es the identity on the boundary \mathbb{S}_∞ .

Then given a leaf h of $\tilde{\mathcal{F}}$, we take a half leah h_0 . If $p(h_0)$ is compact or spiral toward a compact leaf of \mathcal{F} then the first part of the lemme is imediate.

Otherwise, $p(h)$ meet a transverse curve C infinitely often. With an isotopie we can take C to be a geodesic. Now h_0 can meet a connect component of $\tilde{C} = p^{-1}(C)$ only one time. Otherwise there will be a disk bound by an arc of \tilde{C} and an arc of h_0 , which is impossible considering the transversity of C and that \mathcal{F} have no 1-type singularities.

Now every compact of \mathbb{D} meet a finite number of connected components of \tilde{C} so the limit set of h_0 must be on \mathbb{S}_∞ . This limit set is connected and non empty. Moreover it should not contains any end of a connected component of \tilde{C} . But the ends of connected components of \tilde{C} are dense in \mathbb{S}_∞ as \tilde{C} is the image of a geodesic by $\pi_1(X)$. This show the first point of the lemma.

The second assertion is clear if $p(h)$ is compact or if it meet a transverse curve C at least twice since then every connected components of \tilde{C} separate the end of h .

Otherwise $p(h)$ spiral toward two compact leaf f_1 and f_2 . If $f_1 = f_2$ and the two end point of h are the same then there will be a singularity that would not be a saddle. $f_1 \neq f_2$ is impossible since \mathcal{F} follow the condition (*). \square

We can now associate to every leaf h a geodesic $\gamma(g)$ by joining the endpoint. Then $\gamma(\tilde{\mathcal{F}}) = \cup_{h \in \tilde{\mathcal{F}}} \gamma(h)$ is a disjoint union of geodesic invariant by $\pi_1(X)$. We have to show that this set is closed to concluded that we have a lamination.

Lemma 4.4. *$\gamma(\tilde{\mathcal{F}})$ is closed in $\bar{\mathbb{D}}$*

Proof. Let $g_n = \gamma(h_n)$ be a sequence of geodesics in $\gamma(\tilde{\mathcal{F}})$ converging to a geodesic g . We want to show $g \in \gamma(\tilde{\mathcal{F}})$. We can suppose that all the g_n are distinct of g and are all on the same side.

Let L be the limit set in $\bar{\mathbb{D}}$. For all leaf m in $\tilde{\mathcal{F}}$, we call \bar{m} the closure of m by adding the two end point in \mathbb{S}_∞ . Then L meet at least one connected component of $\bar{\mathbb{D}}$ \bar{m} . As the end points of all leaf of $\tilde{\mathcal{F}}$ is a dense subset of \mathbb{S}_∞ , L contain a leaf h . Taking a half-leaf h_0 , we want to show that the end point is the same as one of g .

A first case is if there is a simple closed curve C transverse to \mathcal{F} which meet $p(h_0)$ infinitely often. If h_0 does not converge to the corresponding point at infinity then there would be a connected component of $p^{-1}(C)$ that contains the point of infinity of h_0 but does not contain the point of infiny of h_n which is impossible for large n .

A second case is if $p(h_0)$ spirals toward a compact leaf, then closed leaf close to $p(h_0)$ also spirals toward the same compact leaf. Then h_0 converges to one of the points at infinity of g which is a point at infinity of h_n for n large.

Finally if $p(h)$ is compact then $p(h_n)$ spirals toward it for large n , therefore $\gamma(h)$ and g have one point in common at infinity. If the second was different, by applying a transformation leaving $\gamma(h)$ invariant (but not g), we would separate h from the leaves h_n , and it is a contradiction. □

Now we want to exhibit an inverse construction which takes a lamination λ and gives a foliation μ . To do this we still consider $\tilde{\lambda}$ in the universal cover. We will suppose that every complementary region is an ideal polygon.

We can build a skeleton that it compose of edge between vertex and a chosen point in the center. After building the skeleton for every polygon we fill the complementary region which will be between four edges by line between the two vertices.

This map is often called the "collapsing" map and its inverse the "tightening" map.




Figure 9: In red the skeleton of the ideal triangle in blue, Image from [19]

The measure we put on this foliation is uniform on every region between four edges of the skeleton.

4.2 Correspondance between foliations and quadratic differentials

For a quadratic differential q , one can define two measured foliations, the horizontal $h(q)$ and the vertical $v(q)$ corresponding in local coordinate to $Re(z)$ and $Im(z)$.

Moreover the measure are

$$h_\nu = \int Im(\sqrt{q(z)}dz)$$

and

$$h\mu = \int Re(\sqrt{q(z)}dz)$$

This gives a map from the pair of foliations but it is not the subject if we should restrict to the image. Define $\Delta = (\alpha, \beta) : i(\alpha, \gamma) = i(\beta, \gamma) = 0$, for some $\gamma \in \mathcal{MF}$. Δ contains the diagonal (α, α) and is kind of "fat" diagonal.

Lemma 4.5. *For any $q \in \mathcal{QD}$, $(h(q), v(q)) \notin \Delta$*

Proof. Suppose that there is γ such as $i(h(q), \gamma) = i(v(q), \gamma) = 0$ for some $\gamma \in \mathcal{ML}$. Let take a sequence of simple closed weighted curves γ_i converging to γ . By continuity of the intersection number we have that $i(h(q), \gamma_i) \rightarrow 0$. So there is a sequence of saddle connection in the same homotopy class of γ_i whose x -components is very small. We can have the same argument in the vertical direction and find a contradiction. \square

Theorem 4.6. *The map $q \mapsto (h(q), v(q))$ define a homeomorphism $\mathcal{QD} \rightarrow \mathcal{MF} \times \mathcal{MF} \setminus \Delta$*

Proof. We can describe the inverse map. If we take to measured foliation h and v we can tighten them into lamination (which we also call h and v) as in the previous section. This two laminations do not share any leaf, otherwise we would have $(h, v) \in \Delta$ by considering the leaf as a geodesic. The complementary region of $h \cup v$ are compact polygons, i.e. they do not have a vertex on the boundary of the disk. Now we can fill the polygon to obtain the quadratic differential with a singularity in each complementary region of order the number of side of the polygon. \square

More detail of the proof can be found in [4] Lemma 6.2 and [7]. The injectivity is discuss with more detail in [6] section 3.

4.3 Shear Coordinate

Finally there is a map that, given a hyperbolic structure X and a lamination λ create a measured foliation which is transverse to λ .

For simplicity we will ask that λ is a maximal lamination i.e. if $\tilde{\lambda}$ is the pre-image of λ in the universal cover \mathbb{D} , $\mathbb{D} \setminus \tilde{\lambda}$ is made of ideal triangle. We will first work in this triangles minus a region on the center, then give a measure in this foliation, and finally show that it is a homeomorphism.

So in one the ideal triangle, given two side we can draw an arc perpendicular with the this two sides, which is the intersection of the ideal triangle with the circle tangent to the boundary of \mathbb{D} in the common extremity the two arc chosen. Then by rescaling by a factor $r \in [0; 1]$ and doing the same procedure for the two other pair we get a foliation in the ideal triangle minus a locus in the center.




Figure 10: Image from [14]

Then we can find a full foliation by pitching the resulting.

We have a natural transverse measure to this foliation. For an arc in one ideal triangle of the lamination we can project it along the leaf of the foliation to a segment in the edge of the ideal triangle, the length of the arc will be the length of the segment. As the leaf are horocycle circle based on the same point, it will not depend of which side we choose to project. Then given an arbitrarily arc we decompose it along the ideal triangle it meet.

We want to show that this construction is reversible, that is given $\mu \in \mathcal{MF}_\lambda$, the set of foliation transverse to the lamination, we can construct $X \in \mathcal{T}_g$ whose horocycle foliation is μ .

The idea is that, if we already know X , the lamination λ can be lift to $\tilde{\lambda}$ which is invariant of Γ the fushian group of X . But we can built $\tilde{\lambda}$ only with the information given by μ .

We will note $\tilde{\mu}$ the preimage of μ in \mathbb{D} . If we consider two triangles T_1 and T_2 that are complementary region of $\tilde{\lambda}$ and we suppose we take a segment A in a leaf of $\tilde{\mu}$ that goes to an edge of T_1 to T_2 . We name v_1 and v_2 the two vectors with footprint in the edge and tangent to them. Then there is a Moebuis transformation S which take one to the other. With one more information, the "shear", we can place T_2 on \mathbb{D} , according to the position of T_1




Figure 11: Image from [19]

Indeed given only a Moebuis transformation we still have a one parameter families of triangle T_2 with an edge generated by v_2 . To fix this we trace two orthogeodesic coming from the vertex of the ideal triangles facing the considered edges and from the point of intersection in T_1 we follow a leaf of the foliation, then when we meet T_2 we have to move along the geodesic edge to find the other point of intersection. This length is the shear between T_1 and T_2




Figure 12: Image from [19]

Let I be the set of all triangles in \mathbb{D} that A meet. For each $i \in I$ we can define v_i^+ and v_i^- the vectors tangent to the edge of the corresponding triangle at the intersection of the edges and A . Note that I is a countable totally ordered but non well ordered set. So if we take S_i the Moebuis transformation which take v_i^- to v_i^+ , we have to give a meaning of the expression

$$\prod_{i \in I} S_i$$

Definition 4.4. Given a countable totally ordered set of indice I and element S_i in a Banach algebra, we say that $\prod_i S_i$ is *well defined* and equal to S if for any increasing chain

$$I_0 \subset I_1 \subset \dots \subset I$$

with $\cup_k I_k = I$ we have $\lim_{k \rightarrow \infty} \prod_{i \in I_k} S_i = S$.

Lemma 4.7. For element s_i in a Banach algebra index by a countable totally ordered set, if $\sum \|s_i\| < \infty$, then $\prod(1 + s_i)$ is well-defined.

Proof. For $1 \leq m \leq n$, we have

$$\left\| \prod_{i=1}^n (1 + s_i) - \prod_{i=1, i \neq m}^n (1 + s_i) \right\| \leq \|s_m\| \left\| \prod_{i=1, i \neq m}^n (1 + s_i) \right\| \leq \|s_m\| \prod_{i=1}^n (1 + \|s_i\|)$$

Or with the assumption $\sum \|s_i\| < \infty$ we have that $\prod(1 + \|s_i\|) \leq C < \infty$, so removing or adding $1 + s_m$ produce a change bound by $\|s_m\|C$. \square

Now we want to apply this lemma to $S_i - Id$, with Id the identity matrice.

Lemma 4.8. For the previous S_i , if we note $s_i = S_i - Id$ we have $\sum \|s_i\| < \infty$.

Proof. Each S_i is conjugate to a horocycle transformation of time one. The conjugacy is made by a geodesic flow along the edges of the triangle. We can compute

$$\begin{pmatrix} e^{-t/2} & 0 \\ 0 & e^{t/2} \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & e^{-t} \\ 0 & 0 \end{pmatrix}$$

So the norm of s_i is inversally corrolated to the amount of geodesic flow used in the conjugaison. Now we can parttion the indice set I into finely many subset (I_k) according to to which spike of the lamination the arc of A cross. Then for a spike the sum of $\|s_i\|$ where $i \in I_k$ is finite, indeed the distance between two neighbooring crossing is bound below by a constant and so the amount of time we should do the geodesic flow increase at most lineraly and finnaly the norm of the s_i should decreaese geometrically. \square

So we can conclude that there is an unique Moebuis transformation S equal to the meaningfull expression $\prod_i S_i$.

Now we can conclude the proof. There exist, without any hyperbolic structure X topological classes for $\tilde{\mu}$ and $\tilde{\lambda}$. We choose one arbitrary ideal triangle T_1 in the lamination. For every other triangle T_2 , the Moeubuis transformation and the shear are data that can be compute only using the transverse measure of $\tilde{m}u$. So we can place T_2 , and the other triangle. The closure of this set give the lamination $\tilde{\lambda}$. $\tilde{\lambda}$ will be preserve by a fushian group Γ and we will have $X = \mathbb{D}/\Gamma$.

4.4 Conjugaision between the horocyclic flow and the earthquake flow

We now have to prove that the map between $\mathcal{ML} \times \mathcal{T}_g$ and \mathcal{QD} conjugate the earthquake flow and the horocyclic flow.

Lemma 4.9. Denote by $Shear_X(T_1, T_2)$ the shear for two triangles joined by an arc A of the horocyclic foliation on the hyperbolic surface X . Then

$$Shear_{E_{t\lambda}(X)}(T_1, T_2) = Shear_X(T_1, T_2) + t\lambda(A)$$

where $\lambda(A)$ denote the transverse measure of A and t is sufficiently small.

Proof. T_1 and T_2 are separated by infinitely many leaves of $\tilde{\lambda}$. We want to understand how T_2 moved relatively to T_1 by the action of the earthquake $E_{t\lambda}$. We can approximate the measured lamination between the two triangles by a discrete one. If the earthquake along a leaf of γ of $\tilde{\lambda}$ between T_1 and T_2 by an amount t , then this changes the shear coordinate between T_1 and T_2 by t . Indeed each arc of the horocycle foliation which have endpoint on γ see their endpoints translated by t . Similarly if the earthquake moves finitely many leaves of λ with measure a_i , the shear changes by precisely $t \sum a_i$. So taking a limit, we have the lemma for an arbitrary measured foliation. \square

We will now show that the Mirzakhani's map conjugates the earthquake flow to the horocyclic flow. Let \mathcal{ML}_0 denote the set of maximal lamination and \mathcal{QD}_0 the locus of quadratic differentials with simple zeros and no horizontal saddle connection. We begin with an easy lemma.

Lemma 4.10. *An arc joining two singularities of a quadratic differential have in its isotopy class a path made of horizontal and vertical arc between two singularities of the quadratic differential.*

Corollaire. *Suppose q_t is a path in the space of quadratic differential such that for every t_0 and every path γ on q_{t_0} joining two singularities, the period $x_t + iy_t$ of γ satisfied*

$$\frac{d}{dt}|_{t=t_0} x_t = y_{t_0}, \frac{d}{dt}|_{t=t_0} y_t = 0$$

Then q_t is an orbit of the horocyclic flow.

Proof. We have

$$\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x + ty \\ y \end{pmatrix}$$

And we can integrate the two equations to find the linear action. \square

We can now prove theorem 4.1.

Proof. We want to show that

$$q(\lambda, F_\lambda(E_{t\lambda}(X)))$$

is a horocyclic flow path. We pick an arbitrary time t_0 and look at the derivative of the path, called γ . The coordinate y_t is constant equal to $\lambda(\gamma)$ and the derivative of x_t is by lemma 4.9 equal to y_t . We conclude by the previous lemma. \square

5 Mixing rate

5.1 Mixing proprieties of the geodesic and horocyclic flows

We will first give the behavior of two flows we described before.

Theorem 5.1. *The ergodic flow and the horocycle flow are mixing.*

[15]

Proof. The step will be in four step, first we will show that the ergodic flow is ergodic, the horocyclic flow is ergodic, the ergodic flow is mixing, finally the horocycle flow is mixing.

Ergodicity flow is ergodic

We will look at the time average of a function $f \in L^2$ which is continuous and compactly support. It will be sufficient since this space is dense

$$F(x) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(g_t x) dt$$

We want to show that F is almost everywhere constant.

Since F depend only of the geodesic $\gamma(a, b)$ which pass at x , if $a, b \in S_\infty$ are the two endpoints of γ , we have $F(a, b)$. Then as two geodesic with the same forward endpoint are asymptotic, F is a quantity that depend only of asymptotic average, we have that F do not depend of a . But we can reverse the argument to show that if we consider the inverse geodesic, $t \rightarrow \infty$, we have that F is also independent of b . Hence F is constant almost everywhere and the geodesic flow is ergodic.

Ergodicity of the horocycle flow

Now if we take $f \in L^2(X)$ a function invariant under the horocycle flow and of mean zero, we want to show that $f = 0$ almost everywhere. Let G^t , H^s and E^r correspond to the operators of the different flows. We have the relation

$$H^s = E^r G^t E^{\pi+r}$$

where $r \rightarrow 0$ as $t \rightarrow \infty$. Since $H^s f = f$, we have for any $T > 0$,

$$f = \frac{1}{T} \int_0^T E^r G^t E^{\pi+r} f dt$$

Thus for every g in $L^2(X)$ we have

$$\langle g, f \rangle = \int_X g \frac{1}{T} \int_0^T E^r G^t E^{\pi+r} f dt$$

As $r \rightarrow 0$ then $t \rightarrow \infty$ we can show by controlling the difference that

$$\langle g, f \rangle = \lim_{t \rightarrow \infty} \int_X g \frac{1}{T} \int_0^T G^t E^\pi f dt$$

Then as we have shown that the geodesic flow is ergodic, we have by Von Neumann ergodic theorem

$$\langle g, f \rangle = \langle g, \int_X E^\pi f \rangle = 0$$

This conclude the proof that the horocycle flow is ergodic.

Mixing of the geodesic flow

We have the relation

$$h^s g^t = g^t h^{sexp(2t)}$$

Let us take $f_0, f_1 \in C_0(X)$, we have for small s

$$\langle f_0, g^t f_1 \rangle \equiv \langle h^{-s} f_0, g^t f_1 \rangle = \langle f_0, h^s g^t f_1 \rangle = \langle f_0, g^t h^{sexp(2t)} f_1 \rangle = \langle g^{-t} f_0, h^{sexp(2t)} f_1 \rangle$$

Now we have for small s ,

$$h^{sexp(2t)} f_1 \equiv \frac{1}{S} \int_0^S h^{exp(2t)s} f_1 ds = F_t$$

and for large t , as the horoclic flow is ergodic

$$F_t = \frac{1}{S} \int_0^S h^{exp(2t)s} f_1 ds \equiv \int_X f_1 = \langle f_1, 1 \rangle$$

So we have

$$\langle f_0, g^t f_1 \rangle \equiv \langle g^{-t} f_0, F_t \rangle \equiv \langle g^{-t} f_0, 1 \rangle \langle f_1, 1 \rangle = \langle f_0, 1 \rangle \langle f_1, 1 \rangle$$

Which is the mixing of the geodesic flow.

Mixing of the horocyclic flow

We use again the relation $h^s = e^r g^t e^{\pi+r}$.

$$\langle h^s f_0, f_1 \rangle = \langle g^t e^{\pi+r} f_0, e^{-r} f_1 \rangle \equiv \langle g^t e^{\pi} f_0, f_1 \rangle$$

for t large (and so r small). By the mixing propriety of the geodesic flow, this quantity converges to $\langle f_0, 1 \rangle \langle f_1, 1 \rangle$, and the horocyclic flow is mixing. □

Then we have this elementary corollary

Corollaire. *The Earthquake flow is also ergodic.*

Proof. With the conjugacie of Mirzharani, the earthquake flow is conjugated to the horocycle flow which is ergodic. This propriety is transmitted. □

5.2 Rate of mixing of this flows

Then we want to know at which rate the mixing of the geodesic and horocyclic flow happend. Ratner show in 1986 [18] that the geodesic flow is exponnetially mixing and the horocyclic flow has a polynomial rate of mixing. She used representation theory.

We will first give some definition before abording the main theorem.

Definition 5.1. Let H be a complexe separable Hilbert space, $U(H)$ the group of all unitary transformation of H onto itself and $T : G \rightarrow U(H)$ a unitary representation. We note $T(g) = T_g \in U(H)$.

An element $v \in H$ is called a C^k -vector for T , $k = 0, 1, \dots, \infty$ if $g \mapsto T_g(v)$ is a C^k -map from G to the Hilbert space.

Remark. The space of C^∞ vector is dense in H .

Definition 5.2. If v is a C^1 -vector of H and X is in the Lie algebra of G , the *Lie derivative* $L_X v$ is

$$L_X v = \lim_{t \rightarrow 0} \frac{T(\exp tX)v - v}{t}$$

The Lie algebra of $SL(2, \mathbb{R})$, is the vector space of 2×2 matrices with zero trace. A basis for this space is

$$W = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, Q = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, V = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Which give the Casimir operator on C^2 vectors of $SL(2, \mathbb{R})$,

$$\Omega_T = (L_v^2 + L_Q^2 - L_W^2)/4$$

If T is irreducible then Ω_T is a scalar multiple of the identity, i.e.

$$\Omega_T v = \lambda v$$

for some $\lambda = \lambda(T) \in \mathbb{R}$ and all C^2 -vectors $v \in H(T)$

If Γ is a lattice of $SL(2, \mathbb{R})$, Ω_T create an operator Δ on $\Gamma \backslash SL(2, \mathbb{R})$. We call $\Lambda(\Delta)$ its spectrum and

$$A(\Gamma) = \Lambda(\Gamma) \cup] - 1/4; 0[\quad B(\Gamma) = \sup A(\Gamma) C(\Gamma) = -1 + \sqrt{1 + 4B(\Gamma)}$$

We can now write the theorem about the decay of correlation.

Theorem 5.2. Let Γ be a lattice in $SL(2, \mathbb{R})$, $M = \Gamma \backslash G$ and T be the regular representation of G on $L^2(M, \mu)$. Let $v, w \in K(T, p)$, $p > 0$, $\langle w, 1 \rangle = 0$ and $B(t) = \langle v, w \circ g_t \rangle$, $C(t) = \langle v, w \circ h_t \rangle$. Then there exist $t_0 = t_0(\Gamma) > 0$ such that for all $|t| \geq t_0$ and some $E, F > 0$

$$|B(t)| \leq E(b(|t|))^{\alpha(p)} |C(t)| \leq F(b(\ln|t|))^{\alpha(p)}$$

where

1. $b(t) = e^{\sigma(\Gamma)}$ if $A(\Gamma) \neq \emptyset$
2. $b(t) = e^{-t}$ if $A(\Gamma) = \emptyset$, $\sup(\Lambda \cap] - \infty; -1/4[) < -1/4$ and $-1/4$ is not an eigenvalue of Ω
3. $b(t) = te^{-t}$ if $A(\Gamma) = \emptyset$, $\sup(\Lambda \cap] - \infty; -1/4[) = -1/4$ or $-1/4$ is an eigenvalue of Ω

and $\alpha(p)$ is

1. 1 if $p \geq 3$
2. $\frac{2p}{2p+1}$ if $2 \leq p < 3$
3. $\frac{2p}{2p+3}$ if $1 \leq p < 2$
4. $\frac{p}{p+3}$ if $0 < p < 1$

On the other we can make the path backward, by learning information on the representation via the mixing rate of the flow it generates. This way is taken in the appendice B of [1].

Definition 5.3. Let G be a locally compact σ -compact group. A continuous unitary representation of G is said to have *almost invariant vectors* if for every $\epsilon > 0$ and for every compact subset $K \subset G$, there exists a unit vector V such that $\|g * v - v\| < \epsilon$ for all $g \in K$.

A unitary action which does not have almost invariant vectors is said to be *isolated from trivial representation*.

If G is a semi-simple Lie group, a representation which is isolated from trivial representation is also said to have a spectral gap.

With this definition, we can write the following theorem.

Proposition 5.3. *Let consider a representation of $SL(2, \mathbb{R})$ by a measure preserving action of automorphisms of a probability space. Let ρ be the representation associate on H the space of the L^2 function with zero averages. Assume there is $\delta \in]0; 1[$ and a dense subset of the subspace of $SO(2, \mathbb{R})$ -invariant function $H' \subset H$ consisting of functions ϕ for which the correlations $\langle \phi, \rho(g_t) \rangle$, $g_t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}$, decay like $O(e^{-\delta t})$. Then ρ is isolated from the trivial representation.*

References

- [1] Artur Avila, Sebastien Gouezel, and Jean-Christophe Yoccoz. Exponential mixing for the Teichmüller flow. *arXiv Mathematics e-prints*, page math/0511614, November 2005.
- [2] Francis Bonahon. The geometry of teichmüller space via geodesic currents. *Inventiones mathematicae*, 92(1):139–162, 1988.
- [3] Peter Buser. *Geometry and spectra of compact Riemann surfaces*. Springer Science & Business Media, 2010.
- [4] Andrew J. Casson and Steven A. Bleiler. *Automorphisms of Surfaces after Nielsen and Thurston*. London Mathematical Society Student Texts. Cambridge University Press, 1988.
- [5] B. Farb and D. Margalit. *A Primer on Mapping Class Groups (PMS-49)*. Princeton Mathematical Series. Princeton University Press, 2011.
- [6] Howard Gardiner, Frederick P.; Masur. Extremal length geometry of teichmüller space. *Complex Variables and Elliptic Equations*, 16, 4 1991.
- [7] John Hubbard and Howard Masur. Quadratic differentials and foliations. *Acta Mathematica*, 142:221–274, 01 1979.
- [8] John H. Hubbard. *Teichmüller Theory and Applications to Geometry, Topology, and Dynamics*, volume 2 of *Surface Homeomorphisms and Rational Functions*. Matrix Editions, April 2016.
- [9] M. Kapovich. *Hyperbolic Manifolds and Discrete Groups*. Modern Birkhäuser Classics. Birkhäuser Boston, 2009.
- [10] Steven P. Kerckhoff. The nielsen realization problem. *Annals of Mathematics*, 117(2):235–265, 1983.
- [11] François Labourie and Ser Peow Tan. The probabilistic nature of McShane’s identity: planar tree coding of simple loops. 25 pages, December 2017.
- [12] Gilbert Levitt. Foliations and laminations on hyperbolic surfaces. *Topology*, 22(2):119–135, 1983.
- [13] W. B. R. Lickorish. A finite set of generators for the homeotopy group of a 2-manifold. 1964.
- [14] Bruno Martelli. An introduction to geometric topology. 2016.
- [15] Curt McMullen. Hyperbolic manifolds , discrete groups and ergodic theory course notes. 2017.
- [16] Curt McMullen. Riemann surfaces, dynamics and geometry course notes. 2018.
- [17] Greg Mcshane. Simple geodesics and a series constant over teichmüller space. *Invent. Math.*, 132:607–632, 1995.
- [18] Marina Ratner. The rate of mixing for geodesic and horocycle flows. *Ergodic Theory and Dynamical Systems*, 7(2):267–288, 1987.

- [19] Alex Wright. Mirzakhani's work on earthquake flow. *arXiv preprint arXiv:1810.07571*, 2018.
- [20] Alex Wright. A tour through mirzakhani's work on moduli spaces of riemann surfaces. *Bulletin of the American Mathematical Society*, 2020.