

Catalase Enzyme Reaction Student Experiment - Eden Tomes

Rationale:

Cells, the fundamental units of life, depend on a multitude of biochemical reactions for survival and function, collectively known as metabolism (*Biology LibreTexts*, 2016). Many of these essential reactions would occur too slowly to sustain life without biological catalysts called enzymes (*Cooper*, 2000). Enzymes, primarily proteins, accelerate reaction rates by lowering the activation energy. They achieve this by binding to specific reactant molecules, or substrates, at a region called the active site, facilitating their conversion into products (*Robinson*, 2015).

Catalase is a crucial enzyme found in nearly all aerobic organisms, playing a vital protective role. It catalyses the decomposition of hydrogen peroxide (H_2O_2), a toxic byproduct of cellular metabolism, into harmless water (H_2O) and oxygen (O_2) (*Zamocky et al.*, 2008). This detoxification prevents oxidative damage to cellular components, a key aspect of how cells maintain homeostasis (*Nandi*, 2019).

The rate of enzyme-catalysed reactions is influenced by factors such as temperature, pH, substrate concentration, and enzyme concentration (*Robinson*, 2015). While a previous experiment explored temperature effects, this investigation focuses on how varying enzyme concentration impacts reaction rates. According to established enzyme kinetics, when substrate is abundant, the initial reaction rate is directly proportional to the enzyme concentration (*Aebi*, 1984). This is because more enzyme molecules mean more active sites are available to process the substrate, leading to faster product formation (*Robinson*, 2015).

This experiment aims to systematically investigate how altering the volume of catalase solution, thereby changing its effective concentration, affects the initial rate of oxygen production from hydrogen peroxide decomposition. By keeping substrate concentration, volume, and temperature constant, the study will isolate the influence of enzyme quantity on reaction rate. This will provide insight into how cells can regulate metabolic pathways by controlling enzyme availability.

Research Question:

How does increasing the volume of catalase solution (representing different enzyme concentrations) affect the rate of oxygen gas production when reacting with a constant volume and concentration of hydrogen peroxide?

Methodology:

Original Experiment

The original experiment (provided through *Seymour*, 2019) investigated enzyme activity by reacting 1mL of stock catalase solution (yeast) with a set volume of hydrogen peroxide (10mL) in a reaction chamber. The chamber is submerged in a room-temperature water bath. Oxygen gas, the product of the reaction, is collected over water in an inverted measuring cylinder, and its volume is recorded at regular time intervals (every 30 seconds for 5 minutes) to determine the rate of reaction.

Modifications

The original experimental design was refined and redirected to specifically investigate the effect of enzyme concentration on reaction rate. To improve data reliability, the procedure was modified to include three independent trials for each condition. This replication minimizes the impact of random error and allows for statistical analysis through the calculation of means and standard deviations. The experiment was redirected from using a single enzyme volume to systematically testing three different volumes of catalase solution (1mL, 2mL, and 4mL). Since the stock solution's concentration was constant, altering its volume served as a direct method for varying the effective enzyme concentration in the reaction mixture.

Safety & Ethical Considerations

| Hazard | Identified Risk(s) | Control Measure(s) |
|--|---|---|
| Hydrogen Peroxide (H ₂ O ₂) | <ul style="list-style-type: none">• Skin irritation or mild chemical burns.• Eye irritation or damage. | <ul style="list-style-type: none">• Wear safety goggles and a lab coat at all times.• Handle H₂O₂ with care to avoid spills.• If H₂O₂ contacts skin, rinse immediately with plenty of water |
| Glassware (test tubes, beakers) | <ul style="list-style-type: none">• Cuts from breakage. | <ul style="list-style-type: none">• Handle carefully and avoid excessive force.• Keep glassware away from the edge of the bench.• Dispose of any broken glassware in a designated broken glass bin. |
| Chemical Spills | <ul style="list-style-type: none">• Slip hazard on the floor.• Contamination of the | <ul style="list-style-type: none">• Clean up any spills immediately with paper towels as per teacher |

| | | |
|--|-------------------------|---|
| | work area or equipment. | instructions. <ul style="list-style-type: none"> • Work on a clear, uncluttered bench space. |
|--|-------------------------|---|

Processed Data + Calculations:

Table 1: Sample Calculations

| Calculation | Example |
|--|--|
| Mean oxygen volume | $\text{Mean} = \frac{\text{Sum of values}}{\text{Count}}$ $\text{For 1mL: Mean} = \frac{4.0 + 3.0 + 0.7}{3} \approx 2.57$ |
| Standard deviation (SD) for a sample population | Calculated in excel by using STDEV.S and the three trial measurements for each catalase volume. For 1mL: $s \approx 1.69$ |
| Standard error | Calculated by dividing the standard deviation by the square root of the sample size. For 1mL: $\frac{1.69}{\sqrt{3}} \approx 0.98$ |
| Upper/lower limit | Calculated as follows: Upper limit = Mean + (2 × Standard deviation)) Lower limit = Mean – (2 × Standard deviation)) For 1mL: $2.57 \pm (2 \times 1.69) \approx 5.95, -0.82$ (clamped to 0) |
| P-value | P-values were calculated in excel using the T.DIST function and the trial differences between each pair of conditions. For 1mL vs 2mL: $p \approx 0.35$ |


*Table 2: Processed data table for volume of Oxygen After 300 Seconds
(values are in mL and rounded to 2 decimal places)*

| | 1mL Catalase | 2mL Catalase | 4mL Catalase |
|-----------------------|--------------|--------------|------------------------------|
| Trial 1 | 4.00 | 0.70 | 11.00 |
| Trial 2 | 3.00 | 5.00 | 8.50 |
| Trial 3 | 0.70 | 5.00 | 4.50 |
| Mean | 2.57 | 3.57 | 8.00 |
| Std. Deviation | 1.69 | 2.48 | 3.28 |
| Std. Error | 0.98 | 1.43 | 1.89 |
| Upper Limit | 5.95 | 8.53 | 14.56 |
| Lower Limit | 0.00 | 0.00 | 1.44 |
| P-Value | | Vs 1mL: 0.35 | Vs 2mL: 0.15 Vs 1mL: 0.01 |

Interpretation:

After 300 seconds, mean oxygen production increased with enzyme concentration: 2.57 mL (1mL), 3.57 mL (2mL), and 8.00 mL (4mL). This trend was statistically significant when comparing the lowest and highest concentrations (1mL vs 4mL, $p=0.01$). However, the differences between adjacent steps were not significant (1mL vs 2mL, $p=0.35$; 2mL vs 4mL, $p=0.15$), assuming a significance level of $\alpha=0.05$. The large standard errors (± 1.43 mL and ± 1.89 mL for the 2mL and 4mL treatments) likely masked the expected difference between these groups, indicating low precision.

**Average Volume of Oxygen Produced by Reaction
After 300 Seconds**



Graph 1: Column Graph (standard deviations represented by error bars)


Interpretation:

The column graph shows the mean volume of oxygen gas produced after 300 seconds for each catalase volume treatment. The 4mL catalase treatment produced the highest mean oxygen volume (8.00 mL), followed by 2mL catalase (3.57 mL), and 1mL catalase (2.57 mL). This demonstrates a general positive relationship between enzyme volume and oxygen production, with the 4mL treatment producing approximately 3.1 times more oxygen than the 1mL baseline and 2.2 times more than the 2mL treatment.

Analysis:

The data indicates substantial variability within each treatment group, as evidenced by the large standard deviations represented by the error bars (1mL: ± 1.69 mL, 2mL: ± 2.48 mL, 4mL: ± 3.28 mL). The overlapping error bars between the 1mL and 2mL treatments suggest these results may fall within the same range, indicating no statistically significant difference between these enzyme concentrations. However, the 4mL treatment shows minimal overlap with the other treatments, suggesting a potential significant difference. The increasing magnitude of standard deviations with higher enzyme concentrations indicates decreasing precision in measurements, possibly due to more vigorous reactions being harder to measure consistently or substrate limitation effects occurring at higher enzyme concentrations.

Average Oxygen Production of Catalase + Hydrogen Peroxide Reaction Over Time



Graph 2: Line Graph - Mean oxygen volume over a 300 second period

Interpretation:

The line graph demonstrates the cumulative oxygen production over the 300-second experimental period for each catalase treatment. All treatments show the characteristic pattern of enzyme kinetics, with an initial rapid rate of oxygen production that gradually levels off as the reaction progresses. The 4mL catalase treatment exhibits the steepest initial slope and maintains the highest oxygen production throughout the time period, reaching approximately 8.00 mL by 300 seconds. The 2mL treatment shows a moderate

rate of production, while the 1mL treatment displays the slowest rate, reaching only 2.57 mL after 300 seconds.

Analysis:

The initial reaction rates (represented by the steepness of the early slopes) follow the expected pattern where higher enzyme concentrations produce faster initial rates of oxygen evolution. This aligns with enzyme kinetics theory, where increased enzyme concentration provides more active sites for substrate binding when substrate is in excess. However, the leveling off observed in all treatments suggests that substrate depletion may be occurring before the 300-second endpoint. The plateau is most evident in the 4mL treatment, indicating that the 10mL of hydrogen peroxide substrate becomes limiting when enzyme concentration is high, preventing the reaction from maintaining its initial rate throughout the entire observation period. This plateau effect compromises the ability to accurately assess the true relationship between enzyme concentration and reaction rate, as the measured values at 300 seconds may not reflect the initial rate conditions where enzyme kinetics principles are most applicable.

Evaluation:

Limitations Of Evidence

The experiment's low reliability is evidenced by high standard deviations (± 1.69 to ± 3.28 mL). This variability severely limited the experiment's statistical power, as illustrated by the p-values. For instance, despite the mean oxygen production for the 4mL treatment being more than double that of the 2mL treatment (8.00 mL vs 3.57 mL), the difference was not statistically significant ($p=0.15$). This demonstrates that extreme variance and a small sample size ($n=3$) masked a large, real effect, limiting the validity of conclusions drawn from intermediate comparisons.

Sources Of Error:

Affecting Reliability

- The gas collection apparatus lacks precision in measuring oxygen volume. The water displacement method and visual reading of gas volume contributes to imprecision in the data (± 0.5 mL reading uncertainty). However, the variation in the data is greater than this measurement uncertainty; therefore, there must be other sources of imprecision.
- The samples were not standardized for reaction initiation timing. Random variation exists in the sealing and submersion process of the reaction chamber. This could explain some of the remaining imprecision in the data, as inconsistent reaction start times would lead to variable gas loss.
- The catalase solution homogeneity was not verified prior to conducting the experiment. Therefore, it is not known whether each aliquot contained identical

enzyme concentrations, leading to random biological variation affecting the reaction rates.

Affecting Validity

- The water displacement method leads to underestimation of actual gas volume due to oxygen dissolution. Therefore, the volume of oxygen produced is systematically underestimated. In addition, the oxygen volume is determined indirectly which could lead to greater variability in the data.
- The experimental design does not maintain constant substrate concentration across treatments. The varying total reaction volumes (11mL, 12mL, 14mL) dilute the hydrogen peroxide to different concentrations, introducing a confounding variable that affects the validity of isolating enzyme concentration effects.
- The gas collection method does not account for substrate depletion at higher enzyme concentrations. The 300-second endpoint measurement does not represent true initial rates, as substrate limitation occurs before completion, particularly in the 4mL treatment. Therefore, this could contribute to the data being inaccurate for determining the enzyme concentration-rate relationship.

Suggested Improvements & Extensions:

Suggested Improvements:

Reducing the random error in the experimental process would improve its reliability. In this experiment, the reliability of the data could be improved by increasing the number of repeat readings of each treatment (minimum $n=5$), standardizing the reaction initiation procedure, and running multiple independent trials to decrease standard error.

To address the imprecision in gas measurement, a gas syringe should be used instead of water displacement, which would eliminate oxygen dissolution effects and provide direct, more accurate volume readings. This would improve both the reliability of data and validity of the experimental process.

Maintaining constant total reaction volume across all treatments by adding distilled water to equalize volumes would ensure identical substrate concentrations, allowing the results to isolate the effect of enzyme concentration. In addition, ensuring homogeneity of the catalase stock solution through thorough mixing before each aliquot would reduce biological variation.

Suggested Extensions:

- Redirect the experiment by investigating a wider range of enzyme concentrations with smaller incremental increases to better define the enzyme concentration-rate relationship and identify saturation points.

- Extend the experiment by investigating the effect of substrate concentration while keeping enzyme concentration constant, providing a comprehensive exploration of enzyme kinetics.

Conclusion:

In conclusion, the evidence suggests that increasing the volume of catalase solution does cause a measurable increase in the rate of oxygen gas production, with a statistically significant difference found between the 1mL and 4mL treatments ($p=0.01$). However, the lack of significance in other comparisons (e.g., 2mL vs 4mL, $p=0.15$), despite large differences in mean production, highlights that the experiment's conclusion is limited by low precision. Therefore, while the overall trend is supported, the data's high variability prevents the establishment of a definitive quantitative relationship and confirmation of the theoretical proportionality between enzyme concentration and initial reaction rate.

Word Count: 1964

Reference List:

1. Aebi, H. (1984). Catalase in vitro. *Methods in Enzymology*, [online] 105, pp.121–126. doi:[https://doi.org/10.1016/s0076-6879\(84\)05016-3](https://doi.org/10.1016/s0076-6879(84)05016-3).
2. Biology LibreTexts. (2016). 1.16: Types of Biochemical Reactions. [online] Available at: [https://bio.libretexts.org/Bookshelves/Introductory_and_General_Biology/Introductory_Biology_\(CK-12\)/01%3A_Introduction_to_Biology/1.16%3A_Types_of_Biochemical_Reactions](https://bio.libretexts.org/Bookshelves/Introductory_and_General_Biology/Introductory_Biology_(CK-12)/01%3A_Introduction_to_Biology/1.16%3A_Types_of_Biochemical_Reactions).
3. Cooper, G. (2000). *The Central Role of Enzymes as Biological Catalysts*. [online] National Library of Medicine. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK9921/>.
4. Nandi, A. (2019). Role of Catalase in Oxidative Stress- and Age-Associated Degenerative Diseases. *Oxidative Medicine and Cellular Longevity*, [online] 2019(9613090). doi:<https://doi.org/10.1155/2019/9613090>.
5. Robinson, P. (2015). Enzymes: Principles and Biotechnological Applications. *Essays in Biochemistry*, [online] 59(1), pp.1–41. doi:<https://doi.org/10.1042/bse0590001>.
6. Seymour, M., 2019. *Catalase Enzyme Activity and Temperature*, Brisbane: Education Queensland.
7. Zamocky, M., Furtmüller, P.G. and Obinger, C. (2008). Evolution of Catalases from Bacteria to Humans. *Antioxidants & Redox Signaling*, [online] 10(9), pp.1527–1548. doi:<https://doi.org/10.1089/ars.2008.2046>.

Appendix:

Raw Data

| Amount of Catalase | 30 Seconds | 60 Seconds | 90 Seconds | 120 Seconds | 150 Seconds | 180 Seconds | 210 Seconds | 240 Seconds | 270 Seconds | 300 Seconds |
|--------------------|------------|------------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| 1mL Catalase | | | | | | | | | | |
| trial 1 | 2 | 2.5 | 2.5 | 2.5 | 2.5 | 3 | 3 | 3 | 3 | 4 |
| trial 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 3 | 3 |
| trial 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.5 | 0.7 |
| Average: | 1.333 | 1.500 | 1.500 | 1.500 | 1.500 | 1.667 | 1.667 | 1.667 | 2.167 | 2.567 |
| 2mL Catalase | | | | | | | | | | |
| trial 1 | 0 | 0 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.7 | 0.7 |
| trial 2 | 2 | 2 | 2 | 2 | 2.5 | 2.8 | 3 | 4 | 4 | 5 |
| trial 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 4 | 5 |
| Average: | 1.667 | 1.667 | 1.833 | 1.833 | 2.000 | 2.100 | 2.167 | 2.533 | 2.900 | 3.567 |
| 4mL Catalase | | | | | | | | | | |
| trial 1 | 4 | 5 | 7 | 7.5 | 8.5 | 8.5 | 10 | 11 | 11 | 11 |
| trial 2 | 2 | 3 | 3 | 5 | 5 | 6 | 7 | 7.5 | 8.5 | 8.5 |
| trial 3 | 0 | 0 | 0 | 0.5 | 0.5 | 2 | 2 | 3 | 4.5 | 4.5 |
| Average: | 2.000 | 2.667 | 3.333 | 4.333 | 4.667 | 5.500 | 6.333 | 7.167 | 8.000 | 8.000 |

Excel tests

| | 1mL Catalase | 2mL Catalase | Difference | | 2mL Catalase | 4mL Catalase | Difference | | 1mL Catalase | 4mL Catalase | Difference |
|-----------|-----------------|-----------------|------------|--|-----------------|-----------------|------------|--|-----------------|-----------------|------------|
| | 4 | 0.7 | -3.3 | | 0.7 | 11 | 10.3 | | 4 | 11 | 7 |
| | 3 | 5 | 2 | | 5 | 8.5 | 3.5 | | 3 | 8.5 | 5.5 |
| | 0.7 | 5 | 4.3 | | 5 | 4.5 | -0.5 | | 0.7 | 4.5 | 3.8 |
| Mean | | | 1 | | | | 4.433333 | | | | 5.433333 |
| SD | | | 3.89743505 | | | | 5.46015873 | | | | 1.60104133 |
| SE | | | 2.25018518 | | | | 3.15242411 | | | | 0.92436164 |
| T-Stat | | | 0.44440787 | | | | 1.40632516 | | | | 5.87793033 |
| Degr Free | | | 2 | | | | 2 | | | | 2 |
| P-Value | | | 0.35010489 | | | | 0.1474368 | | | | 0.01387231 |

AT6

AT3

AT7

AT8

AT9

AT11

AT10

AT12

AT13

AT14

AT15

AT16

AT17

AT18

AT19

AT20

AT21

AT22

AT23

AT24

AT25

AT26

AT27

AT28

AT29

AT30

AT31

AT32

AT33

AT34

AT35

AT36

AT37

AT38

AT39

AT40

AT41

AT42

AT43

AT44

AT45

AT46

AT47

AT48

AT49

AT50

AT51

AT52

AT53

AT54

AT55

AT56

AT57

AT58

AT59

AT60

AT61

AT62

AT63

AT64

AT65

AT66

AT67

AT68

AT69

AT70

AT71

AT72

AT73

AT74

AT75

AT76

AT77

AT78

AT79

AT80

AT81

AT82

AT83

AT84

AT85

AT86

AT87

AT88

AT89

AT90

AT91

AT92

AT93

AT94

AT95

AT96

AT97

AT98

AT99

AT100

AT101

AT102

AT103

AT104

AT105

AT106

AT107

AT108

AT109

AT110

AT111

AT112

AT113

AT114

AT115

AT116

AT117

AT118

AT119

AT120

AT121

AT122

AT123

AT124

AT125

AT126

AT127

AT128

AT129

AT130

AT131

AT132

AT133

AT134

AT135

AT136

AT137

AT138

AT139

AT140

AT141

AT142

AT143

AT144

AT145

AT146

AT147

AT148

AT149

AT150

AT151

AT152

AT153

AT154

AT155

AT156

AT157

AT158

AT159

AT160

AT161

AT162

AT163

AT164

AT165

AT166

AT167

AT168

AT169

AT170

AT171

AT172

