

0.1 Calculus cheat sheet

Logs: $\log_b(M * N) = \log_b M + \log_b N$ • $\log_b(\frac{M}{N}) = \log_b M - \log_b N$ • $\log_b(M^k) = k \log_b M$ • $e^n e^m = e^{n+m}$

Derivatives: $(x^n)' = nx^{n-1}$ • $(e^x)' = e^x$ • $(e^{u(x)})' = u'(x)e^x$ • $(\log_e(x))' = (\ln x)' = \frac{1}{x}$ • $(f(g(x)))' = f'(g(x))g'(x)$

Integrals: $\int_a^b f(x)dx = \int_{g(a)}^{g(b)} f(g(u))g'(u)du$ where $g(u) = x$ • $\int_a^b u(x)v'(x)dx = u(b)v(b) - u(a)v(a) - \int_a^b u'(x)v(x)dx$

Infinite series and sums: $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ • $(1 + \frac{a}{n})^n \rightarrow e^a$

• $\ln(1+x) = 1 - x + \frac{x^2}{2} - \frac{x^3}{3} + \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{n}$ • $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots = \sum_{n=0}^{\infty} x^n$ for $|x| < 1$

L'Hopitale: $\lim_{n \rightarrow c} f(x)/g(x) = \lim_{n \rightarrow c} f'(x)/g'(x)$ if $\lim_{n \rightarrow c} f(x) = \lim_{n \rightarrow c} g(x) = 0/\infty/-\infty$

0.2 Expectation

Conditional expectation: $p_{X|Y}(x|y) = \frac{p_{x,y}(x,y)}{p_y(y)}$

Bayes theorem: $P(E_i | B) = \frac{P(B|E_i)P(E_i)}{\sum_{j=1}^{\infty} P(B|E_j)P(E_j)} = \frac{P(B|E_i)P(E_i)}{P(B)}$, where E_1, E_2, \dots form a partition of the sample space.

Expectation: $E(X) = \sum_x xP(X=x)$, also written $E(X) = \sum_{s \in S} X(s)p(s)$, where $p(s)$ is the probability that element $s \in S$

• $E(g(X)) = \sum_i g(x_i)p_X(x_i)$ • $E(aX+b) = aE(X)+b$ • $E(X+Y) = E(X)+E(Y)$

• $P_{Y,X}(Y > X) = E_{Y,X}[\mathbb{I}\{Y > X\}] = E_X E_Y[\mathbb{I}\{Y > X\} | X] = E_X P_Y(Y > X | X) = \int_{\mathbb{R}} P_Y(Y > x | X=x)F_X(dx)$

Variance: $Var(X) = E((X - E(X))^2) = \sigma^2$

• $Var(X) = E(X^2) - \mu^2$ • $Var(aX+b) = a^2 Var(X)$ • $Var(X+Y) = Var(X) + Var(Y) + 2Cov(X,Y)$

Covariance: $Cov(X,Y) = E((X - E(X))(Y - E(Y))) = E(XY) - E(X)E(Y)$

• $Cov(X,X) = Var(X)$ • $Cov(aX,bY) = abCov(X,Y)$ • $Cov(X,Y+Z) = Cov(X,Y) + Cov(X,Z)$

Law of iterated expectation: $E(E(Y | X)) = E(Y)$

Proof: $E(Y | X) = \sum_y y \frac{f_{X,Y}(X,y)}{f_X(X)}$, $E(E(Y | X)) = \sum_x \sum_y \left(y \frac{f_{X,Y}(x,y)}{f_X(x)} \right) f_X(x) = \sum_x \sum_y y f_{X,Y}(x,y) = \sum_y y f_Y(y) = E(Y)$

Law of total probability: $P(E) = \sum_{i=-\infty}^{\infty} P(E | X=x)P(X)$ and $P(E) = \int_{-\infty}^{\infty} P(E | X=x)f(x)dx$

Variance decomposition formula: $Var(Y) = E(Var(Y | X)) + Var(E(Y | X))$

Cauchy-Schwartz inequality: $E(UV)^2 \leq E(U^2)E(V^2)$, with equality if $P(cU=V) = 1$ for some constant, c

Transformations of random variables: For X with density f_X and $Y = g(X)$

$F_Y(y) = P(g(X) \leq y) = P(X \leq g^{-1}(y)) = F_X(g^{-1}(y))$ • $f_Y(y) = \frac{d}{dy} F_Y(g^{-1}(y)) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|$

Stationarity: $\{X_1, \dots, X_n\} \stackrel{d}{=} \{X_{m+1}, \dots, X_{m+n}\}$ • $EX_1 = EX_n$ • $Cov(X_1, X_n) = Cov(X_{m+1}, X_{m+n})$, called $c(n)$ where n is lag. We can prove $c(n) \xrightarrow{p} 0$ using Chebychev and solving for $Var \bar{X}_n$ as a function of $c(n)$: $P(|\bar{X}_n - EX_1| > \epsilon) \leq \frac{2}{n} \sum_i c(i) \rightarrow 0$

0.3 Inequalities

Jensen inequality: $E(g(x)) \geq g(E(x))$ for $g(x)$ convex

Markov inequality: For $X \geq 0$, $P(X \geq t) \leq \frac{E(X)}{t} \quad \forall t > 0$. **Proof:**

Let $Y = \begin{cases} 1 & X \geq t \\ 0 & \text{else} \end{cases}$, then $tY \leq X$ since $\begin{cases} X \geq t & t1 \leq X \\ X < t & t0 < X \end{cases} \implies E(tY) \leq E(X) \implies tP(X \geq t) \leq E(X) \implies P(X \geq t) \leq \frac{E(X)}{t}$

Chebyshev inequality: $P(|X - E(X)| \geq t) \leq \frac{Var(X)}{t^2} \quad \forall t > 0$.

Proof: $P(|X - E(X)| \geq t) = P((X - E(X))^2 \geq t^2) \leq \frac{E((X - E(X))^2)}{t^2} = \frac{Var(X)}{t^2}$, by Markov inequality

Exponential inequality: $P(X > a) \leq e^{-\theta a} E(e^{\theta X})$ for all $\theta > 0$.

Proof: $P(X > a) = P(\theta X > \theta a)$ for $\theta > 0 \implies P(e^{\theta X} > e^{\theta a}) \leq e^{-\theta a} E(e^{\theta X})$, by Markov inequality

(Corrolary) Upper bound on large deviations: $P(S_n < na) \leq e^{-nI(a)}$.

Proof: $P(S_n > a) \leq e^{-\theta a} E(e^{\theta S_n}) = e^{-\theta a} \prod_i E(e^{\theta X_i}) = e^{-\theta a} E(e^{\theta X_1})^n = e^{-\theta a + n\psi(\theta)}$, by exponential inequality, iid, where $\psi(\theta) = \log Ee^{\theta X_1}$
 $P(S_n > na) = e^{-n(\theta(x)a - n\psi(\theta(x)))}$; minimizing RHS w.r.t $\theta \implies e^{-nI(x)}$ where $I(x) = \theta(x)a - n\psi(\theta(x))$

0.4 Generative functions

• $\phi_X(t) = E \exp(itX)$ • $N(\mu, \sigma^2) : \exp(it\mu - \frac{1}{2}\sigma^2 t^2)$ • $Exp(\lambda) : (1 - it\lambda^{-1})^{-1}$ • $Poisson(\lambda) : \exp \lambda (e^{it} - 1)$ • $E[X^k] = i^{-k} E[X^k]$
 • $\phi_{a_1 X_1 + \dots + a_n X_n}(t) = \phi_{X_1}(a_1 t) \dots \phi_{X_n}(A_n t)$ for X_i independent • $M_X(t) = E \exp(tX)$

0.5 Weak law of large numbers

For X_1, X_2, \dots, X_n i.i.d. with $E(X_i) = \mu$, $Var(X_i) = \sigma^2$, $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$, then for any $\epsilon > 0$

$$P(|\bar{X}_n - \mu| > \epsilon) \leq \frac{Var(\bar{X}_n)}{\epsilon^2} = \frac{\sigma^2}{n\epsilon^2} \rightarrow 0 \text{ as } n \rightarrow \infty, \text{ by Chebyshev inequality, } X_1 + \dots + X_n = S_n \approx ES_n, \text{ the "meta result"}$$

0.5.1 Convergence in probability

Convergence in probability: $X_n \xrightarrow{p} X$ when $P(|X_n - X| > \epsilon) \rightarrow 0$ as $n \rightarrow \infty$

Continuous mapping theorem: if $X_n \xrightarrow{p} X$ and g a continuous function then $g(X_n) \xrightarrow{p} g(X)$

Consistent estimator: $T_n = T_n(X_1, \dots, X_n)$ converges in probability to $g(\theta)$, a function of the model parameter

Bounded convergence theorem: $Z_n \xrightarrow{p} Z_\infty, |Z_n| \leq c < \infty \implies EZ_n \xrightarrow{p} EZ_\infty$

Proof starts with $|E(Z_n - Z_\infty)|$ and uses i) triangle inequality, ii) indicator functions for the case when difference is $> \epsilon, < \epsilon$

Dominated convergence theorem: $Z_n \xrightarrow{p} Z_\infty, E\beta < \infty, |Z_n(\omega)| \leq \beta(\omega) \forall \omega \implies EZ_n \xrightarrow{p} EZ_\infty$

Fatous Lemma: for $Z_n > 0$, $E \lim_{n \rightarrow \infty} Z_n \leq \lim_{n \rightarrow \infty} EZ_n$

Almost sure convergence: $P(\omega : \lim_{n \rightarrow \infty} X_n(\omega) \xrightarrow{p} X_\infty(\omega)) = 1$ where ω is element in set of all sequences • $P(\limsup_{n \rightarrow \infty} \{|X_n - X_\infty|\} > \epsilon) = 0$

0.6 Central limit theorem

$$\sqrt{n} \frac{(\bar{X}_n - \mu)}{\sigma} \rightarrow N(0, 1) \iff \sqrt{n}(\bar{X}_n - \mu) \rightarrow N(0, \sigma^2) = \sigma N(0, 1), \quad X_i + \dots + X_n = S_n \approx N(ES_n, Var S_n), \text{ the "meta result"}$$

$$\lim_{n \rightarrow \infty} P\left(\frac{S_n}{\sigma\sqrt{n}} \leq x\right) = \Phi(x), \text{ for } S_n = \sum_{i=1}^n X_i, \quad X_1, X_2, \dots, X_n \text{ i.i.d. with } E(X_i) = 0 \text{ (WLOG), } Var(X_i) = \sigma^2$$

Proof sketch: Start with $M_{S_n}(t)$, plug in $t/(\sigma\sqrt{n})$, use Taylor expansion to show convergence to MGF of a normal, $e^{\frac{t^2}{2}}$

Monte Carlo: • Sample $Y \in \mathbb{R}^d$ • Compute $X = g(Y)$ • Repeat n times • form \bar{X}_n and use CLT for asymptotic behavior.

Generating random data: with $X = F_X^{-1}(U)$ since $P(F_X^{-1}(U) \leq x) = P(F_X(F_X^{-1}(U)) \leq F_X(x)) = P(U \leq F_X(x)) = F_X(x)$

0.6.1 Delta method

If g is a differentiable function at μ , $\sqrt{n}(g(\bar{X}_n) - g(\mu)) \xrightarrow{d} N(0, g'(\mu)^2 \sigma^2) = g'(\mu) \sigma N(0, 1)$. **Proof sketch:** Start with Taylor expansion $g(\bar{X}_n) \approx g(\mu) + g'(\mu)(\bar{X}_n - \mu)$, rearrange to get $\sqrt{n}(g(\bar{X}_n) - g(\mu)) \approx g'(\mu) \sqrt{n}(\bar{X}_n - \mu) \xrightarrow{d} N(0, g'(\mu)^2 \sigma^2)$. **Note:** if we find $g'(\mu) = 0$, then repeat this process with the second derivative, $g''(\mu)$.

0.6.2 Convergence in distribution (a.k.a. weak convergence)

Convergence in distribution: $X_n \xrightarrow{d} X$ when $\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x)$ at all continuity points in F_X

Equalities: $X_n \xrightarrow{d} X \iff Eh(Z_n) \xrightarrow{p} Eh(Z_\infty) \forall h$, bounded/continuous $\iff \phi_{Z_n}(t) \xrightarrow{p} \phi_{Z_\infty}(t) \forall t$

Confidence intervals: $P(Z_{\alpha/2} \leq Z \leq Z_{1-\alpha/2}) = P(\frac{\hat{\sigma}}{\sqrt{n}} Z_{\alpha/2} \leq \bar{X}_n - \mu \leq \frac{\hat{\sigma}}{\sqrt{n}} Z_{1-\alpha/2}) = P(\mu \in [\bar{X}_n \pm \frac{\hat{\sigma}}{\sqrt{n}} Z_{1-\alpha/2}]) = 1 - \alpha$ for $\hat{\sigma} \xrightarrow{p} \sigma$

Slutsky's lemma $A_n X_n + B_n \xrightarrow{d} aX + b$ if $\{X_n\}$ sequence, $X_n \xrightarrow{d} X$, $\{A_n\}$ sequence, $A_n \xrightarrow{d} A$, $\{B_n\}$ sequence, $B_n \xrightarrow{p} b$

0.7 Theory of large deviations

Variance reduction: $E_p h(X)$ where $X \sim P = \int_{-\infty}^{\infty} h(x)p(x)dx = \int_{-\infty}^{\infty} h(x)p(x)q(x)/q(x) = E_q[h(x)p(x)/q(x)]$ where $X \sim Q$.

$\sqrt{n}(\bar{X} - E_p X) \xrightarrow{d} N(0, Var_p(X))$, $\sqrt{n}(\bar{X} - E_q \hat{X}) \xrightarrow{d} N(0, Var_q(\hat{X}))$ where $\hat{X} = Xp(x)/q(X)$

Importance sampling: Choose $h(x)$ to minimize variance. Minimal $H(dx)$ turns out to be the conditional probability of the event happening on event happening: $H^*(dx) = \mathbb{I}\{A\}(x)F(dx)/F(A)$

Exponential tilting: $E f(X_1, \dots, X_n) = E_\theta \exp(-\theta S_n + n\psi(\theta)) f(X_1, \dots, X_n)$, where $\psi(\theta) = \log M_x(\theta)$

Large deviations: • Use exponential tilting with $f(X_1, \dots, X_n) = \mathbb{I}(S_n > an)$: $E \mathbb{I}(S_n > an) = E_\theta [\exp(-\theta S_n + n\psi(\theta)) \mathbb{I}(S_n > an)]$ • Choose optimal $\theta^* = \theta(a)$ which satisfies $\psi'(\theta(a)) = a$, which guarantees $E_{\theta(a)} X_i = a$ • E.g., Gaussian: $M_X(\theta) = \exp(\mu\theta + \sigma^2\theta^2/2) \iff \psi(\theta) = \mu\theta + \sigma^2\theta^2/2 \iff \psi'(\theta) = \mu + \sigma^2\theta \leftarrow$ evaluate at $\psi'(\theta) = a$

0.8 Method of moments estimator

• $E(X^k) = g(\theta)$: Calculate moment with MGF, lower moments typically lead to estimators with lower asymptotic variance • $g^{-1}(E(X^k)) = \theta$: Invert this expression to create an expression for the parameter(s) in terms of the moment • $\hat{\theta} = g^{-1}(\frac{1}{n} \sum X_i^k)$: Insert the sample moment into this expression, thus obtaining estimates of the parameters in terms of data • $\sqrt{n}(g^{-1}(\frac{1}{n} \sum X_i^k) - \theta) \xrightarrow{d} N(0, f'(E(X_i^k))^2 Var(X_i^k)^2)$: Use the delta method • If multiple parameters characterize the distribution, use multiple moments and a system of equations

0.9 Maximum likelihood estimator

• $L(\theta) = \prod_{i=1}^n f(X_i, \theta)$: Construct the likelihood function • $\log(L(\theta)) = l(\theta) = \sum_{i=1}^n \log(f(X_i, \theta))$: Take the log of the likelihood • Find critical points of this function (e.g., $0 = \sum_{i=1}^n \frac{d}{d\theta} \log(f(X_i, \hat{\theta}))$) and determine that one is a maximum

Approach to constructing MLE when indicators, $\mathbb{I}\{U\}$, are present: Logs of indicators and derivatives of indicators are very difficult to work with • Simplify likelihood function (splitting indicators when possible) • Make an argument for why the function is increasing or decreasing

• Determine the value at the bounds of the function

Estimating equations: $E_{\theta_1} g(\theta_2, X_1) = 0 \iff \theta_1 = \theta_2$

0.10 Fisher Information

Fisher information: $I(\theta) = E_\theta \left[\left(\frac{d}{d\theta} \log(\mathcal{L}(X_1, \theta)) \right)^2 \right]$

Properties: $E_\theta \left[\left(\frac{d}{d\theta} \log(\mathcal{L}(X_1, \theta)) \right) \right] = 0$ • $I(\theta) = Var \left(\frac{d}{d\theta} \log(\mathcal{L}(X_1, \theta)) \right)$ • $I(\theta) = -E_\theta \left[\frac{d^2}{d\theta^2} \log(\mathcal{L}(X_1, \theta)) \right]$ • $I_{X_1, X_2}(\theta) = I_{X_1}(\theta) + I_{X_2}(\theta)$ for X_1, X_2 independent (Information increases with larger sample!)

Cramer-Rau bound: $Var(\hat{\theta}) = Var(T(X_1, \dots, X_n)) \geq 1/I_{X_1, \dots, X_n}(\theta)$ where $I_{X_1, \dots, X_n}(\theta) = nE_\theta \left[\left(\frac{d}{d\theta} \log \mathcal{L}(X_1, \theta) \right)^2 \right]$ for $E(\hat{\theta}) = g(\theta)$

0.11 Examples

Newsvendor problem: [Setup] $D \sim Exp(\lambda)$, price p , cost c , buy-back price b , order quantity q , earnings W ; [Solution] $W = pq \min(q, D) - cq + b(q - \min(q, D)) \implies EW = (p - b)E(\min(q, D)) - (c - b)q = (p - b)(xP(D > q) + \int_0^D y f_d(y) dy) - (c - b)q \implies \frac{\partial}{\partial q} E(W)|_{q=q^*}$

Cramer-Rao bound: [Setup] $X \sim Exp(\lambda)$, MLE $= 1/\bar{X}_n$, does $1/\hat{\lambda}_{MLE}$ achieve lower bound?; [Solution] i) Unbiased: $E(1/\hat{\lambda}) = E(\bar{X}_n) = 1/\lambda$, ii) $Var(1/\hat{\lambda}) = Var(\bar{X}_n) = (1/n^2)nVar(X_1) = 1/(\lambda^2 n)$, iii) $I(1/\lambda) = I(\theta) = -E_X \left(\frac{d^2}{d\theta^2} \log \mathcal{L}(\theta) \right) : \mathcal{L}(\theta) = 1/\theta \exp(-x/\theta) \Rightarrow \log \mathcal{L}(\theta) = -\log \theta - x/\theta \Rightarrow \frac{d^2}{d\theta^2} \log \mathcal{L}(\theta) = 1/\theta^2 - 2X/\theta^3 \Rightarrow -E_X \left(\frac{d^2}{d\theta^2} \log \mathcal{L}(\theta) \right) = 1/\theta^2$