

A Unified Framework for Rerandomization using Quadratic Forms

Design and Analysis of Experiments (2024)

Kyle Schindl¹ Zach Branson²

Department of Statistics & Data Science, Carnegie Mellon University

¹schindl@stat.cmu.edu

²zach@stat.cmu.edu

Introduction

- We focus on treatment vs. control randomized experiments

- We focus on treatment vs. control randomized experiments
- We are interested in estimating the Average Treatment Effect (ATE)

$$\tau = \mathbb{E}[Y(1) - Y(0)]$$

where $Y(1)$ and $Y(0)$ are the outcomes under treatment and control.

- We focus on treatment vs. control randomized experiments
- We are interested in estimating the Average Treatment Effect (ATE)

$$\tau = \mathbb{E}[Y(1) - Y(0)]$$

where $Y(1)$ and $Y(0)$ are the outcomes under treatment and control.

- In completely randomized experiments, estimates of the ATE are unbiased.

- We focus on treatment vs. control randomized experiments
- We are interested in estimating the Average Treatment Effect (ATE)

$$\tau = \mathbb{E}[Y(1) - Y(0)]$$

where $Y(1)$ and $Y(0)$ are the outcomes under treatment and control.

- In completely randomized experiments, estimates of the ATE are unbiased.
- Covariate distributions are “balanced” *on average*.

- We focus on treatment vs. control randomized experiments
- We are interested in estimating the Average Treatment Effect (ATE)

$$\tau = \mathbb{E}[Y(1) - Y(0)]$$

where $Y(1)$ and $Y(0)$ are the outcomes under treatment and control.

- In completely randomized experiments, estimates of the ATE are unbiased.
- Covariate distributions are “balanced” *on average*.
- **We can do better than average.**

- We focus on treatment vs. control randomized experiments
- We are interested in estimating the Average Treatment Effect (ATE)

$$\tau = \mathbb{E}[Y(1) - Y(0)]$$

where $Y(1)$ and $Y(0)$ are the outcomes under treatment and control.

- In completely randomized experiments, estimates of the ATE are unbiased.
- Covariate distributions are “balanced” *on average*.
- **We can do better than average.**
- By ensuring covariate distributions are closer, we can obtain more precise estimates of the ATE while maintaining unbiasedness.

The Problem of Covariate Imbalance

The Problem of Covariate Imbalance

- Covariate imbalances between treatment and control increases standard errors and reduces statistical power when estimating causal effects (Senn 1989; Lachin 1988; Lin, Zhu, and Su 2015; Branson, Li, and Ding 2023).

The Problem of Covariate Imbalance

- Covariate imbalances between treatment and control increases standard errors and reduces statistical power when estimating causal effects (Senn 1989; Lachin 1988; Lin, Zhu, and Su 2015; Branson, Li, and Ding 2023).
- This is why classical experimental designs suggest blocking similar subjects together when randomizing.

The Problem of Covariate Imbalance

- Covariate imbalances between treatment and control increases standard errors and reduces statistical power when estimating causal effects (Senn 1989; Lachin 1988; Lin, Zhu, and Su 2015; Branson, Li, and Ding 2023).
- This is why classical experimental designs suggest blocking similar subjects together when randomizing.
- Another method of reducing covariate imbalance is to rerandomize.




Figure 1: Procedure for implementing rerandomization

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as
 - 2^K factorial designs (Branson, Dasgupta, and Rubin 2016; Li, Ding, and Rubin 2020)

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as
 - 2^K factorial designs (Branson, Dasgupta, and Rubin 2016; Li, Ding, and Rubin 2020)
 - Sequential designs (Zhou et al. 2018)

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as
 - 2^K factorial designs (Branson, Dasgupta, and Rubin 2016; Li, Ding, and Rubin 2020)
 - Sequential designs (Zhou et al. 2018)
 - Clustered experiments (Lu et al. 2023)

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as
 - 2^K factorial designs (Branson, Dasgupta, and Rubin 2016; Li, Ding, and Rubin 2020)
 - Sequential designs (Zhou et al. 2018)
 - Clustered experiments (Lu et al. 2023)
 - Bayesian designs (Liu et al. 2023)

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as
 - 2^K factorial designs (Branson, Dasgupta, and Rubin 2016; Li, Ding, and Rubin 2020)
 - Sequential designs (Zhou et al. 2018)
 - Clustered experiments (Lu et al. 2023)
 - Bayesian designs (Liu et al. 2023)
- Most of these extensions utilize the Mahalanobis distance.

- (Morgan and Rubin 2012): Randomize until $M < a$, for some pre-specified $a > 0$, where

$$M := (\bar{X}_T - \bar{X}_C)^T \Sigma^{-1} (\bar{X}_T - \bar{X}_C).$$

where

- $\bar{X}_T, \bar{X}_C \in \mathbb{R}^d$ are the covariate means in treatment and control.
- $\Sigma := \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$
- There have been many extensions to other design settings such as
 - 2^K factorial designs (Branson, Dasgupta, and Rubin 2016; Li, Ding, and Rubin 2020)
 - Sequential designs (Zhou et al. 2018)
 - Clustered experiments (Lu et al. 2023)
 - Bayesian designs (Liu et al. 2023)
- Most of these extensions utilize the Mahalanobis distance.
- We'll consider: What distances are optimal for rerandomization?

Properties of Mahalanobis Rerandomization

- As long as $\sum_{i=1}^n W_i = \sum_{i=1}^n (1 - W_i)$, where $W_i \in \{0, 1\}$ is the treatment assignment, then $\hat{\tau}$ remains unbiased.

Properties of Mahalanobis Rerandomization

- As long as $\sum_{i=1}^n W_i = \sum_{i=1}^n (1 - W_i)$, where $W_i \in \{0, 1\}$ is the treatment assignment, then $\hat{\tau}$ remains unbiased.
- If $\bar{X}_T - \bar{X}_C \mid x \sim \mathcal{N}(0, \Sigma)$ then, Mahalanobis Rerandomization applies an equal-percentage variance reduction to all covariates,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, M \leq a) = v_a \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$$

where $v_a = \mathbb{P}(\chi_{d+2}^2 \leq a) / \mathbb{P}(\chi_d^2 \leq a) \leq 1$.

Properties of Mahalanobis Rerandomization

- As long as $\sum_{i=1}^n W_i = \sum_{i=1}^n (1 - W_i)$, where $W_i \in \{0, 1\}$ is the treatment assignment, then $\hat{\tau}$ remains unbiased.
- If $\bar{X}_T - \bar{X}_C \mid x \sim \mathcal{N}(0, \Sigma)$ then, Mahalanobis Rerandomization applies an equal-percentage variance reduction to all covariates,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, M \leq a) = v_a \text{Cov}(\bar{X}_T - \bar{X}_C \mid x)$$

where $v_a = \mathbb{P}(\chi_{d+2}^2 \leq a) / \mathbb{P}(\chi_d^2 \leq a) \leq 1$.

- When the treatment effect is additive, this leads to a variance reduction of

$$\frac{\mathbb{V}(\hat{\tau} \mid x, M < a)}{\mathbb{V}(\hat{\tau} \mid x)} = 1 - (1 - v_a)R^2 \leq 1$$

where

$$\hat{\tau} = \frac{1}{n_1} \sum_{i=1}^n Y_i(1) W_i - \frac{1}{n_0} \sum_{i=1}^n Y_i(0)(1 - W_i)$$

Properties of Mahalanobis Rerandomization

- As long as $\sum_{i=1}^n W_i = \sum_{i=1}^n (1 - W_i)$, where $W_i \in \{0, 1\}$ is the treatment assignment, then $\hat{\tau}$ remains unbiased.
- If $\bar{X}_T - \bar{X}_C | x \sim \mathcal{N}(0, \Sigma)$ then, Mahalanobis Rerandomization applies an equal-percentage variance reduction to all covariates,

$$\text{Cov}(\bar{X}_T - \bar{X}_C | x, M \leq a) = v_a \text{Cov}(\bar{X}_T - \bar{X}_C | x)$$

where $v_a = \mathbb{P}(\chi_{d+2}^2 \leq a) / \mathbb{P}(\chi_d^2 \leq a) \leq 1$.

- When the treatment effect is additive, this leads to a variance reduction of

$$\frac{\mathbb{V}(\hat{\tau} | x, M < a)}{\mathbb{V}(\hat{\tau} | x)} = 1 - (1 - v_a)R^2 \leq 1$$


where

$$\hat{\tau} = \frac{1}{n_1} \sum_{i=1}^n Y_i(1) W_i - \frac{1}{n_0} \sum_{i=1}^n Y_i(0)(1 - W_i)$$

- Asymptotically, this is true with unequal sample sizes and non-additivity (Li, Ding, and Rubin 2018).


Mahalanobis Rerandomization Visualized

- Geometrically, we can see that Mahalanobis Rerandomization preserves the “shape” of the covariance matrix but reduces the scale.



Mahalanobis Rerandomization Visualized


- Geometrically, we can see that Mahalanobis Rerandomization preserves the “shape” of the covariance matrix but reduces the scale.



- Each eigenvector of the covariance matrix Σ is scaled by v_a .

Mahalanobis Rerandomization Visualized


- Geometrically, we can see that Mahalanobis Rerandomization preserves the “shape” of the covariance matrix but reduces the scale.



- Each eigenvector of the covariance matrix Σ is scaled by v_a .
- Placing equal priority on all covariates create challenges since $v_a \rightarrow 1$ as $d \rightarrow \infty$ (no variance reduction).

Mahalanobis Rerandomization Visualized

- Geometrically, we can see that Mahalanobis Rerandomization preserves the “shape” of the covariance matrix but reduces the scale.



- Each eigenvector of the covariance matrix Σ is scaled by v_a .
- Placing equal priority on all covariates create challenges since $v_a \rightarrow 1$ as $d \rightarrow \infty$ (no variance reduction).
- Mahalanobis Rerandomization may be “wasting” variance reduction on unimportant covariates

- Other recent proposed balance metrics include:

- Other recent proposed balance metrics include:
- A penalized Mahalanobis distance (Branson and Shao 2021):

$$M_\lambda = (\bar{X}_T - \bar{X}_C)^T (\Sigma + \lambda I_d)^{-1} (\bar{X}_T - \bar{X}_C)$$

- Other recent proposed balance metrics include:
- A penalized Mahalanobis distance (Branson and Shao 2021):

$$M_\lambda = (\bar{X}_T - \bar{X}_C)^T (\Sigma + \lambda I_d)^{-1} (\bar{X}_T - \bar{X}_C)$$

- The Mahalanobis distance after dropping the bottom $d - k$ principal components (Zhang, Yin, and Rubin 2023):

$$M_k = (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})^T \Sigma_Z^{-1} (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})$$

where Z are the principal components.

- Other recent proposed balance metrics include:
- A penalized Mahalanobis distance (Branson and Shao 2021):

$$M_\lambda = (\bar{X}_T - \bar{X}_C)^T (\Sigma + \lambda I_d)^{-1} (\bar{X}_T - \bar{X}_C)$$

- The Mahalanobis distance after dropping the bottom $d - k$ principal components (Zhang, Yin, and Rubin 2023):

$$M_k = (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})^T \Sigma_Z^{-1} (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})$$

where Z are the principal components.

- Each method can be written as $(\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$ for some $A \in \mathbb{R}^{d \times d}$, e.g.,

$$M = (\bar{X}_T - \bar{X}_C)^T \underbrace{\Sigma^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

- Other recent proposed balance metrics include:
- A penalized Mahalanobis distance (Branson and Shao 2021):

$$M_\lambda = (\bar{X}_T - \bar{X}_C)^T \underbrace{(\Sigma + \lambda I_d)^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

- The Mahalanobis distance after dropping the bottom $d - k$ principal components (Zhang, Yin, and Rubin 2023):

$$M_k = (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})^T \Sigma_Z^{-1} (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})$$

where Z are the principal components.

- Each method can be written as $(\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$ for some $A \in \mathbb{R}^{d \times d}$, e.g.,

$$M = (\bar{X}_T - \bar{X}_C)^T \underbrace{\Sigma^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

Balancing Metrics as Quadratic Forms

- Other recent proposed balance metrics include:
- A penalized Mahalanobis distance (Branson and Shao 2021):

$$M_\lambda = (\bar{X}_T - \bar{X}_C)^T \underbrace{(\Sigma + \lambda I_d)^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

- The Mahalanobis distance after dropping the bottom $d - k$ principal components (Zhang, Yin, and Rubin 2023):

$$\begin{aligned} M_k &= (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})^T \Sigma_Z^{-1} (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)}) \\ &= (\bar{X}_T - \bar{X}_C)^T \underbrace{V \begin{pmatrix} D_k & 0 \\ 0 & 0 \end{pmatrix} V^T}_A (\bar{X}_T - \bar{X}_C) \end{aligned}$$

where D_k are the singular values from $x = UDV^T$.

- Each method can be written as $(\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$ for some $A \in \mathbb{R}^{d \times d}$, e.g.,

$$M = (\bar{X}_T - \bar{X}_C)^T \underbrace{\Sigma^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

Balancing Metrics as Quadratic Forms

- Other recent proposed balance metrics include:
- A penalized Mahalanobis distance (Branson and Shao 2021):

$$M_\lambda = (\bar{X}_T - \bar{X}_C)^T \underbrace{(\Sigma + \lambda I_d)^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

- The Mahalanobis distance after dropping the bottom $d - k$ principal components (Zhang, Yin, and Rubin 2023):

$$\begin{aligned} M_k &= (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)})^T \Sigma_Z^{-1} (\bar{Z}_T^{(k)} - \bar{Z}_C^{(k)}) \\ &= (\bar{X}_T - \bar{X}_C)^T \underbrace{V \begin{pmatrix} D_k & 0 \\ 0 & 0 \end{pmatrix} V^T}_A (\bar{X}_T - \bar{X}_C) \end{aligned}$$

where D_k are the singular values from $x = UDV^T$.

- Each method can be written as $(\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$ for some $A \in \mathbb{R}^{d \times d}$, e.g.,

$$M = (\bar{X}_T - \bar{X}_C)^T \underbrace{\Sigma^{-1}}_A (\bar{X}_T - \bar{X}_C)$$

- This leads to the question: what is a good choice of A ?

Rerandomization using Quadratic Forms

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- Some key results we cover:

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- Some key results we cover:
 - The covariance of $\bar{X}_T - \bar{X}_C$ conditioning on any quadratic form.

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- Some key results we cover:
 - The covariance of $\bar{X}_T - \bar{X}_C$ conditioning on any quadratic form.
 - The optimal choice of A for covariate balance

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- Some key results we cover:
 - The covariance of $\bar{X}_T - \bar{X}_C$ conditioning on any quadratic form.
 - The optimal choice of A for covariate balance
 - The optimal choice of A for reducing the variance of $\hat{\tau}$

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- Some key results we cover:
 - The covariance of $\bar{X}_T - \bar{X}_C$ conditioning on any quadratic form.
 - The optimal choice of A for covariate balance
 - The optimal choice of A for reducing the variance of $\hat{\tau}$
- Throughout, we assume A is positive-definite and that $\bar{X}_T - \bar{X}_C \mid x \sim \mathcal{N}(0, \Sigma)$.

- In this work, we establish a general framework for randomizing until $Q_A(x) \leq a$ where

$$Q_A(x) := (\bar{X}_T - \bar{X}_C)^T A (\bar{X}_T - \bar{X}_C)$$

- We select a by setting an acceptance probability α such that

$$\mathbb{P}(Q_A(x) \leq a) = \alpha$$

- Some key results we cover:
 - The covariance of $\bar{X}_T - \bar{X}_C$ conditioning on any quadratic form.
 - The optimal choice of A for covariate balance
 - The optimal choice of A for reducing the variance of $\hat{\tau}$
- Throughout, we assume A is positive-definite and that $\bar{X}_T - \bar{X}_C \mid x \sim \mathcal{N}(0, \Sigma)$.
- Also note that we assume the covariates have been standardized.

Covariance Reduction under Quadratic Form Rerandomization

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2}A\Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{(q_{j,\eta})_{1 \leq j \leq d}\} \right) \Gamma^T \Sigma^{1/2}$$

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2}A\Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{(q_{j,\eta})_{1 \leq j \leq d}\} \right) \Gamma^T \Sigma^{1/2}$$

- $\Gamma \in \mathbb{R}^{d \times d}$ is the orthogonal matrix of eigenvectors of Σ

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2}A\Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{(q_{j,\eta})_{1 \leq j \leq d}\} \right) \Gamma^T \Sigma^{1/2}$$

- $\Gamma \in \mathbb{R}^{d \times d}$ is the orthogonal matrix of eigenvectors of Σ
- $q_{j,\eta}$ is defined as

$$q_{j,\eta} = \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \eta_j \mathcal{Z}_j^2 \leq a \right]$$

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2}A\Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{(q_{j,\eta})_{1 \leq j \leq d}\} \right) \Gamma^T \Sigma^{1/2}$$

- $\Gamma \in \mathbb{R}^{d \times d}$ is the orthogonal matrix of eigenvectors of Σ
- $q_{j,\eta}$ is defined as

$$q_{j,\eta} = \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \eta_j \mathcal{Z}_j^2 \leq a \right]$$

- $\eta_1 \geq \dots \geq \eta_d$ are the eigenvalues of $\Sigma^{1/2}A\Sigma^{1/2}$

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2} A \Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{ (q_{j,\eta})_{1 \leq j \leq d} \} \right) \Gamma^T \Sigma^{1/2}$$

- $\Gamma \in \mathbb{R}^{d \times d}$ is the orthogonal matrix of eigenvectors of Σ
- $q_{j,\eta}$ is defined as

$$q_{j,\eta} = \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \eta_j \mathcal{Z}_j^2 \leq a \right]$$

- $\eta_1 \geq \dots \geq \eta_d$ are the eigenvalues of $\Sigma^{1/2} A \Sigma^{1/2}$
- $\mathcal{Z}_1, \dots, \mathcal{Z}_d \sim \mathcal{N}(0, 1)$

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2}A\Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{(q_{j,\eta})_{1 \leq j \leq d}\} \right) \Gamma^T \Sigma^{1/2}$$

- $\Gamma \in \mathbb{R}^{d \times d}$ is the orthogonal matrix of eigenvectors of Σ
- $q_{j,\eta}$ is defined as

$$q_{j,\eta} = \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \eta_j \mathcal{Z}_j^2 \leq a \right]$$

- $\eta_1 \geq \dots \geq \eta_d$ are the eigenvalues of $\Sigma^{1/2}A\Sigma^{1/2}$
- $\mathcal{Z}_1, \dots, \mathcal{Z}_d \sim \mathcal{N}(0, 1)$
- $q_{1,\eta} \leq \dots \leq q_{d,\eta} \leq 1$.

Theorem 1 (Covariance Reduction)

Suppose that $\Sigma^{1/2} A \Sigma^{1/2}$ and Σ share an eigenbasis. Then,

$$\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a) = \Sigma^{1/2} \Gamma \left(\text{diag}\{(q_{j,\eta})_{1 \leq j \leq d}\} \right) \Gamma^T \Sigma^{1/2}$$

- $\Gamma \in \mathbb{R}^{d \times d}$ is the orthogonal matrix of eigenvectors of Σ
- $q_{j,\eta}$ is defined as


$$\begin{aligned} q_{j,\eta} &= \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \eta_j \mathcal{Z}_j^2 \leq a \right] \\ &= \frac{p_d}{\eta_j} \det(\Sigma^{1/2} A \Sigma^{1/2})^{1/d} \alpha^{2/d} + o(\alpha^{2/d}) \end{aligned}$$

for a sufficiently small α , where $p_d = \frac{2\pi}{d+2} \left(\frac{2\pi^{d/2}}{d\Gamma(d/2)} \right)^{-2/d}$ (Lu et al. 2023).


- $\eta_1 \geq \dots \geq \eta_d$ are the eigenvalues of $\Sigma^{1/2} A \Sigma^{1/2}$
- $\mathcal{Z}_1, \dots, \mathcal{Z}_d \sim \mathcal{N}(0, 1)$
- $q_{1,\eta} \leq \dots \leq q_{d,\eta} \leq 1$.

Geometric Intuition behind Quadratic Form Rerandomization

- The eigenvalues of $\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)$ are given by $q_{1,\eta}\lambda_1, \dots, q_{d,\eta}\lambda_d$.




(a) Covariance Matrix before rerandomization



(b) Covariance Matrix after rerandomization


Variance Reduction Factors Visualized

- Depending on the choice of A , there is a different amount of variance reduction applied to each eigenvalue.



Variance Reduction Factors Visualized

- Depending on the choice of A , there is a different amount of variance reduction applied to each eigenvalue.



- We show that different choices of A can lead to significantly different precision levels when estimating $\hat{\tau}$.

Variance Reduction Visualized

- Depending on the relationship between the potential outcomes and x , and the eigenstructure of Σ , different choices of A yield different levels of precision.




Figure 3: Distribution of $\hat{\tau}$

Variance Reduction Visualized

- Depending on the relationship between the potential outcomes and x , and the eigenstructure of Σ , different choices of A yield different levels of precision.




Figure 3: Distribution of $\hat{\tau}$

- Given the clear differences between each of these methods, the question arises: which choice of A is optimal?

- First we consider optimality in terms of covariate balance.

- First we consider optimality in terms of covariate balance.
- We consider two targets for quantifying covariate balance:

Choosing the Optimal Matrix A

- First we consider optimality in terms of covariate balance.
- We consider two targets for quantifying covariate balance:
 - The choice of A that maximizes the variance reduction applied across all covariates,

$$\max_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d (1 - q_{j,\eta}) = \min_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d q_{j,\eta}$$

Choosing the Optimal Matrix A

- First we consider optimality in terms of covariate balance.
- We consider two targets for quantifying covariate balance:
 - The choice of A that maximizes the variance reduction applied across all covariates,

$$\max_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d (1 - q_{j,\eta}) = \min_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d q_{j,\eta}$$

- The choice of A that constrains the covariance matrix after rerandomization the most, which we define as

$$\min_{A \in \mathbb{R}^{d \times d}} \|\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)\|_F^2$$

where $\|\cdot\|_F^2 = \sum_{j=1}^d \sigma_j(\cdot)$ and $\sigma_j(\cdot)$ are the singular values of the matrix.

Choosing the Optimal Matrix A

- First we consider optimality in terms of covariate balance.
- We consider two targets for quantifying covariate balance:
 - The choice of A that maximizes the variance reduction applied across all covariates,

$$\max_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d (1 - q_{j,\eta}) = \min_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d q_{j,\eta}$$

- The choice of A that constrains the covariance matrix after rerandomization the most, which we define as

$$\min_{A \in \mathbb{R}^{d \times d}} \|\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)\|_F^2$$

where $\|\cdot\|_F^2 = \sum_{j=1}^d \sigma_j(\cdot)$ and $\sigma_j(\cdot)$ are the singular values of the matrix.

- Equivalently, $\sigma_j^2(\cdot)$ are the eigenvalues of the sample covariance matrix.

Choosing the Optimal Matrix A

- First we consider optimality in terms of covariate balance.
- We consider two targets for quantifying covariate balance:
 - The choice of A that maximizes the variance reduction applied across all covariates,

$$\max_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d (1 - q_{j,\eta}) = \min_{A \in \mathbb{R}^{d \times d}} \sum_{j=1}^d q_{j,\eta}$$

- The choice of A that constrains the covariance matrix after rerandomization the most, which we define as

$$\min_{A \in \mathbb{R}^{d \times d}} \|\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)\|_F^2$$

where $\|\cdot\|_F^2 = \sum_{j=1}^d \sigma_j(\cdot)$ and $\sigma_j(\cdot)$ are the singular values of the matrix.

- Equivalently, $\sigma_j^2(\cdot)$ are the eigenvalues of the sample covariance matrix.
- Later, we investigate how this impacts the precision of $\hat{\tau}$.

Theorem 2 (Covariate Balance)

For all positive-definite matrices $A \in \mathbb{R}^{d \times d}$

- $A = I_d$ minimizes the Frobenius norm:

$$||\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_I(x) \leq a)||_F \leq ||\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)||_F + o(\alpha^{2/d})$$

- $A = \Sigma^{-1}$ maximizes the total variance reduction applied

$$\sum_{j=1}^d v_a \leq \sum_{j=1}^d q_{j,\eta} + o(\alpha^{2/d})$$

Theorem 2 (Covariate Balance)

For all positive-definite matrices $A \in \mathbb{R}^{d \times d}$

- $A = I_d$ minimizes the Frobenius norm:

$$||\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_I(x) \leq a)||_F \leq ||\text{Cov}(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)||_F + o(\alpha^{2/d})$$

- $A = \Sigma^{-1}$ maximizes the total variance reduction applied

$$\sum_{j=1}^d v_a \leq \sum_{j=1}^d q_{j,\eta} + o(\alpha^{2/d})$$

- These two methods represent opposite sides of the rerandomization spectrum:

Theorem 2 (Covariate Balance)

For all positive-definite matrices $A \in \mathbb{R}^{d \times d}$

- $A = I_d$ minimizes the Frobenius norm:

$$\|Cov(\bar{X}_T - \bar{X}_C \mid x, Q_I(x) \leq a)\|_F \leq \|Cov(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)\|_F + o(\alpha^{2/d})$$

- $A = \Sigma^{-1}$ maximizes the total variance reduction applied

$$\sum_{j=1}^d v_a \leq \sum_{j=1}^d q_{j,\eta} + o(\alpha^{2/d})$$

- These two methods represent opposite sides of the rerandomization spectrum:
 - Mahalanobis Rerandomization ($A = \Sigma^{-1}$) leaves the shape of the covariance matrix unchanged by applying an equal variance reduction to all eigenvectors.

Theorem 2 (Covariate Balance)

For all positive-definite matrices $A \in \mathbb{R}^{d \times d}$

- $A = I_d$ minimizes the Frobenius norm:

$$\|Cov(\bar{X}_T - \bar{X}_C \mid x, Q_I(x) \leq a)\|_F \leq \|Cov(\bar{X}_T - \bar{X}_C \mid x, Q_A(x) \leq a)\|_F + o(\alpha^{2/d})$$

- $A = \Sigma^{-1}$ maximizes the total variance reduction applied

$$\sum_{j=1}^d v_a \leq \sum_{j=1}^d q_{j,\eta} + o(\alpha^{2/d})$$

- These two methods represent opposite sides of the rerandomization spectrum:
 - Mahalanobis Rerandomization ($A = \Sigma^{-1}$) leaves the shape of the covariance matrix unchanged by applying an equal variance reduction to all eigenvectors.
 - Euclidean Rerandomization ($A = I_d$) scales each eigenvector by a different factor such that their magnitude is the same.

Quantifying the variance reduction to $\hat{\tau}$

- When the treatment effect is additive, it follows that

$$Y_i(W_i) = \beta_0 + Z_i\beta_Z + \tau W_i + \varepsilon_i$$

where $\beta_0 + Z_i\beta_Z$ represent the linear projection of Y_i onto the principal components $Z = XV$ and ε_i is a residual that encompasses deviations from the projection.

- When the treatment effect is additive, it follows that

$$Y_i(W_i) = \beta_0 + Z_i\beta_Z + \tau W_i + \varepsilon_i$$

where $\beta_0 + Z_i\beta_Z$ represent the linear projection of Y_i onto the principal components $Z = XV$ and ε_i is a residual that encompasses deviations from the projection.

Theorem 3 (Variance Reduction)

Suppose the treatment effect is additive. Then,

$$\mathbb{V}(\hat{\tau} \mid x) - \mathbb{V}(\hat{\tau} \mid x, Q_A(x) \leq a) = \sum_{j=1}^d \beta_{Z,j}^2 \lambda_j (1 - q_{j,\eta}) \geq 0$$

where $\lambda_1, \dots, \lambda_d$ are the eigenvalues of Σ and $\hat{\tau}$ is the mean-difference estimator.

The Optimal Choice of A for variance reduction to $\hat{\tau}$

Theorem 4 (Optimal A)

Suppose that the treatment effect is additive. Then for all positive-definite matrices $A \in \mathbb{R}^{d \times d}$,

$$A^* = \Gamma \begin{pmatrix} \beta_{Z,1}^2 & & 0 \\ & \ddots & \\ 0 & & \beta_{Z,d}^2 \end{pmatrix} \Gamma^T$$

minimizes the variance of the mean-differences estimator, i.e.,

$$\mathbb{V}(\hat{\tau} \mid x, Q_{A^*}(x) \leq a) \leq \mathbb{V}(\hat{\tau} \mid x, Q_A(x) \leq a) + o(\alpha^{2/d})$$

where Γ is the orthogonal matrix of eigenvectors of Σ .

Which A is optimal? It depends.

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.
- For example:
 - When $\beta_{Z,j}^2 = \frac{1}{\lambda_j}$, $A = \Sigma^{-1}$ (Mahalanobis Rerandomization) is optimal.

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.
- For example:
 - When $\beta_{Z,j}^2 = \frac{1}{\lambda_j}$, $A = \Sigma^{-1}$ (Mahalanobis Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = 1$, $A = I_d$ (Euclidean Rerandomization) is optimal.

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.
- For example:
 - When $\beta_{Z,j}^2 = \frac{1}{\lambda_j}$, $A = \Sigma^{-1}$ (Mahalanobis Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = 1$, $A = I_d$ (Euclidean Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = \lambda_j$, the choice $A = \Sigma$ is optimal.

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.
- For example:
 - When $\beta_{Z,j}^2 = \frac{1}{\lambda_j}$, $A = \Sigma^{-1}$ (Mahalanobis Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = 1$, $A = I_d$ (Euclidean Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = \lambda_j$, the choice $A = \Sigma$ is optimal.
- When the top eigenvectors and potential outcomes are closely related, choices like $A = I_d, \Sigma$ or PCA rerandomization tend to be better

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.
- For example:
 - When $\beta_{Z,j}^2 = \frac{1}{\lambda_j}$, $A = \Sigma^{-1}$ (Mahalanobis Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = 1$, $A = I_d$ (Euclidean Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = \lambda_j$, the choice $A = \Sigma$ is optimal.
- When the top eigenvectors and potential outcomes are closely related, choices like $A = I_d, \Sigma$ or PCA rerandomization tend to be better
- When the bottom eigenvectors and potential outcomes are closely related, methods like $A = \Sigma^{-1}$ are better.

Which A is optimal? It depends.

- For any choice of A there exists some β_Z such that A is optimal.
- For example:
 - When $\beta_{Z,j}^2 = \frac{1}{\lambda_j}$, $A = \Sigma^{-1}$ (Mahalanobis Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = 1$, $A = I_d$ (Euclidean Rerandomization) is optimal.
 - When $\beta_{Z,j}^2 = \lambda_j$, the choice $A = \Sigma$ is optimal.
- When the top eigenvectors and potential outcomes are closely related, choices like $A = I_d, \Sigma$ or PCA rerandomization tend to be better
- When the bottom eigenvectors and potential outcomes are closely related, methods like $A = \Sigma^{-1}$ are better.
- However, different choices may be less risky than others.

Theorem 5 (Minimax A)

The choice $A = I_d$ minimizes the maximum difference between the optimal quadratic form and any other choice of A , given by

$$\min_{A \in \mathbb{R}^{d \times d} \mid \|\beta_Z\|_2 \leq c} \max |\mathbb{V}(\hat{\tau} \mid x, Q_{A^*}(x) \leq a) - \mathbb{V}(\hat{\tau} \mid x, Q_A(x) \leq a)|.$$

Theorem 5 (Minimax A)

The choice $A = I_d$ minimizes the maximum difference between the optimal quadratic form and any other choice of A , given by

$$\min_{A \in \mathbb{R}^{d \times d} \mid \|\beta_Z\|_2 \leq c} \max |\mathbb{V}(\hat{\tau} \mid x, Q_{A^*}(x) \leq a) - \mathbb{V}(\hat{\tau} \mid x, Q_A(x) \leq a)|.$$

- Under $A = I_d$, the precision of $\hat{\tau}$ is never too far from the optimal choice.

The minimax optimal choice of A

Theorem 5 (Minimax A)

The choice $A = I_d$ minimizes the maximum difference between the optimal quadratic form and any other choice of A , given by

$$\min_{A \in \mathbb{R}^{d \times d} \mid \|\beta_Z\|_2 \leq c} \max |\mathbb{V}(\hat{\tau} \mid x, Q_{A^*}(x) \leq a) - \mathbb{V}(\hat{\tau} \mid x, Q_A(x) \leq a)|.$$

- Under $A = I_d$, the precision of $\hat{\tau}$ is never too far from the optimal choice.
- For intuition, compare the variance reduction factors between the two:

$$q_{j,\lambda} = \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \lambda_j \mathcal{Z}_j^2 \leq a \right]$$
$$q_{j,\beta\lambda}^* = \mathbb{E} \left[\mathcal{Z}_j^2 \mid x, \sum_{j=1}^d \beta_{Z,j}^2 \lambda_j \mathcal{Z}_j^2 \leq a \right]$$

- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.

- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.
- That way $\sum_{j=1}^d \lambda_j = d = \text{tr}(\Sigma)$

- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.
- That way $\sum_{j=1}^d \lambda_j = d = \text{tr}(\Sigma)$
- We simulate $n = 500$ observations where 250 units are sent to treatment and 250 units are sent to control.

- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.
- That way $\sum_{j=1}^d \lambda_j = d = \text{tr}(\Sigma)$
- We simulate $n = 500$ observations where 250 units are sent to treatment and 250 units are sent to control.
- Potential outcomes $Y_i(0)$ and $Y_i(1)$ are defined as

$$Y_i(0) = \mathcal{N}(Z\beta_Z, 1)$$

$$Y_i(1) = Y_i(0) + \tau$$

where $\tau = 1$, $\beta_Z \in \mathbb{R}^d$, and $Z = XV$ are the principal components of X .

- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.
- That way $\sum_{j=1}^d \lambda_j = d = \text{tr}(\Sigma)$
- We simulate $n = 500$ observations where 250 units are sent to treatment and 250 units are sent to control.
- Potential outcomes $Y_i(0)$ and $Y_i(1)$ are defined as

$$Y_i(0) = \mathcal{N}(Z\beta_Z, 1)$$

$$Y_i(1) = Y_i(0) + \tau$$

where $\tau = 1$, $\beta_Z \in \mathbb{R}^d$, and $Z = XV$ are the principal components of X .

- In this presentation we consider two choices of β_Z :

- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.
- That way $\sum_{j=1}^d \lambda_j = d = \text{tr}(\Sigma)$
- We simulate $n = 500$ observations where 250 units are sent to treatment and 250 units are sent to control.
- Potential outcomes $Y_i(0)$ and $Y_i(1)$ are defined as

$$Y_i(0) = \mathcal{N}(Z\beta_Z, 1)$$

$$Y_i(1) = Y_i(0) + \tau$$

where $\tau = 1$, $\beta_Z \in \mathbb{R}^d$, and $Z = XV$ are the principal components of X .

- In this presentation we consider two choices of β_Z :
 - β_Z such that Mahalanobis is optimal ($A = \Sigma^{-1}$)


- Because the optimal choice of A depends on β_Z as well as the distribution of eigenvalues, we simulate $\lambda_1, \dots, \lambda_d \sim d \cdot \text{Dirichlet}(\gamma)$.
- That way $\sum_{j=1}^d \lambda_j = d = \text{tr}(\Sigma)$
- We simulate $n = 500$ observations where 250 units are sent to treatment and 250 units are sent to control.
- Potential outcomes $Y_i(0)$ and $Y_i(1)$ are defined as


$$Y_i(0) = \mathcal{N}(Z\beta_Z, 1)$$

$$Y_i(1) = Y_i(0) + \tau$$


where $\tau = 1$, $\beta_Z \in \mathbb{R}^d$, and $Z = XV$ are the principal components of X .

- In this presentation we consider two choices of β_Z :
 - β_Z such that Mahalanobis is optimal ($A = \Sigma^{-1}$)
 - β_Z such that Euclidean is optimal ($A = I_d$)





- In these settings, Euclidean is either the best or second best choice of A



- In these settings, Euclidean is either the best or second best choice of A
- We tend to find asymmetry: Mahalanobis is either the best in settings favorable to it, or it is the worst.

- We derived general results for rerandomization using any quadratic form.

- We derived general results for rerandomization using any quadratic form.
- This allows researchers to quickly and easily derive new rerandomization schemes not previously considered.

- We derived general results for rerandomization using any quadratic form.
- This allows researchers to quickly and easily derive new rerandomization schemes not previously considered.
- We find that $A = \Sigma^{-1}$ and $A = I_d$ optimize different measurements of covariate balance.




- We derived general results for rerandomization using any quadratic form.
- This allows researchers to quickly and easily derive new rerandomization schemes not previously considered.
- We find that $A = \Sigma^{-1}$ and $A = I_d$ optimize different measurements of covariate balance.
- However, $A = I_d$ is never too far away from the optimal choice, and tends to be more precise on average.








- We derived general results for rerandomization using any quadratic form.
- This allows researchers to quickly and easily derive new rerandomization schemes not previously considered.
- We find that $A = \Sigma^{-1}$ and $A = I_d$ optimize different measurements of covariate balance.
- However, $A = I_d$ is never too far away from the optimal choice, and tends to be more precise on average.
- In future work, we plan to extend this framework to sequential experiments as well as multi-valued treatments.




Thank you!

- Questions?

References

-  Branson, Zach, Tirthankar Dasgupta, and Donald B Rubin (2016). **“Improving covariate balance in 2 K factorial designs via rerandomization with an application to a New York City Department of Education High School Study”**. In.
-  Branson, Zach, Xinran Li, and Peng Ding (May 2023). **“Power and sample size calculations for rerandomization”**. In: *Biometrika* 111.1, pp. 355–363. ISSN: 1464-3510. eprint: <https://academic.oup.com/biomet/article-pdf/111/1/355/56665596/asad027.pdf>.
-  Branson, Zach and Stephane Shao (2021). **“Ridge rerandomization: An experimental design strategy in the presence of covariate collinearity”**. In: *Journal of Statistical Planning and Inference* 211, pp. 287–314.

-  Lachin, John M (1988). “**Statistical properties of randomization in clinical trials**”. In: *Controlled clinical trials* 9.4, pp. 289–311.
-  Li, Xinran, Peng Ding, and Donald B Rubin (2018). “**Asymptotic theory of rerandomization in treatment–control experiments**”. In: *Proceedings of the National Academy of Sciences* 115.37, pp. 9157–9162.
-  — (2020). “**Rerandomization in 2^K factorial experiments**”. In.
-  Lin, Yunzhi, Ming Zhu, and Zheng Su (2015). “**The pursuit of balance: an overview of covariate-adaptive randomization techniques in clinical trials**”. In: *Contemporary clinical trials* 45, pp. 21–25.
-  Liu, Zhaoyang et al. (2023). “**Bayesian Criterion for Re-randomization**”. In: *arXiv preprint arXiv:2303.07904*.
-  Lu, Xin et al. (2023). “**Design-based theory for cluster rerandomization**”. In: *Biometrika* 110.2, pp. 467–483.
-  Morgan, Kari Lock and Donald B Rubin (2012). “**Rerandomization to improve covariate balance in experiments**”. In.

-  Senn, SJ (1989). “**Covariate imbalance and random allocation in clinical trials**”. In: *Statistics in medicine* 8.4, pp. 467–475.
-  Zhang, Hengtao, Guosheng Yin, and Donald B Rubin (2023). “**PCA rerandomization**”. In: *Canadian Journal of Statistics*.
-  Zhou, Quan et al. (2018). “**Sequential rerandomization**”. In: *Biometrika* 105.3, pp. 745–752.