Adaptive Experimental Design Using Shrinkage Estimators

Evan T. R. Rosenman[†], Kristen B. Hunter[⋆]

[†]Claremont McKenna College, [⋆]University of New South Wales

October 9, 2026

Outline

1 Background and Motivation

2 Estimators and Risk

3 Oracle Designs

4 Adaptation & Inference

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control
  - ... units arrive one at a time. We assign them to a treatment arm and observe their outcomes immediately

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control
  - ... units arrive one at a time. We assign them to a treatment arm and observe their outcomes immediately
  - ... seek to *adapt* treatment assignments for later units, based on outcomes observed in previous units, to improve estimation

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control
  - ... units arrive one at a time. We assign them to a treatment arm and observe their outcomes immediately
  - ... seek to *adapt* treatment assignments for later units, based on outcomes observed in previous units, to improve estimation
- **Simple example:** testing many candidate redesigns for a website ("A/B/C/D..." test)

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control
  - ... units arrive one at a time. We assign them to a treatment arm and observe their outcomes immediately
  - ... seek to *adapt* treatment assignments for later units, based on outcomes observed in previous units, to improve estimation
- **Simple example:** testing many candidate redesigns for a website ("A/B/C/D..." test)
  - assign website visitors to a different UI as they arrive

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control
  - ... units arrive one at a time. We assign them to a treatment arm and observe their outcomes immediately
  - ... seek to *adapt* treatment assignments for later units, based on outcomes observed in previous units, to improve estimation
- **Simple example:** testing many candidate redesigns for a website ("A/B/C/D..." test)
  - assign website visitors to a different UI as they arrive
  - measure time spent on site

Setting

- Consider a **multiarmed adaptive experiment**. This means:
  - ... have $K > 1$ treatments we want to compare vs. a control
  - ... units arrive one at a time. We assign them to a treatment arm and observe their outcomes immediately
  - ... seek to *adapt* treatment assignments for later units, based on outcomes observed in previous units, to improve estimation
- **Simple example:** testing many candidate redesigns for a website ("A/B/C/D..." test)
  - assign website visitors to a different UI as they arrive
  - measure time spent on site
- **Goal:** Obtain best estimates of *all* treatment effects (vs. the control) at the end of the trial

Building Blocks

Our approach builds on two classical statistical ideas:

Building Blocks

Our approach builds on two classical statistical ideas:


1. Neyman allocation
2. Shrinkage estimation

Idea #1: Neyman Allocation

Neyman Allocation (Neyman, 1934) gives how to choose optimal (expected-squared-error-minimizing) sample sizes for surveys

Idea #1: Neyman Allocation


Neyman Allocation (Neyman, 1934) gives how to choose optimal (expected-squared-error-minimizing) sample sizes for surveys



Problem: estimate avg. height of residents within each of Iowa's four congressional districts.

Idea #1: Neyman Allocation

Neyman Allocation (Neyman, 1934) gives how to choose optimal (expected-squared-error-minimizing) sample sizes for surveys




Problem: estimate avg. height of residents within each of Iowa's four congressional districts.

Can only sample n people. Should we sample districts proportionally to...

Idea #1: Neyman Allocation

Neyman Allocation (Neyman, 1934) gives how to choose optimal (expected-squared-error-minimizing) sample sizes for surveys




Problem: estimate avg. height of residents within each of Iowa's four congressional districts.

Can only sample n people. Should we sample districts proportionally to...

1. 1 (i.e. $n/4$ per district)?
2. # of people in each district?
3. $\mathbb{E}$(height) in each district?
4. var(height) in each district?
5. sd(height) in each district?

Idea #1: Neyman Allocation

Neyman Allocation (Neyman, 1934) gives how to choose optimal (expected-squared-error-minimizing) sample sizes for surveys



Problem: estimate avg. height of residents within each of Iowa's four congressional districts.

Can only sample n people. Should we sample districts proportionally to...

1. 1 (i.e. $n/4$ per district)?
2. # of people in each district?
3. $\mathbb{E}$(height) in each district?
4. var(height) in each district?
5. sd(height) in each district?

Idea #1: Neyman Allocation

- Neyman Allocation formalizes a simple idea: draw more samples from noisier strata

Idea #1: Neyman Allocation

- Neyman Allocation formalizes a simple idea: draw more samples from noisier strata
- In multiarmed trials, analogous idea is to **oversample treatments** that induce outcomes with **larger standard deviations**

Idea #1: Neyman Allocation

- Neyman Allocation formalizes a simple idea: draw more samples from noisier strata
- In multiarmed trials, analogous idea is to **oversample treatments** that induce outcomes with **larger standard deviations**
- Idea can be extended to **adaptive designs**

Idea #1: Neyman Allocation

- Neyman Allocation formalizes a simple idea: draw more samples from noisier strata
- In multiarmed trials, analogous idea is to **oversample treatments** that induce outcomes with **larger standard deviations**
- Idea can be extended to **adaptive designs**
  - Learn standard deviations in early samples

Idea #1: Neyman Allocation

- Neyman Allocation formalizes a simple idea: draw more samples from noisier strata
- In multiarmed trials, analogous idea is to **oversample treatments** that induce outcomes with **larger standard deviations**
- Idea can be extended to **adaptive designs**
  - Learn standard deviations in early samples
  - Later, allocate extra units to treatments that yield high-standard-deviation outcomes

Idea #1: Neyman Allocation

- Neyman Allocation formalizes a simple idea: draw more samples from noisier strata
- In multiarmed trials, analogous idea is to **oversample treatments** that induce outcomes with **larger standard deviations**
- Idea can be extended to **adaptive designs**
  - Learn standard deviations in early samples
  - Later, allocate extra units to treatments that yield high-standard-deviation outcomes
  - Variants of this approach appear in Blackwell et al. (2023), Dai et al. (2023), Zhao (2023), Li and Owen (2024) and Cai and Rafi (2024)

Idea #2: Shrinkage Estimation

Idea #2: Shrinkage Estimation

- James-Stein estimator (Stein, 1956)
  - Suppose $\boldsymbol{X} \sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2 \boldsymbol{I}_K) \in \mathbb{R}^K$, and we want to estimate $\boldsymbol{\mu}$.

Idea #2: Shrinkage Estimation

- James-Stein estimator (Stein, 1956)
  - Suppose $\boldsymbol{X} \sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2 \boldsymbol{I}_K) \in \mathbb{R}^K$, and we want to estimate $\boldsymbol{\mu}$.
  - Define $\text{JS}(\boldsymbol{X}) = \left(1 - \frac{(K-2)\sigma^2}{||\boldsymbol{X}||_2^2}\right) \boldsymbol{X}$.

Idea #2: Shrinkage Estimation

- James-Stein estimator (Stein, 1956)
  - Suppose $\boldsymbol{X} \sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2 \boldsymbol{I}_K) \in \mathbb{R}^K$, and we want to estimate $\boldsymbol{\mu}$.
  - Define $\text{JS}(\boldsymbol{X}) = \left(1 - \frac{(K-2)\sigma^2}{||\boldsymbol{X}||_2^2}\right) \boldsymbol{X}$.
  - If $K \geq 3$, $\text{JS}(\boldsymbol{X})$ has *lower expected total error* estimating $\boldsymbol{\mu}$ than does $\boldsymbol{X}$

Idea #2: Shrinkage Estimation


- James-Stein estimator (Stein, 1956)
  - Suppose $\boldsymbol{X} \sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2 \boldsymbol{I}_K) \in \mathbb{R}^K$, and we want to estimate $\boldsymbol{\mu}$.
  - Define $\text{JS}(\boldsymbol{X}) = \left(1 - \frac{(K-2)\sigma^2}{||\boldsymbol{X}||_2^2}\right) \boldsymbol{X}$.
  - If $K \geq 3$, $\text{JS}(\boldsymbol{X})$ has *lower expected total error* estimating $\boldsymbol{\mu}$ than does $\boldsymbol{X}$
- JS estimator gave birth to a whole literature on shrinkage estimators!

Idea #2: Shrinkage Estimation

- James-Stein estimator (Stein, 1956)
  - Suppose $\boldsymbol{X} \sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2 \boldsymbol{I}_K) \in \mathbb{R}^K$, and we want to estimate $\boldsymbol{\mu}$.
  - Define $\text{JS}(\boldsymbol{X}) = \left(1 - \frac{(K-2)\sigma^2}{||\boldsymbol{X}||_2^2}\right) \boldsymbol{X}$.
  - If $K \geq 3$, $\text{JS}(\boldsymbol{X})$ has *lower expected total error* estimating $\boldsymbol{\mu}$ than does $\boldsymbol{X}$
- JS estimator gave birth to a whole literature on shrinkage estimators!
- **Key Idea:** When estimating multiple quantities, use data to learn how much to "shrink" estimates toward 0.

Idea #2: Shrinkage Estimation

- James-Stein estimator (Stein, 1956)
  - Suppose $\boldsymbol{X} \sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2 \boldsymbol{I}_K) \in \mathbb{R}^K$, and we want to estimate $\boldsymbol{\mu}$.
  - Define $\text{JS}(\boldsymbol{X}) = \left(1 - \frac{(K-2)\sigma^2}{||\boldsymbol{X}||_2^2}\right) \boldsymbol{X}$.
  - If $K \geq 3$, $\text{JS}(\boldsymbol{X})$ has *lower expected total error* estimating $\boldsymbol{\mu}$ than does $\boldsymbol{X}$
- JS estimator gave birth to a whole literature on shrinkage estimators!
- **Key Idea:** When estimating multiple quantities, use data to learn how much to "shrink" estimates toward 0.



Idea #2: Shrinkage Estimation

Recent work uses shrinkage estimation in **causal inference**:

- Synthesizing observational and experimental causal estimates (Chen et al., 2015; Rosenman et al., 2023; Lin et al., 2025)
- Improved precision in online experiments (Dimmery et al., 2019; Deng et al., 2021)

Our Approach

We propose using shrinkage estimation with adaptive designs.

Our Approach

We propose using shrinkage estimation with adaptive designs.

1. Suggest a set of **shrinkage estimators** to obtain stabilized estimates of treatment effects in multiarmed trials

Our Approach

We propose using shrinkage estimation with adaptive designs.

1. Suggest a set of **shrinkage estimators** to obtain stabilized estimates of treatment effects in multiarmed trials
2. Develop a scheme to **adapt** later treatment assignments to minimize the estimated statistical error of these estimators

Our Approach

We propose using shrinkage estimation with adaptive designs.

1. Suggest a set of **shrinkage estimators** to obtain stabilized estimates of treatment effects in multiarmed trials
2. Develop a scheme to **adapt** later treatment assignments to minimize the estimated statistical error of these estimators

Simulations demonstrate significant performance improvements, especially in low signal-to-noise ratio regime!

Outline

1 Background and Motivation

2 Estimators and Risk

3 Oracle Designs

4 Adaptation & Inference

Assumptions and Set-Up

- Units $i = 1, 2, \ldots, n$ in trial are sampled iid

Assumptions and Set-Up

- Units $i = 1, 2, \ldots, n$ in trial are sampled iid
  - Each unit has $K + 1$ associated potential outcomes:

$$\underbrace{(Y_i(0), Y_i(1), \ldots, Y_i(K))}_{\substack{\text{unit's outcome under control arm } (k = 0) \\ \text{or under treatment arm } k = 1, 2, \ldots K}}$$

Assumptions and Set-Up

- Units $i = 1, 2, \ldots, n$ in trial are sampled iid
  - Each unit has $K + 1$ associated potential outcomes:

$$\underbrace{(Y_i(0), Y_i(1), \ldots, Y_i(K))}_{\substack{\text{unit's outcome under control arm } (k = 0) \\ \text{or under treatment arm } k = 1, 2, \ldots K}}$$

  - Define $\mu_k = \mathbb{E}(Y(k))$ and $V_k = \mathrm{var}(Y(k))$.

Assumptions and Set-Up

- Units $i = 1, 2, \ldots, n$ in trial are sampled iid
  - Each unit has $K + 1$ associated potential outcomes:

$$\underbrace{(Y_i(0), Y_i(1), \ldots, Y_i(K))}_{\substack{\text{unit's outcome under control arm } (k = 0) \\ \text{or under treatment arm } k = 1, 2, \ldots K}}$$

  - Define $\mu_k = \mathbb{E}(Y(k))$ and $V_k = \mathrm{var}(Y(k))$.
- **SUTVA:** units receive treatment $W_i \in \{0, 1, \ldots, K\}$, and observed outcomes are $Y_i = Y_i(W_i)$.

Assumptions and Set-Up

- Units $i = 1, 2, \ldots, n$ in trial are sampled iid
  - Each unit has $K + 1$ associated potential outcomes:

$$\underbrace{(Y_i(0), Y_i(1), \ldots, Y_i(K))}_{\substack{\text{unit's outcome under control arm } (k=0) \\ \text{or under treatment arm } k = 1, 2, \ldots K}}$$

  - Define $\mu_k = \mathbb{E}(Y(k))$ and $V_k = \mathrm{var}(Y(k))$.
- **SUTVA:** units receive treatment $W_i \in \{0, 1, \ldots, K\}$, and observed outcomes are $Y_i = Y_i(W_i)$.
- Seek to estimate vector of causal effects $\boldsymbol{\tau}$, where

$$\begin{aligned} \boldsymbol{\tau} &= (\tau_1, \ldots, \tau_K) \\ &= (\mu_1 - \mu_0, \ldots, \mu_K - \mu_0) \in \mathbb{R}^K \end{aligned}$$

Difference-in-Means Estimator

Let's first consider the *non-adaptive RCT* case.

Difference-in-Means Estimator

Let's first consider the *non-adaptive RCT* case.

- Define n_k as # of units assigned to treatment k

Difference-in-Means Estimator

Let's first consider the *non-adaptive RCT* case.

- Define n_k as # of units assigned to treatment k
- Consider the diff-in-means estimator, $\hat{\tau}$, where

$$\hat{\tau}_k = \underbrace{\frac{1}{n_k} \sum_{i=1}^{n} \mathbb{1}(W_i = k) Y_i}_{\overline{Y}_k} - \underbrace{\frac{1}{n_0} \sum_{i=1}^{n} \mathbb{1}(W_i = 0) Y_i}_{\overline{Y}_0}$$

Difference-in-Means Estimator

Let's first consider the *non-adaptive RCT* case.

- Define n_k as # of units assigned to treatment k
- Consider the diff-in-means estimator, $\hat{\boldsymbol{\tau}}$, where

$$\hat{\tau}_k = \underbrace{\frac{1}{n_k}\sum_{i=1}^{n} \mathbb{1}(W_i = k)\, Y_i}_{\overline{Y}_k} - \underbrace{\frac{1}{n_0}\sum_{i=1}^{n} \mathbb{1}(W_i = 0)\, Y_i}_{\overline{Y}_0}$$

- In a standard trial, with large n_k, approx. $\boxed{\hat{\boldsymbol{\tau}} \sim \mathcal{N}(\boldsymbol{\tau}, \boldsymbol{\Sigma})}$,

$$\boldsymbol{\Sigma} = \begin{pmatrix} \frac{V_1}{n_1} + \frac{V_0}{n_0} & \frac{V_0}{n_0} & \cdots & \frac{V_0}{n_0} \\ \frac{V_0}{n_0} & \frac{V_2}{n_2} + \frac{V_0}{n_0} & \cdots & \frac{V_0}{n_0} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{V_0}{n_0} & \frac{V_0}{n_0} & \cdots & \frac{V_K}{n_K} + \frac{V_0}{n_0} \end{pmatrix}.$$

Can we improve performance?

Note: Goal is to estimate *all* effects, not just best treatment!

Can we improve performance?

Note: Goal is to estimate *all* effects, not just best treatment!
Seek to minimize expectation of squared error loss ("risk" $\mathcal{R}$)

Can we improve performance?

Note: Goal is to estimate *all* effects, not just best treatment!
Seek to minimize expectation of squared error loss ("risk" $\mathcal{R}$)

- With diff-in-means estimator $\hat{\boldsymbol{\tau}}$,

$$\mathcal{R}_{\hat{\boldsymbol{\tau}}}\left(\boldsymbol{\tau}, \boldsymbol{\Sigma}\right) = \mathbb{E}\left(||\hat{\boldsymbol{\tau}} - \boldsymbol{\tau}||_2^2\right) = \mathrm{Tr}(\boldsymbol{\Sigma})$$

$$= \underbrace{\frac{KV_0}{n_0}}_{\substack{\text{"excess" contribution} \\ \text{of control}}} + \sum_{k=1}^{K} \frac{V_k}{n_k}.$$

Can we improve performance?

Note: Goal is to estimate *all* effects, not just best treatment!
Seek to minimize expectation of squared error loss ("risk" $\mathcal{R}$)

- With diff-in-means estimator $\hat{\boldsymbol{\tau}}$,

$$\mathcal{R}_{\hat{\boldsymbol{\tau}}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}\left(||\hat{\boldsymbol{\tau}} - \boldsymbol{\tau}||_2^2\right) = \mathrm{Tr}(\boldsymbol{\Sigma})$$

$$= \underbrace{\frac{KV_0}{n_0}}_{\substack{\text{"excess" contribution} \\ \text{of control}}} + \sum_{k=1}^{K} \frac{V_k}{n_k}.$$

- Can do better by *borrowing strength* across treatment effects

Can we improve performance?

Note: Goal is to estimate *all* effects, not just best treatment!
Seek to minimize expectation of squared error loss ("risk" $\mathcal{R}$)

- With diff-in-means estimator $\hat{\boldsymbol{\tau}}$,

$$\mathcal{R}_{\hat{\boldsymbol{\tau}}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}\left(||\hat{\boldsymbol{\tau}} - \boldsymbol{\tau}||_2^2\right) = \mathrm{Tr}(\boldsymbol{\Sigma})$$

$$= \underbrace{\frac{KV_0}{n_0}}_{\substack{\text{"excess" contribution} \\ \text{of control}}} + \sum_{k=1}^{K} \frac{V_k}{n_k}.$$

- Can do better by *borrowing strength* across treatment effects
  - Use a **shrinkage estimator** $\boldsymbol{\delta}$

Can we improve performance?

Note: Goal is to estimate *all* effects, not just best treatment!
Seek to minimize expectation of squared error loss ("risk" $\mathcal{R}$)

- With diff-in-means estimator $\hat{\boldsymbol{\tau}}$,

$$\mathcal{R}_{\hat{\boldsymbol{\tau}}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}\left(||\hat{\boldsymbol{\tau}} - \boldsymbol{\tau}||_2^2\right) = \mathrm{Tr}(\boldsymbol{\Sigma})$$

$$= \underbrace{\frac{KV_0}{n_0}}_{\substack{\text{"excess" contribution} \\ \text{of control}}} + \sum_{k=1}^{K} \frac{V_k}{n_k}.$$

- Can do better by *borrowing strength* across treatment effects
  - Use a **shrinkage estimator** $\boldsymbol{\delta}$
  - Risk $\mathcal{R}_{\boldsymbol{\delta}}(\boldsymbol{\tau}, \boldsymbol{\Sigma})$ will depend on both $\boldsymbol{\tau}$ and $\boldsymbol{\Sigma}$.

Candidate Estimators

Consider shrinkage estimators that work with *dependent* and *heteroscedastic* data.

Candidate Estimators

Consider shrinkage estimators that work with *dependent* and *heteroscedastic* data.

- Bock's estimator (Bock, 1975)

$$\boldsymbol{\delta_B} = \left(1 - \frac{\tilde{p} - 2}{\hat{\boldsymbol{\tau}}^{\mathsf{T}} \boldsymbol{\Sigma}^{-1} \hat{\boldsymbol{\tau}}}\right) \hat{\boldsymbol{\tau}} \qquad \text{where} \qquad \tilde{p} = \frac{\mathrm{Tr}(\boldsymbol{\Sigma})}{\lambda_{\max}(\boldsymbol{\Sigma})}$$

Candidate Estimators

Consider shrinkage estimators that work with *dependent* and *heteroscedastic* data.

- Bock's estimator (Bock, 1975)

$$\boldsymbol{\delta_B} = \left(1 - \frac{\tilde{p} - 2}{\hat{\boldsymbol{\tau}}^{\mathsf{T}} \boldsymbol{\Sigma}^{-1} \hat{\boldsymbol{\tau}}}\right) \hat{\boldsymbol{\tau}} \qquad \text{where} \qquad \tilde{p} = \frac{\mathrm{Tr}(\boldsymbol{\Sigma})}{\lambda_{\max}(\boldsymbol{\Sigma})}$$

- Dimmery's estimator (Dimmery et al., 2019)

$$\boldsymbol{\delta_D} = \left(\boldsymbol{I}_K - \frac{(K-2)\boldsymbol{\Sigma}_\star}{||\hat{\boldsymbol{\tau}}||_2^2}\right) \hat{\boldsymbol{\tau}}, \qquad \boldsymbol{\Sigma}_\star = \mathrm{diag}\left(\frac{V_1}{n_1} + \frac{V_0}{n_0}, \ldots, \frac{V_K}{n_K} + \frac{V_0}{n_0}\right)$$

Candidate Estimators

Consider shrinkage estimators that work with *dependent* and *heteroscedastic* data.

- Bock's estimator (Bock, 1975)

$$\boldsymbol{\delta_B} = \left(1 - \frac{\tilde{p} - 2}{\hat{\boldsymbol{\tau}}^{\mathsf{T}} \boldsymbol{\Sigma}^{-1} \hat{\boldsymbol{\tau}}}\right) \hat{\boldsymbol{\tau}} \qquad \text{where} \qquad \tilde{p} = \frac{\mathrm{Tr}(\boldsymbol{\Sigma})}{\lambda_{\max}(\boldsymbol{\Sigma})}$$

- Dimmery's estimator (Dimmery et al., 2019)

$$\boldsymbol{\delta_D} = \left(\boldsymbol{I}_K - \frac{(K-2)\boldsymbol{\Sigma}_\star}{||\hat{\boldsymbol{\tau}}||_2^2}\right) \hat{\boldsymbol{\tau}}, \qquad \boldsymbol{\Sigma}_\star = \mathrm{diag}\left(\frac{V_1}{n_1} + \frac{V_0}{n_0}, \ldots, \frac{V_K}{n_K} + \frac{V_0}{n_0}\right)$$

- SURE-minimizing estimator (Li, 1985)

$$\boldsymbol{\delta_S} = \left(1 - \frac{\mathrm{Tr}(\boldsymbol{\Sigma})}{||\hat{\boldsymbol{\tau}}||_2^2}\right) \hat{\boldsymbol{\tau}}.$$

Candidate Estimators

Consider shrinkage estimators that work with *dependent* and *heteroscedastic* data.

- Bock's estimator (Bock, 1975)

$$\delta_{\boldsymbol{B}} = \left(1 - \frac{\tilde{p} - 2}{\hat{\boldsymbol{\tau}}^{\mathsf{T}} \boldsymbol{\Sigma}^{-1} \hat{\boldsymbol{\tau}}}\right) \hat{\boldsymbol{\tau}} \qquad \text{where} \qquad \tilde{p} = \frac{\mathrm{Tr}(\boldsymbol{\Sigma})}{\lambda_{\max}(\boldsymbol{\Sigma})}$$

- Dimmery's estimator (Dimmery et al., 2019)

$$\delta_{\boldsymbol{D}} = \left(\boldsymbol{I}_K - \frac{(K-2)\boldsymbol{\Sigma}_\star}{||\hat{\boldsymbol{\tau}}||_2^2}\right) \hat{\boldsymbol{\tau}}, \qquad \boldsymbol{\Sigma}_\star = \mathrm{diag}\left(\frac{V_1}{n_1} + \frac{V_0}{n_0}, \ldots, \frac{V_K}{n_K} + \frac{V_0}{n_0}\right)$$

- SURE-minimizing estimator (Li, 1985)

$$\delta_{\boldsymbol{S}} = \left(1 - \frac{\mathrm{Tr}(\boldsymbol{\Sigma})}{||\hat{\boldsymbol{\tau}}||_2^2}\right) \hat{\boldsymbol{\tau}}.$$

Shrinker Risk

Under known parameters, a two-stage process to compute

$$\mathcal{R}_{\boldsymbol{\delta_S}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}_{\mathcal{F}}\left(||\boldsymbol{\delta_S} - \boldsymbol{\tau}||_2^2\right) = \mathbb{E}_{\mathcal{F}}\left(\sum_{k=1}^{K}(\delta_{sk} - \tau_k)^2\right).$$

Shrinker Risk

Under known parameters, a two-stage process to compute

$$\mathcal{R}_{\boldsymbol{\delta_S}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}_{\mathcal{F}}\left(||\boldsymbol{\delta_S} - \boldsymbol{\tau}||_2^2\right) = \mathbb{E}_{\mathcal{F}}\left(\sum_{k=1}^{K}(\delta_{sk} - \tau_k)^2\right).$$

1. Compute an unbiased risk estimate (modified version of SURE; Stein, 1981)

Step 1: Unbiased Risk Estimate

Corollary (Estimator Risk)

Suppose we have $\hat{\boldsymbol{\tau}} \sim \mathcal{N}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) \in \mathbb{R}^K$ *for general* $\boldsymbol{\Sigma}$. *Consider*

$$\boldsymbol{\kappa}(\hat{\boldsymbol{\tau}}) = \hat{\boldsymbol{\tau}} - \boldsymbol{g}(\hat{\boldsymbol{\tau}})$$

where $\boldsymbol{g}(\hat{\boldsymbol{\tau}})$ *is differentiable, satisfying* $\mathbb{E}(||\boldsymbol{g}||^2) < \infty$.

Step 1: Unbiased Risk Estimate

Corollary (Estimator Risk)

Suppose we have $\hat{\boldsymbol{\tau}} \sim \mathcal{N}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) \in \mathbb{R}^K$ *for general* $\boldsymbol{\Sigma}$. *Consider*

$$\boldsymbol{\kappa}(\hat{\boldsymbol{\tau}}) = \hat{\boldsymbol{\tau}} - \boldsymbol{g}(\hat{\boldsymbol{\tau}})$$

where $\boldsymbol{g}(\hat{\boldsymbol{\tau}})$ *is differentiable, satisfying* $\mathbb{E}(||\boldsymbol{g}||^2) < \infty$. *Then*

$$\begin{aligned}\mathcal{R}_{\boldsymbol{\kappa}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) &= \mathbb{E}\left(||\boldsymbol{\tau} - \boldsymbol{\kappa}(\hat{\boldsymbol{\tau}})||_2^2\right) \\ &= Tr(\boldsymbol{\Sigma}) + \mathbb{E}\Big(||\boldsymbol{g}(\hat{\boldsymbol{\tau}})||_2^2 - 2 \cdot Tr(\boldsymbol{\Sigma}\boldsymbol{\mathcal{J}}_{\boldsymbol{g}}(\hat{\boldsymbol{\tau}}))\Big)\end{aligned}$$

where $\boldsymbol{\mathcal{J}}_{\boldsymbol{g}}(\hat{\boldsymbol{\tau}})$ *is the Jacobian of* $\boldsymbol{g}(\cdot)$ *evaluated at* $\hat{\boldsymbol{\tau}}$.

Step 1: Unbiased Risk Estimate

Corollary (Estimator Risk)

Suppose we have $\hat{\boldsymbol{\tau}} \sim \mathcal{N}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) \in \mathbb{R}^K$ *for general* $\boldsymbol{\Sigma}$. *Consider*

$$\boldsymbol{\kappa}(\hat{\boldsymbol{\tau}}) = \hat{\boldsymbol{\tau}} - \boldsymbol{g}(\hat{\boldsymbol{\tau}})$$

where $\boldsymbol{g}(\hat{\boldsymbol{\tau}})$ *is differentiable, satisfying* $\mathbb{E}(||\boldsymbol{g}||^2) < \infty$. *Then*

$$\begin{aligned}\mathcal{R}_{\boldsymbol{\kappa}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) &= \mathbb{E}\left(||\boldsymbol{\tau} - \boldsymbol{\kappa}(\hat{\boldsymbol{\tau}})||_2^2\right) \\ &= Tr(\boldsymbol{\Sigma}) + \mathbb{E}\Big(||\boldsymbol{g}(\hat{\boldsymbol{\tau}})||_2^2 - 2 \cdot Tr(\boldsymbol{\Sigma}\boldsymbol{\mathcal{J}}_{\boldsymbol{g}}(\hat{\boldsymbol{\tau}}))\Big)\end{aligned}$$

where $\boldsymbol{\mathcal{J}}_{\boldsymbol{g}}(\hat{\boldsymbol{\tau}})$ *is the Jacobian of* $\boldsymbol{g}(\cdot)$ *evaluated at* $\hat{\boldsymbol{\tau}}$.

Upshot: slight modification of classical SURE to account for heteroscedasticity and dependence across causal estimates.

Shrinker Risk

Under known parameters, a two-stage process to compute

$$\mathcal{R}_{\boldsymbol{\delta_S}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}_{\mathcal{F}}\left(||\boldsymbol{\delta_S} - \boldsymbol{\tau}||_2^2\right) = \mathbb{E}_{\mathcal{F}}\left(\sum_{k=1}^{K}(\delta_{sk} - \tau_k)^2\right).$$

1. Compute an unbiased risk estimate (modified version of SURE; Stein, 1981)

$$\mathcal{R}_{\boldsymbol{\delta_S}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathrm{Tr}(\boldsymbol{\Sigma}) + \mathrm{Tr}(\boldsymbol{\Sigma})\underbrace{\mathbb{E}\left(\frac{\hat{\boldsymbol{\tau}}^{\mathsf{T}}\left(4\boldsymbol{\Sigma} - \mathrm{Tr}(\boldsymbol{\Sigma})\boldsymbol{I}_K\right)\hat{\boldsymbol{\tau}}}{||\hat{\boldsymbol{\tau}}||_2^4}\right)}_{\substack{\text{expectation of ratio of} \\ \text{Gaussian quadratic forms}}}$$

Shrinker Risk

Under known parameters, a two-stage process to compute

$$\mathcal{R}_{\boldsymbol{\delta_S}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathbb{E}_{\mathcal{F}}\left(||\boldsymbol{\delta_S} - \boldsymbol{\tau}||_2^2\right) = \mathbb{E}_{\mathcal{F}}\left(\sum_{k=1}^{K}(\delta_{sk} - \tau_k)^2\right).$$

1. Compute an unbiased risk estimate (modified version of SURE; Stein, 1981)

$$\mathcal{R}_{\boldsymbol{\delta_S}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathrm{Tr}(\boldsymbol{\Sigma}) + \mathrm{Tr}(\boldsymbol{\Sigma})\underbrace{\mathbb{E}\left(\frac{\hat{\boldsymbol{\tau}}^{\mathsf{T}}\left(4\boldsymbol{\Sigma} - \mathrm{Tr}(\boldsymbol{\Sigma})\boldsymbol{I}_K\right)\hat{\boldsymbol{\tau}}}{||\hat{\boldsymbol{\tau}}||_2^4}\right)}_{\substack{\text{expectation of ratio of} \\ \text{Gaussian quadratic forms}}}$$

2. Implement expectation as a *univariate* numerical integral via the method of Bao and Kan (2013)

Outline

1 Background and Motivation

2 Estimators and Risk

3 Oracle Designs

4 Adaptation & Inference

Oracle Setting

- With a way to compute risks of shrinkage estimators, we can optimize experimental designs to minimize these quantities

Oracle Setting

- With a way to compute risks of shrinkage estimators, we can optimize experimental designs to minimize these quantities
- To build intuition, let's consider the non-adaptive setting:

Oracle Setting

- With a way to compute risks of shrinkage estimators, we can optimize experimental designs to minimize these quantities
- To build intuition, let's consider the non-adaptive setting:
  - Values of μ_k, V_k are known for $k = 0, \ldots, K$

Oracle Setting

- With a way to compute risks of shrinkage estimators, we can optimize experimental designs to minimize these quantities
- To build intuition, let's consider the non-adaptive setting:
  - Values of μ_k, V_k are known for $k = 0, \ldots, K$
  - Allocate fixed sample size n across the $K + 1$ arms

Oracle Setting

- With a way to compute risks of shrinkage estimators, we can optimize experimental designs to minimize these quantities
- To build intuition, let's consider the non-adaptive setting:
  - Values of μ_k, V_k are known for $k = 0, \ldots, K$
  - Allocate fixed sample size n across the $K + 1$ arms
- Let's start by optimizing allocation over risk of $\hat{\boldsymbol{\tau}}$:

$$\mathcal{R}_{\hat{\boldsymbol{\tau}}}(\boldsymbol{\tau}, \boldsymbol{\Sigma}) = \mathrm{Tr}(\boldsymbol{\Sigma}) = \underbrace{\frac{KV_0}{n_0}}_{\substack{\text{“excess” contribution} \\ \text{of control}}} + \sum_{k=1}^{K} \frac{V_k}{n_k}.$$

(Oracle) Neyman Allocation

Risk-minimizing allocations is solution to Problem 1:

$$\begin{aligned} \min_{n_0,\ldots,n_K} \quad & \frac{KV_0}{n_0} + \sum_{k=1}^{K} \frac{V_k}{n_k} \\ \text{subject to} \quad & \sum_{k=0}^{K} n_k = n, \\ & 0 < n_0, \ldots, n_K. \end{aligned} \tag{1}$$

(Oracle) Neyman Allocation

Risk-minimizing allocations is solution to Problem 1:

$$\begin{aligned} \min_{n_0,\ldots,n_K} \quad & \frac{KV_0}{n_0} + \sum_{k=1}^{K} \frac{V_k}{n_k} \\ \text{subject to} \quad & \sum_{k=0}^{K} n_k = n, \\ & 0 < n_0, \ldots, n_K. \end{aligned} \tag{1}$$

Solution works out to be

$$n_0^\star = \frac{n\sqrt{KV_0}}{\sqrt{KV_0} + \sum_{\ell=1}^{K} \sqrt{V_\ell}}, \qquad \text{and}$$

$$n_k^\star = \frac{n\sqrt{V_k}}{\sqrt{KV_0} + \sum_{\ell=1}^{K} \sqrt{V_\ell}} \qquad \text{for } k = 1, \ldots, K.$$

What about the oracle δ_S allocation?

Risk-minimizing allocation for shrinker δ_S is solution to Problem 2:

$$\begin{aligned} \min_{n_0,\ldots,n_K} \quad & \mathrm{Tr}(\boldsymbol{\Sigma}) + \mathrm{Tr}(\boldsymbol{\Sigma})\mathbb{E}\left(\frac{\hat{\boldsymbol{\tau}}^{\mathsf{T}}\left(4\boldsymbol{\Sigma} - \mathrm{Tr}(\boldsymbol{\Sigma})\boldsymbol{I}_K\right)\hat{\boldsymbol{\tau}}}{||\hat{\boldsymbol{\tau}}||_2^4}\right) \\ \text{subject to} \quad & \sum_{k=0}^{K} n_k = n, \\ & 0 < n_0, \ldots, n_K. \end{aligned} \tag{2}$$

What about the oracle δ_S allocation?

Risk-minimizing allocation for shrinker δ_S is solution to Problem 2:

$$\begin{aligned} \min_{n_0,\ldots,n_K} \quad & \mathrm{Tr}(\boldsymbol{\Sigma}) + \mathrm{Tr}(\boldsymbol{\Sigma})\mathbb{E}\left(\frac{\hat{\boldsymbol{\tau}}^{\mathsf{T}}\left(4\boldsymbol{\Sigma} - \mathrm{Tr}(\boldsymbol{\Sigma})\boldsymbol{I}_K\right)\hat{\boldsymbol{\tau}}}{||\hat{\boldsymbol{\tau}}||_2^4}\right) \\ \text{subject to} \quad & \sum_{k=0}^{K} n_k = n, \\ & 0 < n_0, \ldots, n_K. \end{aligned} \tag{2}$$

- Not convex $\Longrightarrow$ hard to solve globally!


What about the oracle δ_S allocation?

Risk-minimizing allocation for shrinker δ_S is solution to Problem 2:


$$\begin{aligned} \min_{n_0,\ldots,n_K} \quad & \mathrm{Tr}(\boldsymbol{\Sigma}) + \mathrm{Tr}(\boldsymbol{\Sigma})\mathbb{E}\left(\frac{\hat{\boldsymbol{\tau}}^{\mathsf{T}}\left(4\boldsymbol{\Sigma} - \mathrm{Tr}(\boldsymbol{\Sigma})\boldsymbol{I}_K\right)\hat{\boldsymbol{\tau}}}{||\hat{\boldsymbol{\tau}}||_2^4}\right) \\ \text{subject to} \quad & \sum_{k=0}^{K} n_k = n, \\ & 0 < n_0, \ldots, n_K. \end{aligned} \tag{2}$$

- Not convex $\Longrightarrow$ hard to solve globally!
- **Greedy minimization**: start at Neyman allocation; swap units across arms until no further improvements possible

Simulated Results: Low Signal-to-Noise Ratio (SNR)




Sparse signal; low SNR




Dense signal; low SNR

- Simulated allocations for $K = 6$
- Heteroscedastic, $n = 1,000$

Simulated Results: Low Signal-to-Noise Ratio (SNR)




Sparse signal; low SNR




Dense signal; low SNR

- Simulated allocations for $K = 6$
- Heteroscedastic, $n = 1,000$
- Shrinkers allocate more units to control than Neyman
- Sparse signal: $\boldsymbol{\delta_D}, \boldsymbol{\delta_S}$ reallocate more units to strong-effect arm

Simulated Results: High SNR



Sparse signal; high SNR



Dense signal; high SNR

- Stronger signal $\rightarrow$ less reallocation to control for $\boldsymbol{\delta_B}$
- More oversampling of high-signal arm by $\boldsymbol{\delta_D}, \boldsymbol{\delta_S}$

Outline

1 Background and Motivation

2 Estimators and Risk

3 Oracle Designs

4 Adaptation & Inference

Adaptivity: Key Ideas

Upshot: With online estimates of $\boldsymbol{\Sigma}$ and $\boldsymbol{\tau}$, can estimate $\mathcal{R}_{\boldsymbol{\delta}}(\boldsymbol{\tau}, \boldsymbol{\Sigma})$ for desired shrinker $\boldsymbol{\delta}$

$\longrightarrow$ adapt future assignments to minimize estimated risk

Adaptivity: Key Ideas

Upshot: With online estimates of $\boldsymbol{\Sigma}$ and $\boldsymbol{\tau}$, can estimate $\mathcal{R}_{\boldsymbol{\delta}}(\boldsymbol{\tau}, \boldsymbol{\Sigma})$ for desired shrinker $\boldsymbol{\delta}$

$\longrightarrow$ adapt future assignments to minimize estimated risk

1. **Greedy algorithm:** for each new arrival i, choose whichever arm $k = 0, 1, \ldots, K$ minimizes current risk estimate.
2. **Batch update:** minimize treatment probabilities for each batch over current risk estimate. Apply IPW weights.
3. **Gradient update:** iterative update of treatment probabilities via risk gradient step, a la Dai et al. (2023). IPW weights.
4. ...

Adaptivity: Key Ideas

Upshot: With online estimates of $\boldsymbol{\Sigma}$ and $\boldsymbol{\tau}$, can estimate $\mathcal{R}_{\boldsymbol{\delta}}(\boldsymbol{\tau}, \boldsymbol{\Sigma})$ for desired shrinker $\boldsymbol{\delta}$

$\longrightarrow$ adapt future assignments to minimize estimated risk

1. **Greedy algorithm:** for each new arrival i, choose whichever arm $k = 0, 1, \ldots, K$ minimizes current risk estimate.
2. **Batch update:** minimize treatment probabilities for each batch over current risk estimate. Apply IPW weights.
3. **Gradient update:** iterative update of treatment probabilities via risk gradient step, a la Dai et al. (2023). IPW weights.
4. ...


Rcpp Implementation

To achieve required speeds, need extremely fast risk computation


$\longrightarrow$ use Rcpp!

```
81  // [[Rcpp::export]]
82  double calc_stein_min_risk(const arma::vec& tau_hat, const arma::mat& Sigma_hat) {
83    int K = tau_hat.n_elem;
84
85    // Compute xi
86    mat Sigma_sqrt_inv = inv(sqrtmat_sympd(Sigma_hat));
87    vec xi = Sigma_sqrt_inv * tau_hat;
88
89    // Compute matrices
90    mat A = trans(Sigma_hat) * Sigma_hat;
91    mat B = Sigma_hat;
92    double trSigma = trace(Sigma_hat);
93
94    // Eigen decomposition
95    cx_vec eigval;
96    cx_mat eigvec;
97    eig_gen(eigval, eigvec, B);
98    vec D = real(eigval);
99    mat V = real(eigvec);
100
101   // Perform integration
102   Integrand_delta_s integrand(trSigma, A, B, V, xi, D, K);
103   double err_est;
104   int err_code;
105   double integral = integrate(integrand, 0.0, std::numeric_limits<double>::infinity(), err_est, err_code);
106
107   return (1.0 / K) * (trSigma + trSigma * integral);
108 }
```

Simulated Results: δ_S at Low SNR




Sparse signal; low SNR




Dense signal; low SNR

- $K = 6$, heteroscedastic
- 2,800 units; first 100 assigned randomly
- Risk trajectories averaged over 1,000 simulations

Simulated Results: δ_S at Low SNR



Sparse signal; low SNR




Dense signal; low SNR


- $K = 6$, heteroscedastic
- 2,800 units; first 100 assigned randomly
- Risk trajectories averaged over 1,000 simulations

- Big gains from using shrinkage estimators
- Adaptivity benefits over Neyman allocation bigger when signal is sparse (i.e. one large effect)

Simulated Results: δ_S at High SNR



Sparse signal; high SNR



Dense signal; high SNR

- At high SNR, shrinkage & adaptivity benefits remain for sparse signals
- For dense signals, benefits are more limited

What about inference?

- Shrinkage & adaptivity have exclusively sought to reduce risk for point estimation.

What about inference?

- Shrinkage & adaptivity have exclusively sought to reduce risk for point estimation.
- But what about CIs: can we also obtain shorter intervals?

What about inference?

- Shrinkage & adaptivity have exclusively sought to reduce risk for point estimation.
- But what about CIs: can we also obtain shorter intervals?
- Yes!

What about inference?

- Shrinkage & adaptivity have exclusively sought to reduce risk for point estimation.
- But what about CIs: can we also obtain shorter intervals?
- Yes!
  - Adapt "Frequentist Assisted by Bayes" (FAB) intervals from Yu and Hoff (2018)

What about inference?

- Shrinkage & adaptivity have exclusively sought to reduce risk for point estimation.
- But what about CIs: can we also obtain shorter intervals?
- Yes!
  - Adapt "Frequentist Assisted by Bayes" (FAB) intervals from Yu and Hoff (2018)
  - Shorter CIs for the causal effects nonetheless maintain frequentist coverage

What about inference?

- Shrinkage & adaptivity have exclusively sought to reduce risk for point estimation.
- But what about CIs: can we also obtain shorter intervals?
- Yes!
  - Adapt "Frequentist Assisted by Bayes" (FAB) intervals from Yu and Hoff (2018)
  - Shorter CIs for the causal effects nonetheless maintain frequentist coverage
  - Intervals account for shrinkage, but are not always-valid nor corrected for adaptivity

Shorter Confidence Intervals

Adapt "FAB" intervals (Yu and Hoff, 2018) to build shorter CIs for the causal effects that nonetheless maintain frequentist coverage.

Shorter Confidence Intervals

Adapt "FAB" intervals (Yu and Hoff, 2018) to build shorter CIs for the causal effects that nonetheless maintain frequentist coverage.

SNR	τ setting	Allocation Scheme	Min Wald Coverage	Min FAB Coverage	Avg. FAB length Δ
$\kappa = 0$	–	SURE-min	95%	95%	−15%
		Dimmery	95%	95%	−15%
		Bock	94%	94%	−15%
$\kappa = 4$	sparse	SURE-min	95%	95%	−14%
		Dimmery	95%	94%	−14%
		Bock	93%	93%	−14%
$\kappa = 4$	dense	SURE-min	95%	94%	−12%
		Dimmery	96%	95%	−13%
		Bock	94%	93%	−12%

Δ is mean FAB interval length divided by mean Wald interval length, minus one.

Takeaways

- Shrinkage estimators **improve efficiency** in multiarmed trials.

Takeaways

- Shrinkage estimators **improve efficiency** in multiarmed trials.
- We propose **adaptive experimental designs** that directly minimize the estimated risk of shrinkage estimators.

Takeaways

- Shrinkage estimators **improve efficiency** in multiarmed trials.
- We propose **adaptive experimental designs** that directly minimize the estimated risk of shrinkage estimators.
  - Efficient risk computation via SURE, Laplace transforms, `Rcpp`.

Takeaways

- Shrinkage estimators **improve efficiency** in multiarmed trials.
- We propose **adaptive experimental designs** that directly minimize the estimated risk of shrinkage estimators.
  - Efficient risk computation via SURE, Laplace transforms, `Rcpp`.
  - Plays a role analogous to Neyman allocation for difference-in-means estimator.

Takeaways

- Shrinkage estimators **improve efficiency** in multiarmed trials.
- We propose **adaptive experimental designs** that directly minimize the estimated risk of shrinkage estimators.
  - Efficient risk computation via SURE, Laplace transforms, `Rcpp`.
  - Plays a role analogous to Neyman allocation for difference-in-means estimator.
- Risk-minimizing designs consistently allocate more units to the control arm.

Takeaways

- Shrinkage estimators **improve efficiency** in multiarmed trials.
- We propose **adaptive experimental designs** that directly minimize the estimated risk of shrinkage estimators.
  - Efficient risk computation via SURE, Laplace transforms, `Rcpp`.
  - Plays a role analogous to Neyman allocation for difference-in-means estimator.
- Risk-minimizing designs consistently allocate more units to the control arm.
- The largest gains occur in the low signal-to-noise regime, where shrinkage is most effective.

Thanks!





Figure 1: Evan T. R. Rosenman (CMC) & Kristen B. Hunter (UNSW)

arXiv: 2602.07404

arXiv draft

References (I)

Bao, Y. and Kan, R. (2013). On the moments of ratios of quadratic forms in normal random variables. *Journal of Multivariate Analysis*, 117:229–245.

Blackwell, M., Pashley, N. E., and Valentino, D. (2023). Batch adaptive designs to improve efficiency in social science experiments. Working paper, Harvard University.

Bock, M. E. (1975). Minimax estimators of the mean of a multivariate normal distribution. *The Annals of Statistics*, pages 209–218.

Cai, Y. and Rafi, A. (2024). On the performance of the neyman allocation with small pilots. *Journal of Econometrics*, 242(1):105793.

Chen, A., Owen, A. B., and Shi, M. (2015). Data enriched linear regression. *Electronic Journal of Statistics*, 9(1):1078–1112.

Dai, J., Gradu, P., and Harshaw, C. (2023). Clip-ogd: An experimental design for adaptive neyman allocation in sequential experiments. *Advances in Neural Information Processing Systems*, 36:32235–32269.

Deng, A., Li, Y., Lu, J., and Ramamurthy, V. (2021). On post-selection inference in a/b testing. In *Proceedings of the 27th ACM SIGKDD Conference on Knowledge Discovery & Data Mining*, pages 2743–2752.

Dimmery, D., Bakshy, E., and Sekhon, J. (2019). Shrinkage estimators in online experiments. In *Proceedings of the 25th ACM SIGKDD international conference on knowledge discovery & data mining*, pages 2914–2922.

References (II)

Li, H. H. and Owen, A. B. (2024). Double machine learning and design in batch adaptive experiments. *Journal of Causal Inference*, 12(1):20230068.

Li, K.-C. (1985). From Stein's unbiased risk estimates to the method of generalized cross validation. *The Annals of Statistics*, 13(4):1352–1377.

Lin, X., Tarp, J. M., and Evans, R. J. (2025). Combining experimental and observational data through a power likelihood. *Biometrics*, 81(1):ujaf008.

Neyman, J. (1934). On the two different aspects of the representative method: The method of stratified sampling and the method of purposive selection. *Journal of the Royal Statistical Society*, 97(4):558–606.

Pratt, J. W. (1963). Shorter confidence intervals for the mean of a normal distribution with known variance. *The Annals of Mathematical Statistics*, 34(2):574 – 586.

Rosenman, E. T., Basse, G., Owen, A. B., and Baiocchi, M. (2023). Combining observational and experimental datasets using shrinkage estimators. *Biometrics*, 79(4):2961–2973.

Stein, C. (1956). Inadmissibility of the usual estimator for the mean of a multivariate normal distribution. Technical report, Stanford University, Stanford, CA.

Stein, C. M. (1981). Estimation of the mean of a multivariate normal distribution. *The Annals of Statistics*, 9:1135–1151.

Yu, C. and Hoff, P. D. (2018). Adaptive multigroup confidence intervals with constant coverage. *Biometrika*, 105(2):319–335.

Zhao, J. (2023). Adaptive neyman allocation. *arXiv preprint arXiv:2309.08808*.

Why use δ_S?

Lemma

δ_S *dominates* $\hat{\tau}$ *if*

$$4 \cdot \lambda_{max}(\mathbf{\Sigma}) < Tr(\mathbf{\Sigma}).$$

Why use δ_S?

Lemma

δ_S dominates $\hat{\tau}$ if

$$4 \cdot \lambda_{max}(\mathbf{\Sigma}) < Tr(\mathbf{\Sigma}).$$

- Testable condition under which estimator has provably lower risk than $\hat{\tau}$ (requires at least $K > 4$)
- Estimator frequently outperforms $\hat{\tau}$ even if condition not met

FAB Intervals: Pratt's results

- Pratt (1963) considered hierarchical model:

$$Y \sim \mathcal{N}(\theta, \sigma^2) \qquad \text{where} \qquad \theta \sim \mathcal{N}(m, \eta^2)$$

- Results depend on idea of constructing *asymmetric intervals*:

$$C_w(y) = \left\{\theta : y + \sigma z_{\alpha(1-w(\theta))} < \theta < y + \sigma z_{1-\alpha w(\theta)}\right\}$$

  - $z_q = \Phi^{-1}(q)$. For any fixed w, coverage is $1 - \alpha$
  - $w \equiv 1/2$ gives the usual Wald interval.

- Under prior $\theta \sim \mathcal{N}(m, \eta^2)$, the optimal rule is

$$w(\theta) = g^{-1}\left(\frac{2\sigma(\theta - m)}{\eta^2}\right), \qquad g(w) = z_{\alpha w} - z_{\alpha(1-w)}.$$

- **Upshot:** rule produces intervals that are shorter near prior's center and longer in the tails. Intervals have $1 - \alpha$ coverage, but are shorter averaged over prior!

FAB Intervals: Our Adaptation

- **Yu & Hoff** (Yu and Hoff, 2018): proposed to *learn* the prior from other groups in the multigroup setting
- Implied prior in our setting (for $\boldsymbol{\delta_B}, \boldsymbol{\delta_S}$) is

$$\boldsymbol{\tau} \sim \mathcal{N}(\mathbf{0}, a\boldsymbol{\Sigma}), \qquad \hat{\boldsymbol{\tau}} \mid \boldsymbol{\tau} \sim \mathcal{N}(\boldsymbol{\tau}, \boldsymbol{\Sigma}), \qquad \mathbb{E}(\boldsymbol{\tau} \mid \hat{\boldsymbol{\tau}}) = \frac{a}{a+1}\hat{\boldsymbol{\tau}}.$$

- **Borrow strength across arms** to estimate the prior scale:

$$\hat{a} = \left(\frac{\hat{\boldsymbol{\tau}}^{\mathsf{T}} \hat{\boldsymbol{\Sigma}}^{-1} \hat{\boldsymbol{\tau}}}{K} - 1 \right)_{+}, \qquad \hat{\sigma}_k^2 = \hat{\boldsymbol{\Sigma}}_{kk} = \frac{\hat{V}_0}{n_0} + \frac{\hat{V}_k}{n_k}.$$

- Can then use updated FAB rule using prior variance $a\sigma_k^2$:

$$\hat{w}_k(t) = g^{-1}\left(\frac{2t}{\hat{a}\,\hat{\sigma}_k} \right).$$

- Solve two scalar equations to obtain the interval endpoints.