

(Actual) Shortest Paths

Consider the following:

$$\text{SP} \equiv \text{AddZero}(\infty, (\mathbb{R}^+, \min, +) \overrightarrow{\times} (\mathcal{P}(E^*), \cup, \hat{\cdot}))$$

we combine paths by adding their weights and concatenating their edges (thus, second component can be seen as set of paths). We choose by using min; if both have the same weight, we take the union the path sets.

| ref | S | \oplus | \otimes | $\bar{0}$ | $\bar{1}$ |
|-----|--|--|---|----------------------|---------------------------------|
| (1) | \mathbb{R}^+ | \min | $+$ | | 0 |
| (2) | $\mathcal{P}(E^*)$ | \cup | $\hat{\cdot}$ | \emptyset | $\{\epsilon\}$ |
| (3) | $\mathbb{R}^+ \times \mathcal{P}(E^*)$ | $\min \overrightarrow{\times} \cup$ | $+\times \hat{\cdot}$ | | $(0, \{\epsilon\})$ |
| SP | $\{\infty\} \uplus (\mathbb{R}^+ \times \mathcal{P}(E^*))$ | $(\min \overrightarrow{\times} \cup)_{\infty}^{\text{id}}$ | $(+\overrightarrow{\times} \hat{\cdot})_{\infty}^{\text{an}}$ | $\text{inl}(\infty)$ | $\text{inl}((0, \{\epsilon\}))$ |

We show that SP is a semiring.

First, We show that (2) is \mathbb{LD} (\mathbb{RD} is similar):

$$\begin{aligned} (A \hat{\cdot} B) \cup (A \hat{\cdot} C) &= \{a \cdot b : a \in A, b \in B\} \cup \{a \cdot c : a \in A, c \in C\} \\ &= \{a \cdot x : a \in A, x \in B \cup C\} \\ &= A \hat{\cdot} (B \cup C) \end{aligned}$$

\mathbb{LD} for (3) requires:

$$\mathbb{LD}((1) \overrightarrow{\times} (2)) \equiv \underbrace{\mathbb{LD}((1))}_{\text{yes}} \wedge \underbrace{\mathbb{LD}((2))}_{\text{yes}} \wedge \underbrace{(\mathbb{LC}(\mathbb{R}^+, +) \vee \mathbb{LK}(\mathcal{P}(E^*), \hat{\cdot}))}_{\text{yes!}}$$

\mathbb{LD} for SP follows, since AddZero preserves \mathbb{LD} .

\mathbb{RD} is similar. Thus, SP is both \mathbb{LD} and \mathbb{RD} .

We need \oplus to be commutative:

$$\mathbb{CM}((\mathbb{R}^+, \min) \overrightarrow{\times} (\mathcal{P}(E^*), \cup)) \equiv \underbrace{\mathbb{CM}(\mathcal{P}(E^*), \cup)}_{\text{yes!}}$$

We have a zero by construction, annihilating \otimes . $\text{inr}((0, \{\epsilon\}))$ is a multiplicative ID, since $((0, \{\epsilon\}))$ is an ID for $+\times \hat{\cdot}$.

It follows that SP is a semiring.

SP adjacency matrix, A for graph $G = (V, E)$ with weights $w : E \mapsto \mathbb{R}^+$:

$$A_{ij} = \begin{cases} \text{inr}((w(ij), \{ij\})) & \text{if } ij \in E \\ \text{inl}(\infty) & \text{otherwise} \end{cases}$$