

ScatteringKernelsin 3D

This is code to accompany the book:

A Hitchhiker's Guide to Multiple Scattering

© 2022 Eugene d'Eon

www.eugenedeon.com/hitchhikers

Draine

Draine, B.T. (2003) 'Scattering by interstellar dust grains. 1: Optical and ultraviolet', ApJ., 598, 1017–25.

$$\text{In[10790]:= } \text{pDraine}[u_ , g_ , \alpha_] := \frac{1}{4 \text{ Pi}} \left(\frac{1 - g^2}{(1 + g^2 - 2 g u)^{3/2}} \frac{1 + \alpha u^2}{1 + \alpha (1 + 2 g^2) / 3} \right)$$

Normalization condition

`In[*]:= Integrate[2 Pi pDraine[u, g, a], {u, -1, 1}, Assumptions → 0 < a < 1 && -1 < g < 1]`

`Out[*]= 1`

Mean-cosine

`In[*]:= Integrate[2 Pi pDraine[u, g, a] u, {u, -1, 1}, Assumptions → 0 < a < 1 && -1 < g < 1]`

$$\text{Out[*]= } \frac{3}{5} \left(g + \frac{2 (1 + a) g}{3 + a + 2 a g^2} \right)$$

$$\text{In[*]:= } \frac{3}{5} \left(g + \frac{2 (1 + a) g}{3 + a + 2 a g^2} \right) /. a \rightarrow 0$$

`Out[*]= g`

Legendre expansion coefficients

`In[*]:= Integrate[2 Pi (2 k + 1) pDraine[Cos[y], g, a] LegendreP[k, Cos[y]] Sin[y] /. k → 0, {y, 0, Pi}, Assumptions → 0 < a < 1 && -1 < g < 1]`

`Out[*]= 1`

`In[*]:= Integrate[2 Pi (2 k + 1) pDraine[Cos[y], g, a] LegendreP[k, Cos[y]] Sin[y] /. k → 1, {y, 0, Pi}, Assumptions → 0 < a < 1 && -1 < g < 1]`

$$\text{Out[*]= } \frac{9 g (5 + a (3 + 2 g^2))}{5 (3 + a + 2 a g^2)}$$

```
In[*]:= Integrate[2 Pi (2 k + 1) pDraine[Cos[y], g, a] LegendreP[k, Cos[y]] Sin[y] /. k -> 2,
  {y, 0, Pi}, Assumptions -> 0 < a < 1 && -1 < g < 1]
```

$$\text{Out[*]} = \frac{14 a + 5 (21 + 11 a) g^2 + 36 a g^4}{7 (3 + a + 2 a g^2)}$$

```
In[*]:= Integrate[2 Pi (2 k + 1) pDraine[Cos[y], g, a] LegendreP[k, Cos[y]] Sin[y] /. k -> 3,
  {y, 0, Pi}, Assumptions -> 0 < a < 1 && -1 < g < 1]
```

$$\text{Out[*]} = \frac{g (54 a + 7 (45 + 23 a) g^2 + 100 a g^4)}{15 (3 + a + 2 a g^2)}$$

sampling

```
In[*]:= cdf = Integrate[2 Pi pDraine[u, g, a],
  {u, -1, x}, Assumptions -> 0 < a < 1 && -1 < g < 1 && -1 < x < 1]
```

$$\begin{aligned} \text{Out[*]} = & \left(3 (-1 + g) g^2 (-1 - g + \sqrt{1 + g^2 - 2 g x}) + \right. \\ & a \left(2 - 2 g^6 - 2 g x - 2 \sqrt{1 + g^2 - 2 g x} + g^3 \sqrt{1 + g^2 - 2 g x} + g^4 (-2 + x^2) + \right. \\ & \left. \left. 2 g^5 (x + \sqrt{1 + g^2 - 2 g x}) - g^2 (-2 + x^2 + \sqrt{1 + g^2 - 2 g x}) \right) \right) / \\ & \left(2 g^3 (3 + a + 2 a g^2) \sqrt{1 + g^2 - 2 g x} \right) \end{aligned}$$

general case

```
In[*]:= sampleDraineSimplified[xi_, g_, a_] := Module[{T1, T2, T3, T4, T4a, T5, T6},
```

$$T1 = (-1 + g^2) (4 g^2 + a (1 + g^2)^2);$$

$$T2 = -1296 (a - a g^2) (-a + a g^4) T1;$$

$$T3 = 3 g^2 (1 + g (-1 + 2 x i)) + a (2 + g^2 + g^3 (1 + 2 g^2) (-1 + 2 x i));$$

$$T4a = 432 (-a + a g^4)^3 + T2 + 432 (a - a g^2) T3^2;$$

$$T4 = T4a + \sqrt{-4 (-144 a g^2 + 288 a g^4 - 144 a g^6)^3 + (T4a)^2};$$


$$T6 = \frac{1}{(a - a g^2)} \left(2 (-a + a g^4) + \frac{48 \times 2^{1/3} (-a g^2 + 2 a g^4 - a g^6)}{T4^{1/3}} + \frac{T4^{1/3}}{3 \times 2^{1/3}} \right);$$

$$T5 = 6 (1 + g^2) + T6;$$

$$\frac{1 + g^2 - \left(\frac{1}{2} \sqrt{6 (1 + g^2)} - T6 - \frac{8 T3}{a (-1 + g^2) \sqrt{T5}} - \frac{\sqrt{T5}}{2} \right)^2}{2 g}$$

```
]
```

```
In[10999]:= With[{a = 6.7, g = .46}, Show[
  Histogram[Table[sampleDraineSimplified[RandomReal[], g, a], {i, 1, 100 000}],
    100, "PDF", ScalingFunctions -> "Log"],
  LogPlot[2 Pi pDraine[u, g, a], {u, -1, 1}, PlotRange -> All]
]
```



special case $g = 1/2$, $a = 12$ (useful for approximating Mie scattering of water spheres in air)

```
In[*]:= pDraine[μ, 1/2, 12] // FullSimplify
```

$$\text{Out[*]} = \frac{3 (1 + 12 \mu^2)}{14 \pi (5 - 4 \mu)^{3/2}}$$

Simplification of an exact CDF inverse:

```
In[*]:= sampleDraineFog[xi_] := Module[{T1, T2, T3},
  T1 =  $\sqrt{(67 + 14 xi)^4 (5239 - 102376 xi + 492072 xi^2 + 105056 xi^3 + 5488 xi^4)}$  ;
  T2 =  $(883297 - 8820952 xi + 2597784 xi^2 + 735392 xi^3 + 38416 xi^4 + \sqrt{7} T1)^{1/3}$  ;
  T3 =  $\sqrt{((3954 \times 2^{2/3} \times 3^{1/3} - 56280 \times 2^{2/3} \times 3^{1/3} xi - 5880 \times 2^{2/3} \times 3^{1/3} xi^2 + 912 T2 + 2^{1/3} \times 3^{2/3} T2^2) / T2)}$  ;
   $\frac{1}{12} (-30 - T3 + \sqrt{(1824 + (6 \times 2^{2/3} \times 3^{1/3} (-659 + 9380 xi + 980 xi^2)) / T2 - 2^{1/3} \times 3^{2/3} T2 + (12 (67 + 14 xi)^2) / (T3))})$ 
]
```

```

In[ ]:= Show[Histogram[Table[sampleDraineFog[RandomReal[]], {i, 1, 100 000}],
  100, "PDF", ScalingFunctions -> "Log"],
  LogPlot[2 Pi pDraine[u, .5, 12.], {u, -1, 1}, PlotRange -> All]

```

